(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 25 May 2001 (25.05.2001)

PCT

(10) International Publication Number WO 01/36471 A2

| (51) International Patent Classification?: | C07K 14/00 | 60/242,332 | 20 October 2000 (20.10.2000) | US |
|--|----------------|--------------------------|--|----------|
| (21) International Application Number: | PCT/US00/31509 | 60/242,343 60/243,019 | 20 October 2000 (20.10.2000) 24 October 2000 (24.10.2000) | US US |

(22) International Filing Date: 16 November 2000 (16.11.2000)

(25) Filing Language:

(30

English

(26) Publication Language: English

|) | Priority Data: | | |
|---|----------------|--------------------------------|----|
| | 60/166,088 | 17 November 1999 (17.11.1999) | US |
| | 60/166,099 | 17 November 1999 (17.11.1999) | US |
| | 60/166,369 | 17 November 1999 (17.11.1999) | US |
| | 60/171,900 | 23 December 1999 (23.12.1999) | US |
| | 60/171,901 | 23 December 1999 (23.12.1999) | US |
| | 60/171,902 | 23 December 1999 (23.12.1999) | US |
| | 60/181,749 | 11 February 2000 (11.02.2000) | US |
| | 60/189,258 | 14 March 2000 (14.03.2000) | US |
| | 60/189,259 | 14 March 2000 (14.03.2000) | US |
| | 60/195,898 | 10 April 2000 (10.04.2000) | US |
| | 60/195,899 | 10 April 2000 (10.04.2000) | US |
| | 60/196,078 | 10 April 2000 (10.04.2000) | US |
| | 60/200,419 | 28 April 2000 (28.04.2000) | US |
| | 60/203,630 | 12 May 2000 (12.05.2000) | US |
| | 60/210,741 | 12 June 2000 (12.06.2000) | US |
| | 60/210,982 | 12 June 2000 (12.06.2000) | US |
| | 60/226,760 | 21 August 2000 (21.08.2000) | US |
| | 60/235,418 | 26 September 2000 (26.09.2000) | US |
| | 60/235,779 | 26 September 2000 (26.09.2000) | US |
| | | | |

(71) Applicant (for all designated States except US): ARENA PHARMACEUTICALS, INC. [US/US]; 6166 Nancy Ridge Drive, San Diego, CA 92121 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): CHEN, Ruoping [CN/US]; 5296 Timber Branch Way, San Diego, CA 92130 (US). DANG, Huong, T. [US/US]; 5352 Oak Park Drive, San Diego, CA 92105 (US). LOWITZ, Kevin, P. [US/US]; 8031 Caminito de Pizza #C, San Diego, CA 82108 (US).

(74) Agents: MILLER, Suzanne, E. et al.; Woodcock Washburn Kurtz Mackiewicz & Norris LLP, One Liberty Place -46th Floor, Philadelphia, PA 19103 (US).


(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GII, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European

[Continued on next page]

(54) Title: ENDOGENOUS AND NON-ENDOGENOUS VERSIONS OF HUMAN G PROTEIN-COUPLED RECEPTORS

IP3 Assay in 293 Cells



(57) Abstract: The invention disclosed in this patent document relates to transmembrane receptors, more particularly to a human G protein-coupled receptor for which the endogenous ligand is unknown ("orphan GPCR receptors"), and most particularly to mutated (non-endogenous) versions of the human GPCRs for evidence of constitutive activity.



patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

 Without international search report and to be republished upon receipt of that report.

ENDOGENOUS AND NON-ENDOGENOUS VERSIONS OF HUMAN G PROTEIN-COUPLED RECEPTORS

FIELD OF THE INVENTION

5

The invention disclosed in this patent document relates to transmembrane receptors, and more particularly to human G protein-coupled receptors, and specifically to endogenous human GPCRs with particular emphasis on non-endogenous versions of the GPCRs that have been altered to establish or enhance constitutive activity of the receptor. Preferably, the altered GPCRs are used for the direct identification of candidate compounds as receptor agonists, inverse agonists or partial agonists having potential applicability as therapeutic agents.

15

20

25

10

BACKGROUND OF THE INVENTION

Although a number of receptor classes exist in humans, by far the most abundant and therapeutically relevant is represented by the G protein-coupled receptor (GPCR or GPCRs) class. It is estimated that there are some 100,000 genes within the human genome, and of these, approximately 2%, or 2,000 genes, are estimated to code for GPCRs. Receptors, including GPCRs, for which the endogenous ligand has been identified are referred to as "known" receptors, while receptors for which the endogenous ligand has not been identified are referred to as "orphan" receptors. GPCRs represent an important area for the development of pharmaceutical products: from approximately 20 of the 100 known GPCRs, approximately 60% of all prescription pharmaceuticals have been developed.

GPCRs share a common structural motif. All these receptors have seven sequences of between 22 to 24 hydrophobic amino acids that form seven alpha helices, each of which spans the membrane (each span is identified by number, *i.e.*, transmembrane-1 (TM-1), transmebrane-2 (TM-2), etc.). The transmembrane helices are joined by strands of amino acids between transmembrane-2 and transmembrane-3,

transmembrane-4 and transmembrane-5, and transmembrane-6 and transmembrane-7 on the exterior, or "extracellular" side, of the cell membrane (these are referred to as "extracellular" regions 1, 2 and 3 (EC-1, EC-2 and EC-3), respectively). The transmembrane helices are also joined by strands of amino acids between transmembrane-1 and transmembrane-2, transmembrane-3 and transmembrane-4, and transmembrane-5 and transmembrane-6 on the interior, or "intracellular" side, of the cell membrane (these are referred to as "intracellular" regions 1, 2 and 3 (IC-1, IC-2 and IC-3), respectively). The "carboxy" ("C") terminus of the receptor lies in the intracellular space within the cell, and the "amino" ("N") terminus of the receptor lies in the extracellular space outside of the cell.

Generally, when an endogenous ligand binds with the receptor (often referred to as "activation" of the receptor), there is a change in the conformation of the intracellular region that allows for coupling between the intracellular region and an intracellular "G-protein." It has been reported that GPCRs are "promiscuous" with respect to G proteins, *i.e.*, that a GPCR can interact with more than one G protein. *See*, Kenakin, T., 43 *Life Sciences* 1095 (1988). Although other G proteins exist, currently, Gq, Gs, Gi, Gz and Go are G proteins that have been identified. Endogenous ligand-activated GPCR coupling with the G-protein begins a signaling cascade process (referred to as "signal transduction"). Under normal conditions, signal transduction ultimately results in cellular activation or cellular inhibition. It is thought that the IC-3 loop as well as the carboxy terminus of the receptor interact with the G protein.

Under physiological conditions, GPCRs exist in the cell membrane in equilibrium between two different conformations: an "inactive" state and an "active" state. A receptor in an inactive state is unable to link to the intracellular signaling transduction pathway to produce a biological response. Changing the receptor

25

5

10

15

conformation to the active state allows linkage to the transduction pathway (via the G-protein) and produces a biological response.

A receptor may be stabilized in an active state by an endogenous ligand or a compound such as a drug. Recent discoveries, including but not exclusively limited to modifications to the amino acid sequence of the receptor, provide means other than endogenous ligands or drugs to promote and stabilize the receptor in the active state conformation. These means effectively stabilize the receptor in an active state by simulating the effect of an endogenous ligand binding to the receptor. Stabilization by such ligand-independent means is termed "constitutive receptor activation."

10

5

SUMMARY OF THE INVENTION

Disclosed herein are endogenous and non-endogenous versions of human GPCRs and uses thereof.

15

20

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides an illustration of second messenger IP₃ production from endogenous version RUP12 ("RUP12") as compared with the control ("CMV").

Figure 2 is a graphic representation of the results of a second messenger cell-based cyclic AMP assay providing comparative results for constitutive signaling of endogenous RUP13 ("RUP13") and a control vector ("CMV").

Figure 3 is a diagrammatic representation of the signal measured comparing CMV, endogenous RUP13 ("RUP13 wt") and non-endogenous, constitutively activated RUP13 ("RUP13(A268K)"), utilizing 8XCRE-Luc reporter plasmid.

Figure 4 is a graphic representation of the results of a [35S]GTPγS assay providing comparative results for constitutive signaling by RUP13:Gs Fusion Protein ("RUP13-Gs") and a control vector ("CMV").

Figure 5 is a diagrammatic representation of the signal measured comparing CMV, endogenous RUP14 ("RUP14 wt") and non-endogenous, constitutively activated RUP13 ("RUP14(L246K)"), utilizing 8XCRE-Luc reporter plasmid.

Figure 6 is a diagrammatic representation of the signal measured comparing CMV, endogenous RUP15 ("RUP15 wt") and non-endogenous, constitutively activated RUP15 ("RUP15(A398K)"), utilizing 8XCRE-Luc reporter plasmid.

Figure 7 is a graphic representation of the results of a second messenger cell-based cyclic AMP assay providing comparative results for constitutive signaling of endogenous RUP15 ("RUP15 wt"), non-endogenous, constitutively activated version of RUP15 ("RUP15(A398K)") and a control vector ("CMV").

Figure 8 is a graphic representation of the results of a [35]GTPγS assay providing comparative results for constitutive signaling by RUP15:Gs Fusion Protein ("RUP15-Gs") and a control vector ("CMV").

Figure 9 provides an illustration of second messenger IP₃ production from endogenous version RUP17 ("RUP17") as compared with the control ("CMV").

Figure 10 provides an illustration of second messenger IP₃ production from endogenous version RUP21 ("RUP21") as compared with the control ("CMV").

Figure 11 is a diagrammatic representation of the signal measured comparing CMV, endogenous RUP23 ("RUP23 wt") and non-endogenous, constitutively activated RUP23 ("RUP23(W275K)"), utilizing 8XCRE-Luc reporter plasmid.

5

10

15

Figure 12 is a graphic representation of results from a primary screen of several candidate compounds against RUP13; results for "Compound A" are provided in well A2 and "Compound "B" are provided in well G9.

5

10

DETAILED DESCRIPTION

The scientific literature that has evolved around receptors has adopted a number of terms to refer to ligands having various effects on receptors. For clarity and consistency, the following definitions will be used throughout this patent document. To the extent that these definitions conflict with other definitions for these terms, the following definitions shall control:

AGONISTS shall mean materials (e.g., ligands, candidate compounds) that activate the intracellular response when they bind to the receptor, or enhance GTP binding to membranes.

AMINO ACID ABBREVIATIONS used herein are set out in Table A:

TABLE A

| ALANINE | ALA | A |
|---------------|-----|---|
| ARGININE | ARG | R |
| ASPARAGINE | ASN | N |
| ASPARTIC ACID | ASP | D |
| CYSTEINE | CYS | С |
| GLUTAMIC ACID | GLU | E |
| GLUTAMINE | GLN | Q |
| GLYCINE | GLY | G |
| HISTIDINE | HIS | Н |
| ISOLEUCINE | ILE | I |
| LEUCINE | LEU | L |
| LYSINE | LYS | K |
| METHIONINE | MET | M |

| PHENYLALANINE | PHE | F |
|---------------|-----|---|
| PROLINE | PRO | P |
| SERINE | SER | S |
| THREONINE | THR | T |
| TRYPTOPHAN | TRP | W |
| TYROSINE | TYR | Y |
| VALINE | VAL | V |

PARTIAL AGONISTS shall mean materials (e.g., ligands, candidate compounds) that activate the intracellular response when they bind to the receptor to a lesser degree/extent than do agonists, or enhance GTP binding to membranes to a lesser degree/extent than do agonists.

ANTAGONIST shall mean materials (e.g., ligands, candidate compounds) that competitively bind to the receptor at the same site as the agonists but which do not activate the intracellular response initiated by the active form of the receptor, and can thereby inhibit the intracellular responses by agonists or partial agonists. ANTAGONISTS do not diminish the baseline intracellular response in the absence of an agonist or partial agonist.

CANDIDATE COMPOUND shall mean a molecule (for example, and not limitation, a chemical compound) that is amenable to a screening technique. Preferably, the phrase "candidate compound" does not include compounds which were publicly known to be compounds selected from the group consisting of inverse agonist, agonist or antagonist to a receptor, as previously determined by an indirect identification process ("indirectly identified compound"); more preferably, not including an indirectly identified compound which has previously been determined to have therapeutic efficacy in at least one mammal; and, most preferably, not including an indirectly identified compound which has previously been determined to have therapeutic utility in humans.

20

5

10

COMPOSITION means a material comprising at least one component; a "pharmaceutical composition" is an example of a composition:

COMPOUND EFFICACY shall mean a measurement of the ability of a compound to inhibit or stimulate receptor functionality, as opposed to receptor binding affinity. Exemplary means of detecting compound efficacy are disclosed in the Example section of this patent document:

CODON shall mean a grouping of three nucleotides (or equivalents to nucleotides) which generally comprise a nucleoside (adenosine (A), guanosine (G), cytidine (C), uridine (U) and thymidine (T)) coupled to a phosphate group and which, when translated, encodes an amino acid.

CONSTITUTIVELY ACTIVATED RECEPTOR shall mean a receptor subject to constitutive receptor activation. A constitutively activated receptor can be endogenous or non-endogenous.

CONSTITUTIVE RECEPTOR ACTIVATION shall mean stabilization of a receptor in the active state by means other than binding of the receptor with its endogenous ligand or a chemical equivalent thereof.

CONTACT or CONTACTING shall mean bringing at least two moieties together, whether in an in vitro system or an in vivo system.

the phrase "candidate compound", shall mean the screening of a candidate compound against a constitutively activated receptor, preferably a constitutively activated orphan receptor, and most preferably against a constitutively activated G protein-coupled cell surface orphan receptor, and assessing the compound efficacy of such compound. This phrase is, under no circumstances, to be interpreted or understood to be encompassed by or to encompass the phrase "indirectly identifying" or "indirectly identified."

5

10

ENDOGENOUS shall mean a material that a mammal naturally produces. ENDOGENOUS in reference to, for example and not limitation, the term "receptor," shall mean that which is naturally produced by a mammal (for example, and not limitation, a human) or a virus. By contrast, the term NON-ENDOGENOUS in this context shall mean that which is not naturally produced by a mammal (for example, and not limitation, a human) or a virus. For example, and not limitation, a receptor which is not constitutively active in its endogenous form, but when manipulated becomes constitutively active, is most preferably referred to herein as a "non-endogenous, constitutively activated receptor." Both terms can be utilized to describe both "in vivo" and "in vitro" systems. For example, and not limitation, in a screening approach, the endogenous or non-endogenous receptor may be in reference to an in vitro screening system. As a further example and not limitation, where the genome of a mammal has been manipulated to include a non-endogenous constitutively activated receptor, screening of a candidate compound by means of an in vivo system is viable.

FUSION PROTEIN, in the context of the invention disclosed herein, each mean a non-endogenous protein comprising an endogenous, constitutively activate GPCR or a non-endogenous, constitutively activated GPCR fused to at least one G protein, most preferably the alpha (α) subunit of such G protein (this being the subunit that binds GTP), with the G protein preferably being of the same type as the G protein that naturally couples with endogenous orphan GPCR. For example, and not limitation, in an endogenous state, if the G protein "Gs α " is the predominate G protein that couples with the GPCR, a GPCR Fusion Protein based upon the specific GPCR would be a non-endogenous protein comprising the GPCR fused to Gs α ; in some circumstances, as will be set forth below, a non-predominant G protein can be fused to the GPCR. The G

25

5

10

15

protein can be fused directly to the c-terminus of the constitutively active GPCR or there may be spacers between the two.

HOST CELL shall mean a cell capable of having a Plasmid and/or Vector incorporated therein. In the case of a prokaryotic Host Cell, a Plasmid is typically replicated as a autonomous molecule as the Host Cell replicates (generally, the Plasmid is thereafter isolated for introduction into a eukaryotic Host Cell); in the case of a eukaryotic Host Cell, a Plasmid is integrated into the cellular DNA of the Host Cell such that when the eukaryotic Host Cell replicates, the Plasmid replicates. Preferably, for the purposes of the invention disclosed herein, the Host Cell is eukaryotic, more preferably, mammalian, and most preferably selected from the group consisting of 293, 293T and COS-7 cells.

INDIRECTLY IDENTIFYING or INDIRECTLY IDENTIFIED means the traditional approach to the drug discovery process involving identification of an endogenous ligand specific for an endogenous receptor, screening of candidate compounds against the receptor for determination of those which interfere and/or compete with the ligand-receptor interaction, and assessing the efficacy of the compound for affecting at least one second messenger pathway associated with the activated receptor.

INHIBIT or INHIBITING, in relationship to the term "response" shall mean that a response is decreased or prevented in the presence of a compound as opposed to in the absence of the compound.

INVERSE AGONISTS shall mean materials (e.g., ligand, candidate compound) which bind to either the endogenous form of the receptor or to the constitutively activated form of the receptor, and which inhibit the baseline intracellular response initiated by the active form of the receptor below the normal base level of activity which

25

5

10

15

is observed in the absence of agonists or partial agonists, or decrease GTP binding to membranes. Preferably, the baseline intracellular response is inhibited in the presence of the inverse agonist by at least 30%, more preferably by at least 50%, and most preferably by at least 75%, as compared with the baseline response in the absence of the inverse agonist.

KNOWN RECEPTOR shall mean an endogenous receptor for which the endogenous ligand specific for that receptor has been identified.

LIGAND shall mean an endogenous, naturally occurring molecule specific for an endogenous, naturally occurring receptor.

MUTANT or MUTATION in reference to an endogenous receptor's nucleic acid and/or amino acid sequence shall mean a specified change or changes to such endogenous sequences such that a mutated form of an endogenous, non-constitutively activated receptor evidences constitutive activation of the receptor. In terms of equivalents to specific sequences, a subsequent mutated form of a human receptor is considered to be equivalent to a first mutation of the human receptor if (a) the level of constitutive activation of the subsequent mutated form of a human receptor is substantially the same as that evidenced by the first mutation of the receptor; and (b) the percent sequence (amino acid and/or nucleic acid) homology between the subsequent mutated form of the receptor and the first mutation of the receptor is at least about 80%, more preferably at least about 90% and most preferably at least 95%. Ideally, and owing to the fact that the most preferred cassettes disclosed herein for achieving constitutive activation includes a single amino acid and/or codon change between the endogenous and the non-endogenous forms of the GPCR, the percent sequence homology should be at least 98%.

5

10

15

NON-ORPHAN RECEPTOR shall mean an endogenous naturally occurring molecule specific for an endogenous naturally occurring ligand wherein the binding of a ligand to a receptor activates an intracellular signaling pathway.

ORPHAN RECEPTOR shall mean an endogenous receptor for which the endogenous ligand specific for that receptor has not been identified or is not known.

PHARMACEUTICAL COMPOSITION shall mean a composition comprising at least one active ingredient, whereby the composition is amenable to investigation for a specified, efficacious outcome in a mammal (for example, and not limitation, a human). Those of ordinary skill in the art will understand and appreciate the techniques appropriate for determining whether an active ingredient has a desired efficacious outcome based upon the needs of the artisan.

PLASMID shall mean the combination of a Vector and cDNA. Generally, a Plasmid is introduced into a Host Cell for the purposes of replication and/or expression of the cDNA as a protein.

SECOND MESSENGER shall mean an intracellular response produced as a result of receptor activation. A second messenger can include, for example, inositol triphosphate (IP₃), diacycglycerol (DAG), cyclic AMP (cAMP), and cyclic GMP (cGMP). Second messenger response can be measured for a determination of receptor activation. In addition, second messenger response can be measured for the direct identification of candidate compounds, including for example, inverse agonists, agonists, partial agonists and antagonists.

STIMULATE or STIMULATING, in relationship to the term "response" shall mean that a response is increased in the presence of a compound as opposed to in the absence of the compound.

5

10

15

VECTOR in reference to cDNA shall mean a circular DNA capable of incorporating at least one cDNA and capable of incorporation into a Host Cell.

The order of the following sections is set forth for presentational efficiency and is not intended, nor should be construed, as a limitation on the disclosure or the claims to follow.

A. Introduction

5

10

15

20

The traditional study of receptors has always proceeded from the a priori assumption (historically based) that the endogenous ligand must first be identified before discovery could proceed to find antagonists and other molecules that could affect the receptor. Even in cases where an antagonist might have been known first, the search immediately extended to looking for the endogenous ligand. This mode of thinking has persisted in receptor research even after the discovery of constitutively activated receptors. What has not been heretofore recognized is that it is the active state of the receptor that is most useful for discovering agonists, partial agonists, and inverse agonists of the receptor. For those diseases which result from an overly active receptor or an under-active receptor, what is desired in a therapeutic drug is a compound which acts to diminish the active state of a receptor or enhance the activity of the receptor, respectively, not necessarily a drug which is an antagonist to the endogenous ligand. This is because a compound that reduces or enhances the activity of the active receptor state need not bind at the same site as the endogenous ligand. Thus, as taught by a method of this invention, any search for therapeutic compounds should start by screening compounds against the ligand-independent active state.

B. Identification of Human GPCRs

The efforts of the Human Genome project has led to the identification of a plethora of information regarding nucleic acid sequences located within the human genome; it has been the case in this endeavor that genetic sequence information has been made available without an understanding or recognition as to whether or not any particular genomic sequence does or may contain open-reading frame information that translate human proteins. Several methods of identifying nucleic acid sequences within the human genome are within the purview of those having ordinary skill in the art. For example, and not limitation, a variety of human GPCRs, disclosed herein, were discovered by reviewing the GenBankTM database. Table B, below, lists several endogenous GPCRs that we have discovered, along with other GPCR's that are homologous to the disclosed GPCR.

TABLE B

| Disclosed Human Orphan GPCRs | Accession Number Identified | Open Reading Frame (Base Pairs) | Reference To Homologous GPCR | Per Cent Homology To Designated GPCR |
|------------------------------------|-----------------------------------|---------------------------------------|------------------------------------|--------------------------------------|
| hRUP8 | AL121755 | 1,152bp | NPY2R | 27% |
| hRUP9 | AC0113375 | 1,260bp | GAL2R | 22% |
| hRUP10 | AC008745 | 1,014bp | C5aR | 40% |
| hRUP11 | AC013396 | 1,272bp | HM74 | 36% |
| hRUP12 | AP000808 | 966bp | Masl | 34% |
| hRUP13 | AC011780 | 1,356bp | Fish GPRX- ORYLA | 43% |
| hRUP14 | AL137118 | 1,041bp | CysLT1R | 35% |
| hRUP15 | AL016468 | 1,527bp | RE2 | 30% |
| hRUP16 | AL136106 | 1,068bp | GLR101 | 37% |
| hRUP17 | AC023078 | 969bp | Masl | 37% |
| hRUP18 | AC008547 | 1,305bp | Oxytocin | 31% |
| hRUP19 | AC026331 | 1,041bp | HM74 | 52% |
| hRUP20 | AL161458 | 1,011bp | GPR34 | 25% |
| hRUP21 | AC026756 | 1,014bp | P2Y1R | 37% |
| hRUP22 | AC027026 | 993bp | RUP17 Mas1 | 67% 37% |

5

| hRUP23 | AC007104 | 1,092bp | Rat GPR26 | 31% |
|--------|----------|---------|--------------|-----|
| hRUP24 | AL355388 | 1,125bp | SALPR | 44% |
| hRUP25 | AC026331 | 1,092bp | HM74 | 95% |
| hRUP26 | AC023040 | 1,044bp | Rabbit 5HT1D | 27% |
| hRUP27 | AC027643 | 158,700 | МСН | 38% |

Receptor homology is useful in terms of gaining an appreciation of a role of the receptors within the human body. As the patent document progresses, we will disclose techniques for mutating these receptors to establish non-endogenous, constitutively activated versions of these receptors.

The techniques disclosed herein have also been applied to other human, orphan GPCRs known to the art, as will be apparent as the patent document progresses.

C. Receptor Screening

Screening candidate compounds against a non-endogenous, constitutively activated version of the human GPCRs disclosed herein allows for the direct identification of candidate compounds which act at this cell surface receptor, without requiring use of the receptor's endogenous ligand. Using routine, and often commercially available techniques, one can determine areas within the body where the endogenous version of human GPCRs disclosed herein is expressed and/or over-expressed. It is also possible using these techniques to determine related disease/disorder states which are associated with the expression and/or over-expression of the receptor; such an approach is disclosed in this patent document.

With respect to creation of a mutation that may evidence constitutive activation of the human GPCR disclosed herein is based upon the distance from the proline residue at which is presumed to be located within TM6 of the GPCR; this algorithmic technique is disclosed in co-pending and commonly assigned patent document PCT Application

5

10

15

Number PCT/US99/23938, published as WO 00/22129 on April 20, 2000, which, along with the other patent documents listed herein, is incorporated herein by reference. The algorithmic technique is not predicated upon traditional sequence "alignment" but rather a specified distance from the aforementioned TM6 proline residue (or, of course, endogenous constitutive substitution for such proline residue). By mutating the amino acid residue located 16 amino acid residues from this residue (presumably located in the IC3 region of the receptor) to, most preferably, a lysine residue, such activation may be obtained. Other amino acid residues may be useful in the mutation at this position to achieve this objective.

10

15

20

5

D. Disease/Disorder Identification and/or Selection

As will be set forth in greater detail below, most preferably inverse agonists and agonists to the non-endogenous, constitutively activated GPCR can be identified by the methodologies of this invention. Such inverse agonists and agonists are ideal candidates as lead compounds in drug discovery programs for treating diseases related to this receptor. Because of the ability to directly identify inverse agonists to the GPCR, thereby allowing for the development of pharmaceutical compositions, a search for diseases and disorders associated with the GPCR is relevant. For example, scanning both diseased and normal tissue samples for the presence of the GPCR now becomes more than an academic exercise or one which might be pursued along the path of identifying an endogenous ligand to the specific GPCR. Tissue scans can be conducted across a broad range of healthy and diseased tissues. Such tissue scans provide a preferred first step in associating a specific receptor with a disease and/or disorder.

Preferably, the DNA sequence of the human GPCR is used to make a probe for (a) dot-blot analysis against tissue-mRNA, and/or (b) RT-PCR identification of the expression of the receptor in tissue samples. The presence of a receptor in a tissue

source, or a diseased tissue, or the presence of the receptor at elevated concentrations in diseased tissue compared to a normal tissue, can be preferably utilized to identify a correlation with a treatment regimen, including but not limited to, a disease associated with that disease. Receptors can equally well be localized to regions of organs by this technique. Based on the known functions of the specific tissues to which the receptor is localized, the putative functional role of the receptor can be deduced.

E. Screening of Candidate Compounds

1. Generic GPCR screening assay techniques

When a G protein receptor becomes constitutively active, it binds to a G protein (e.g., Gq, Gs, Gi, Gz, Go) and stimulates the binding of GTP to the G protein. The G protein then acts as a GTPase and slowly hydrolyzes the GTP to GDP, whereby the receptor, under normal conditions, becomes deactivated. However, constitutively activated receptors continue to exchange GDP to GTP. A non-hydrolyzable analog of GTP, [35S]GTPγS, can be used to monitor enhanced binding to membranes which express constitutively activated receptors. It is reported that [35S]GTPγS can be used to monitor G protein coupling to membranes in the absence and presence of ligand. An example of this monitoring, among other examples well-known and available to those in the art, was reported by Traynor and Nahorski in 1995. The preferred use of this assay system is for initial screening of candidate compounds because the system is generically applicable to all G protein-coupled receptors regardless of the particular G protein that interacts with the intracellular domain of the receptor.

2. Specific GPCR screening assay techniques

Once candidate compounds are identified using the "generic" G protein-coupled receptor assay (i.e., an assay to select compounds that are agonists, partial agonists, or inverse agonists), further screening to confirm that the compounds have interacted at the

25

5

10

15

receptor site is preferred. For example, a compound identified by the "generic" assay may not bind to the receptor, but may instead merely "uncouple" the G protein from the intracellular domain.

a. Gs, Gz and Gi.

5

10

15

20

Gs stimulates the enzyme adenylyl cyclase. Gi (and Gz and Go), on the other hand, inhibit this enzyme. Adenylyl cyclase catalyzes the conversion of ATP to cAMP; thus, constitutively activated GPCRs that couple the Gs protein are associated with increased cellular levels of cAMP. On the other hand, constitutively activated GPCRs that couple Gi (or Gz, Go) protein are associated with decreased cellular levels of cAMP. See. generally, "Indirect Mechanisms of Synaptic Transmission," Chpt. 8, From Neuron To Brain (3rd Ed.) Nichols, J.G. et al eds. Sinauer Associates, Inc. (1992). Thus, assays that detect cAMP can be utilized to determine if a candidate compound is, e.g., an inverse agonist to the receptor (i.e., such a compound would decrease the levels of cAMP). A variety of approaches known in the art for measuring cAMP can be utilized; a most preferred approach relies upon the use of anti-cAMP antibodies in an ELISAbased format. Another type of assay that can be utilized is a whole cell second messenger reporter system assay. Promoters on genes drive the expression of the proteins that a particular gene encodes. Cyclic AMP drives gene expression by promoting the binding of a cAMP-responsive DNA binding protein or transcription factor (CREB) that then binds to the promoter at specific sites called cAMP response elements and drives the expression of the gene. Reporter systems can be constructed which have a promoter containing multiple cAMP response elements before the reporter gene, e.g., β-galactosidase or luciferase. Thus, a constitutively activated Gs-linked receptor causes the accumulation of cAMP that then activates the gene and expression of

the reporter protein. The reporter protein such as β -galactosidase or luciferase can then be detected using standard biochemical assays (Chen et al. 1995).

b. Go and Gq.

5

10

15

20

25

Gq and Go are associated with activation of the enzyme phospholipase C, which in turn hydrolyzes the phospholipid PIP₂, releasing two intracellular messengers: diacycloglycerol (DAG) and inistol 1,4,5-triphoispha¹ & (IP₃). Increased accumulation of IP₃ is associated with activation of Gq- and Go-associated receptors. See, generally, "Indirect Mechanisms of Synaptic Transmission," Chpt. 8, From Neuron To Brain (3rd Ed.) Nichols, J.G. et al eds. Sinauer Associates, Inc. (1992). Assays that detect IP₃ accumulation can be utilized to determine if a candidate compound is, e.g., an inverse agonist to a Gq- or Go-associated receptor (i.e., such a compound would decrease the levels of IP₃). Gq-associated receptors can also been examined using an AP1 reporter assay in that Gq-dependent phospholipase C causes activation of genes containing AP1 elements; thus, activated Gq-associated receptors will evidence an increase in the expression of such genes, whereby inverse agonists thereto will evidence a decrease in such expression, and agonists will evidence an increase in such expression. Commercially available assays for such detection are available.

3. GPCR Fusion Protein

The use of an endogenous, constitutively activate orphan GPCR or a non-endogenous, constitutively activated orphan GPCR, for use in screening of candidate compounds for the direct identification of inverse agonists, agonists and partial agonists provide an interesting screening challenge in that, by definition, the receptor is active even in the absence of an endogenous ligand bound thereto. Thus, in order to differentiate between, e.g., the non-endogenous receptor in the presence of a candidate compound and the non-endogenous receptor in the absence of that compound, with an

aim of such a differentiation to allow for an understanding as to whether such compound may be an inverse agonist, agonist, partial agonist or have no affect on such a receptor, it is preferred that an approach be utilized that can enhance such differentiation. A preferred approach is the use of a GPCR Fusion Protein.

Generally, once it is determined that a non-endogenous orphan GPCR has been constitutively activated using the assay techniques set forth above (as well as others), it is possible to determine the predominant G protein that couples with the endogenous GPCR. Coupling of the G protein to the GPCR provides a signaling pathway that can be assessed. Because it is most preferred that screening take place by use of a mammalian expression system, such a system will be expected to have endogenous G protein therein. Thus, by definition, in such a system, the non-endogenous, constitutively activated orphan GPCR will continuously signal. In this regard, it is preferred that this signal be enhanced such that in the presence of, e.g., an inverse agonist to the receptor, it is more likely that it will be able to more readily differentiate, particularly in the context of screening, between the receptor when it is contacted with the inverse agonist.

The GPCR Fusion Protein is intended to enhance the efficacy of G protein coupling with the non-endogenous GPCR. The GPCR Fusion Protein is preferred for screening with a non-endogenous, constitutively activated GPCR because such an approach increases the signal that is most preferably utilized in such screening techniques. This is important in facilitating a significant "signal to noise" ratio; such a significant ratio is import preferred for the screening of candidate compounds as disclosed herein.

The construction of a construct useful for expression of a GPCR Fusion Protein is within the purview of those having ordinary skill in the art. Commercially available expression vectors and systems offer a variety of approaches that can fit the particular

25

5

10

15

needs of an investigator. The criteria of importance for such a GPCR Fusion Protein construct is that the endogenous GPCR sequence and the G protein sequence both be inframe (preferably, the sequence for the endogenous GPCR is upstream of the G protein sequence) and that the "stop" codon of the GPCR must be deleted or replaced such that upon expression of the GPCR, the G protein can also be expressed. The GPCR can be linked directly to the G protein, or there can be spacer residues between the two (preferably, no more than about 12, although this number can be readily ascertained by one of ordinary skill in the art). We have a preference (based upon convenience) of use of a spacer in that some restriction sites that are not used will, effectively, upon expression, become a spacer. Most preferably, the G protein that couples to the nonendogenous GPCR will have been identified prior to the creation of the GPCR Fusion Protein construct. Because there are only a few G proteins that have been identified, it is preferred that a construct comprising the sequence of the G protein (i.e., a universal G protein construct) be available for insertion of an endogenous GPCR sequence therein; this provides for efficiency in the context of large-scale screening of a variety of different endogenous GPCRs having different sequences.

As noted above, constitutively activated GPCRs that couple to Gi, Gz and Go are expected to inhibit the formation of cAMP making assays based upon these types of GPCRs challenging (i.e., the cAMP signal decreases upon activation thus making the direct identification of, e.g, inverse agonists (which would further decrease this signal), interesting. As will be disclosed herein, we have ascertained that for these types of receptors, it is possible to create a GPCR Fusion Protein that is not based upon the endogenous GPCR's endogenous G protein, in an effort to establish a viable cyclase-based assay. Thus, for example, an endogenous Gi coupled receptor can be fused to a Gs protein – we believe that such a fusion construct, upon expression, "drives" or "forces"

25

5

10

15

the endogenous GPCR to couple with, e.g., Gs rather than the "natural" Gi protein, such that a cyclase-based assay can be established. Thus, for Gi, Gz and Go coupled receptors, we prefer that that when a GPCR Fusion Protein is used and the assay is based upon detection of adenylyl cyclase activity, that the fusion construct be established with Gs (or an equivalent G protein that stimulates the formation of the enzyme adenylyl cyclase).

Equally effective is a G Protein Fusion construct that utilizes a Gq Protein fused with a Gs, Gi, Gz or Go Protein. A most preferred fusion construct can be accomplished with a Gq Protein wherein the first six (6) amino acids of the G-protein α -subunit ("G α q") is deleted and the last five (5) amino acids at the C-terminal end of G α q is replaced with the corresponding amino acids of the G α of the G protein of interest. For example, a fusion construct can have a Gq (6 amino acid deletion) fused with a Gi Protein, resulting in a "Gq/Gi Fusion Construct". We believe that this fusion construct will force the endogenous Gi coupled receptor to couple to its non-endogenous G protein, Gq, such that the second messenger, for example, inositol triphosphate or diacylgycerol, can be measured in lieu of cAMP production.

4. Co-transfection of a Target Gi Coupled GPCR with a Signal-Enhancer Gs Coupled GPCR (cAMP Based Assays)

20

25

15

5

10

A Gi coupled receptor is known to inhibit adenylyl cyclase, and, therefore, decrease the level of cAMP production, which can make assessment of cAMP levels challenging. An effective technique in measuring the decrease in production of cAMP as an indication of constitutive activation of a receptor that predominantly couples Gi upon activation can be accomplished by co-transfecting a signal enhancer, e.g., a non-endogenous, constitutively activated receptor that predominantly couples with Gs upon activation (e.g., TSHR-A623I, disclosed below), with the Gi linked GPCR. As is

apparent, constitutive activation of a Gs coupled receptor can be determined based upon an increase in production of cAMP. Constitutive activation of a Gi coupled receptor leads to a decrease in production cAMP. Thus, the co-transfection approach is intended to advantageously exploit these "opposite" affects. For example, co-transfection of a non-endogenous, constitutively activated Gs coupled receptor (the "signal enhancer") with the endogenous Gi coupled receptor (the "target receptor") provides a baseline cAMP signal (i.e., although the Gi coupled receptor will decrease cAMP levels, this "decrease" will be relative to the substantial increase in cAMP levels established by constitutively activated Gs coupled signal enhancer). By then co-transfecting the signal enhancer with a constitutively activated version of the target receptor, cAMP would be expected to further decrease (relative to base line) due to the increased functional activity of the Gi target (i.e., which decreases cAMP).

Screening of candidate compounds using a cAMP based assay can then be accomplished, with two provisos: first, relative to the Gi coupled target receptor, "opposite" effects will result, *i.e.*, an inverse agonist of the Gi coupled target receptor will increase the measured cAMP signal, while an agonist of the Gi coupled target receptor will decrease this signal; second, as would be apparent, candidate compounds that are directly identified using this approach should be assessed independently to ensure that these do not target the signal enhancing receptor (this can be done prior to or after screening against the co-transfected receptors).

F. Medicinal Chemistry

Generally, but not always, direct identification of candidate compounds is preferably conducted in conjunction with compounds generated via combinatorial chemistry techniques, whereby thousands of compounds are randomly prepared for such analysis. Generally, the results of such screening will be compounds having

25

5

10

15

unique core structures; thereafter, these compounds are preferably subjected to additional chemical modification around a preferred core structure(s) to further enhance the medicinal properties thereof. Such techniques are known to those in the art and will not be addressed in detail in this patent document.

5

10

15

20

G. Pharmaceutical compositions

Candidate compounds selected for further development can be formulated into pharmaceutical compositions using techniques well known to those in the art. Suitable pharmaceutically-acceptable carriers are available to those in the art; for example, see Remington's Pharmaceutical Sciences, 16th Edition, 1980, Mack Publishing Co., (Oslo et al., eds.).

H. Other Utility

Although a preferred use of the non-endogenous versions the human GPCRs disclosed herein may be for the direct identification of candidate compounds as inverse agonists, agonists or partial agonists (preferably for use as pharmaceutical agents), these versions of human GPCRs can also be utilized in research settings. For example, *in vitro* and *in vivo* systems incorporating GPCRs can be utilized to further elucidate and understand the roles these receptors play in the human condition, both normal and diseased, as well as understanding the role of constitutive activation as it applies to understanding the signaling cascade. The value in non-endogenous human GPCRs is that their utility as a research tool is enhanced in that, because of their unique features, non-endogenous human GPCRs can be used to understand the role of these receptors in the human body before the endogenous ligand therefore is identified. Other uses of the disclosed receptors will become apparent to those in the art based upon, *inter alia*, a review of this patent document.

EXAMPLES

The following examples are presented for purposes of elucidation, and not limitation, of the present invention. While specific nucleic acid and amino acid sequences are disclosed herein, those of ordinary skill in the art are credited with the ability to make minor modifications to these sequences while achieving the same or substantially similar results reported below. The traditional approach to application or understanding of sequence cassettes from one sequence to another (e.g. from rat receptor to human receptor or from human receptor A to human receptor B) is generally predicated upon sequence alignment techniques whereby the sequences are aligned in an effort to determine areas of commonality. The mutational approach disclosed herein does not rely upon this approach but is instead based upon an algorithmic approach and a positional distance from a conserved proline residue located within the TM6 region of human GPCRs. Once this approach is secured, those in the art are credited with the ability to make minor modifications thereto to achieve substantially the same results (i.e., constitutive activation) disclosed herein. Such modified approaches are considered within the purview of this disclosure.

//

5

10

15

//

//

20 //

Example 1 ENDOGENOUS HUMAN GPCRS

1. Identification of Human GPCRs

The disclosed endogenous human GPCRs were identified based upon a review
of the GenBankTM database information. While searching the database, the following
cDNA clones were identified as evidenced below (Table C).

TABLE C

| Disclosed Human Orphan GPCRs | Accession Number Identified | Complete DNA Sequence (Base Pairs) | Open Reading Frame (Base Pairs) | Nucleic Acid SEQ.ID. NO. | Amino Acid SEQ.ID. NO. |
|---------------------------------------|-----------------------------------|--|---------------------------------------|-----------------------------------|---------------------------------|
| hRUP8 | AL121755 | 147,566bp | 1,152bp | 1 | 2 |
| hRUP9 | AC0113375 | 143,181bp | 1,260bp | 3 | 4 |
| hRUP10 | AC008745 | 94,194bp | 1,014bp | 5 | 6 |
| hRUP11 | AC013396 | 155,086bp | 1,272bp | 7 | 8 |
| hRUP12 | AP000808 | 177,764bp | 966bp | 9 | 10 |
| hRUP13 | AC011780 | 167,819bp | 1,356bp | 11 | 12 |
| hRUP14 | AL137118 | 168,297bp | 1,041bp | 13 | 14 |
| hRUP15 | AL016468 | 138,828bp | 1,527bp | 15 | 16 |
| hRUP16 | AL136106 | 208,042bp | 1,068bp | 17 | 18 |
| hRUP17 | AC023078 | 161,735bp | 969bp | 19 | 20 |
| hRUP18 | AC008547 | 117,304bp | 1,305bp | 21 | 22 |
| hRUP19 | AC026331 | 145,183bp | 1,041bp | 23 | 24 |
| hRUP20 | AL161458 | 163,511bp | 1,011bp | 25 | 26 |
| hRUP21 | AC026756 | 156,534bp | 1,014bp | 27 | 28 |
| hRUP22 | AC027026 | 151,811bp | 993bp | 29 | 30 |
| hRUP23 | AC007104 | 200,000bp | 1,092bp | 31 | 32 |
| hRUP24 | AL355388 | 190,538bp | - 1,125bp | 33 | 34 |
| hRUP25 | AC026331 | 145,183bp | 1,092bp | 35 | 36 |
| hRUP26 | AC023040 | 178,508bp | 1,044bp | 37 | 38 |
| hRUP27 | AC027643 | 158,700bp | 1,020bp | 39 | 40 |

2. Full Length Cloning

a. hRUP8 (Seq. Id. Nos. 1 & 2)

The disclosed human RUP8 was identified based upon the use of EST database (dbEST) information. While searching the dbEST, a cDNA clone with accession number

AL121755 was identified to encode a novel GPCR. The following PCR primers were used for RT-PCR with human testis Marathon-Ready cDNA (Clontech) as templates: 5'-CTTGCAGACATCACCATGGCAGCC-3' (SEQ.ID.NO.:41; sense) and 5'-GTGATGCTCTGAGTACTGGACTGG-3' (SEQ.ID.NO.: 42; antisense).

PCR was performed using Advantage cDNA polymerase (Clontech; manufacturing instructions will be followed) in 50ul reaction by the following cycles: 94°C for 30 sec; 94°C for 10 sec; 65°C for 20 sec, 72°C for 1.5 min, and 72°C for 7 min. Cycles 2 through 4 were repeated 35 times.

A 1.2kb PCR fragment was isolated and cloned into the pCRII-TOPO vector (Invitrogen) and sequenced using the ABI Big Dye Terminator kit (P.E. Biosystem). See, SEQ.ID.NO.:1. The putative amino acid sequence for RUP8 is set forth in SEQ.ID.NO.:2.

b. hRUP9 (Seq. Id. Nos. 3 & 4)

The disclosed human RUP9 was identified based upon the use of GeneBank

database information. While searching the database, a cDNA clone with Accession

Number AC011375 was identified as a human genomic sequence from chromosome

5. The full length RUP9 was cloned by PCR using primers:

- 5'-GAAGCTGTGAAGAGTGATGC-3' (SEQ.ID.NO.:43; sense),
- 5'-GTCAGCAATATTGATAAGCAGCAG-3' (SEQ.ID.NO.:44; antisense)
- and human genomic DNA (Promega) as a template. Taq Plus Precision polymerase (Stratagene) was used for the amplification in a 100µl reaction with 5% DMSO by the following cycle with step 2 to step 4 repeated 35 times: 94°C for 1 minute; 94°C for 30 seconds; 56°C for 30 seconds; 72°C for 2 minutes; 72°C for 5 minutes.

A 1.3 Kb PCR fragment was isolated and cloned into the pCRII-TOPO vector

(Invitrogen) from 1% agarose gel and completely sequenced using the ABI Big Dye

5

Terminator kit (P.E. Biosystem). *See*, SEQ.ID.NO.:3. The putative amino acid sequence for RUP8 is set forth in SEQ.ID.NO.:4. The sequence of RUP9 clones isolated from human genomic DNA matched with the sequence obtained from data base.

c. hRUP10 (Seq. Id. Nos. 5 & 6)

The disclosed human RUP10 was identified based upon the use of GenBank database information. While searching the database, a cDNA clone with accession number AC008754 was identified as a human genomic sequence from chromosome 19. The full length RUP10 was cloned by RT-PCR using primers:

5'-CCATGGGGAACGATTCTGTCAGCTACG-3' (SEQ.ID.NO.:45; sense) and

5'-GCTATGCCTGAAGCCAGTCTTGTG-3' (SEQ.ID.NO.:46; antisense) and human leukocyte Marathon-Ready cDNA (Clontech) as a template. Advantage cDNA polymerase (Clontech) was used for the amplification in a 50μl reaction by the following cycle with step 2 to step 4 repeated 35 times: 94°C for 30 seconds; 94°C for 10 seconds; 62°C for 20 seconds; 72°C for 1.5 minutes; 72°C for 7 minutes. A 1.0 Kb PCR fragment was isolated and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Terminator kit (P.E. Biosystem). The nucleic acid sequence of the novel human receptor RUP10 is set forth in SEQ.ID.NO.:5 and the putative amino acid sequence thereof is set forth in SEQ.ID.NO.:6.

20

5

10

15

d. hRUP11 (Seq. Id. Nos. 7 & 8)

The disclosed human RUP11 was identified based upon the use of GenBank database information. While searching the database, a cDNA clone with accession number AC013396 was identified as a human genomic sequence from chromosome 2.

The full length RUP11 was cloned by PCR using primers:

5'-CCAGGATGTTGTGTCACCGTGGTGGC-3' (SEQ.ID.NO.:47; sense),

5'-CACAGCGCTGCAGCCCTGCAGCTGGC-3' (SEQ.ID.NO.:48; antisense)

and human genomic DNA (Clontech) as a template. TaqPlus Precision DNA polymerase (Stratagene) was used for the amplification in a 50µl reaction by the following cycle with step 2 to step 4 repeated 35 times: 94°C for 3 minutes; 94°C for 20 seconds; 67°C for 20 seconds; 72°C for 1.5 minutes; 72°C for 7 minutes. A 1.3 Kb PCR fragment was isolated and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Terminator kit (P.E. Biosystem). The nucleic acid sequence of the novel human receptor RUP11 is set forth in SEQ.ID.NO.:7 and the putative amino acid sequence thereof is set forth in SEQ.ID.NO.:8.

e. hRUP12 (Seq. Id. Nos. 9 & 10)

The disclosed human RUP12 was identified based upon the use of GenBank database. While searching the database, a cDNA clone with accession number AP000808 was identified to encode a new GPCR, having significant homology with rat RTA and human mas1 oncogene GPCRs. The full length RUP12 was cloned by PCR using primers:

- 5'-CTTCCTCTCGTAGGGATGAACCAGAC-3' (SEQ.ID.NO.:49; sense)
- 5'-CTCGCACAGGTGGGAAGCACCTGTGG-3' (SEQ.ID.NO.:50; antisense)
- and human genomic DNA (Clontech) as template. TaqPlus Precision DNA polymerase (Stratagene) was used for the amplification by the following cycle with step 2 to step 4 repeated 35 times: 94°C for 3 min; 94°C for 20 sec; 65°C for 20sec; 72°C for 2 min and 72°C for 7 min. A 1.0kb PCR fragment was isolated and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Terminator kit

5

10

15

(P.E. Biosystem) (see, SEQ.ID.NO.:9 for nucleic acid sequence and SEQ.ID.NO.:10 for deduced amino acid sequence).

f. hRUP13 (Seq. Id. Nos. 11 & 12)

The disclosed human RUP13 was identified based upon the use of GenBank database. While searching the database, a cDNA clone with accession number 5 AC011780 was identified to encode a new GPCR, having significant homology with GPCR fish GPRX-ORYLA. The full length RUP13 was cloned by PCR using primers: 5'-GCCTGTGACAGGAGGTACCCTGG-3' (SEQ.ID.NO.:51; sense) 5'-CATATCCCTCCGAGTGTCCAGCGGC-3' (SEQ.ID.NO.:52; antisense)

and human genomic DNA (Clontech) as template. TaqPlus Precision DNA polymerase 10 (Stratagene) was used for the amplification by the following cycle with step 2 to step 4 repeated 35 times: 94°C for 3 min; 94°C for 20 sec; 65°C for 20sec; 72°C for 2 min and 72°C for 7 min. A 1.35kb PCR fragment was isolated and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Terminator kit (P.E. Biosystem) (see, SEQ.ID.NO.:11 for nucleic acid sequence and SEQ.ID.NO.:12 for deduced amino acid sequence).

g. hRUP14 (Seq. Id. Nos. 13 & 14)

The disclosed human RUP14 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession Number AL137118 was identified as a human genomic sequence from chromosome 13. The full length RUP14 was cloned by PCR using primers:

- 5'-GCATGGAGAAAATTTATGTCCTTGCAACC-3' (SEQ.ID.NO.:53; sense)
- 5'-CAAGAACAGGTCTCATCTAAGAGCTCC-3' (SEQ.ID.NO.:54; antisense)

and human genomic DNA (Promega) as a template. Taq Plus Precision polymerase

(Stratagene) and 5% DMSO were used for the amplification by the following cycle 25

15

with step 2 and step 3 repeated 35 times: 94°C for 3 minute; 94°C for 20 seconds; 58°C for 2 minutes; 72°C for 10 minutes.

A 1.1 Kb PCR fragment was isolated and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Terminator kit (P.E. Biosystem) (see, SEQ.ID.NO.:13 for nucleic acid sequence and SEQ.ID.NO.:14 for deduced amino acid sequence). The sequence of RUP14 clones isolated from human genomic DNA matched with the sequence obtained from database.

h. hRUP15 (Seq. Id. Nos. 15 & 16)

The disclosed human RUP15 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession Number AC016468 was identified as a human genomic sequence. The full length RUP15 was cloned by PCR using primers:

5'-GCTGTTGCCATGACGTCCACCTGCAC-3' (SEQ.ID.NO.:55; sense)

5'-GGACAGTTCAAGGTTTGCCTTAGAAC-3' (SEQ.ID.NO.:56; antisense)

and human genomic DNA (Promega) as a template. Taq Plus Precision polymerase (Stratagene) was used for the amplification by the following cycle with step 2 to 4 repeated 35 times: 94°C for 3 minute; 94°C for 20 seconds; 65°C for 20 seconds; 72°C for 2 minutes and 72°C for 7 minutes.

A 1.5 Kb PCR fragment was isolated and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Terminator kit (P.E. Biosystem). See, SEQ.ID.NO.:15 for nucleic acid sequence and SEQ.ID.NO.:16 for deduced amino acid sequence. The sequence of RUP15 clones isolated from human genomic DNA matched with the sequence obtained from database.

i. hRUP16 (Seq. Id. Nos. 17 & 18)

5

10

The disclosed human RUP16 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession Number AL136106 was identified as a human genomic sequence from chromosome 13. The full length RUP16 was cloned by PCR using primers:

- 5 5'-CTTTCGATACTGCTCCTATGCTC-3' (SEQ.ID.NO.:57; sense, 5' of initiation codon),
 5'-GTAGTCCACTGAAAGTCCAGTGATCC-3' (SEQ.ID.NO.:58; antisense, 3' of stop codon)
 and human skeletal muscle Marathon-Ready cDNA (Clontech) as template. Advantage
 cDNA polymerase (Clontech) was used for the amplification in a 50ul reaction by the
 following cycle with step 2 to 4 repeated 35 times: 94°C for 30 seconds; 94°C for 5
 seconds; 69°C for 15 seconds; 72°C for 1 minute and 72°C for 5 minutes.
 - A 1.1 Kb PCR fragment was isolated and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the T7 sequenase kit (Amsham). See, SEQ.ID.NO.:17 for nucleic acid sequence and SEQ.ID.NO.:18 for deduced amino acid sequence. The sequence of RUP16 clones matched with four unordered segments of AL136106, indicating that the RUP16 cDNA is composed of 4 exons.

j. hRUP17 (Seq. Id. Nos. 19 & 20)

The disclosed human RUP17 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession Number AC023078 was identified as a human genomic sequence from chromosome

- 20 11. The full length RUP17 was cloned by PCR using primers:
 - 5'-TTTCTGAGC<u>ATG</u>GATCCAACCATCTC-3' (SEQ.ID.NO.:59; sense, containing initiation codon)
 - 5'-CTGTCTGACAGGGCAGAGGCTCTTC-3' (SEQ.ID.NO.:60; antisense, 3' of stop codon) . and human genomic DNA (Promega) as template. Advantage cDNA polymerase mix
- 25 (Clontech) was used for the amplification in a 100ul reaction with 5% DMSO by the

following cycle with step 2 to 4 repeated 30 times: 94°C for 1 min; 94°C for 15 sec; 67°C for 20 sec; 72°C for 1 min and 30 sec; and 72°C for 5 min.

A 970bp PCR fragment was isolated from 1% agarose gel and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Termiantor Kit (P.E. Biosystem). *See*, SEQ.ID.NO.:19 for nucleic acid sequence and SEQ.ID.NO.:20 for deduced amino acid sequence.

k. hRUP18 (Seq. Id. Nos. 21 & 22)

The disclosed human RUP18 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession Number AC008547 was identified as a human genomic sequence from chromosome 5. The full length RUP18 was cloned by PCR using primers:

5'-GGAACTCGTATAGACCCAGCGTCGCTCC-3' (SEQ.ID.NO.:61; sense, 5' of the initiation codon),

5'-GGAGGTTGCGCCTTAGCGACAGATGACC-3' (SEQ.ID.NO.:62; antisense, 3' of stop codon)

and human genomic DNA (Promega) as template. TaqPlus precision DNA polymerase (Stratagene) was used for the amplification in a 100ul reaction with 5% DMSO by the following cycle with step 2 to 4 repeated 35 times: 95°C for 5 min; 95°C for 30 sec; 65°C for 30 sec; 72°C for 2 min; and 72°C for 5 min.

A 1.3kb PCR fragment was isolated from 1% agarose gel and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Termiantor Kit (P.E. Biosystem). *See*, SEQ.ID.NO.:21 for nucleic acid sequence and SEQ.ID.NO.:22 for deduced amino acid sequence.

l. hRUP19 (Seq. Id. Nos. 23 & 24)

20

5

The disclosed human RUP19 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession Number AC026331 was identified as a human genomic sequence from chromosome 12. The full length RUP19 was cloned by PCR using primers:

5 5'-CTGCACCCGGACACTTGCTCTG-3' (SEQ.ID.NO.:63; sense, 5' of initiation codon), 5'-GTCTGCTTGTTCAGTGCCACTCAAC-3' (SEQ.ID.NO.:64; antisense, containing the stop codon)

and human genomic DNA (Promega) as template. TaqPlus Precision DNA polymerase (Stratagene) was used for the amplification with 5% DMSO by the following cycle with step 2 to 4 repeated 35 times: 94°C for 1 min; 94°C for 15 sec; 70°C for 20 sec; 72°C for 1 min and 30 sec; and 72°C for 5 min.

A 1.1kp PCR fragment was isolated from 1% agarose gel and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Termiantor Kit (P.E. Biosystem). *See*, SEQ.ID.NO.:23 for nucleic acid sequence and SEQ.ID.NO.:24 for deduced amino acid sequence.

m. hRUP20 (Seq. Id. Nos. 25 & 26)

The disclosed human RUP20 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession Number AL161458 was identified as a human genomic sequence from chromosome

20 1. The full length RUP20 was cloned by PCR using primers:

5'-TATCTGCAATTCTATTCTAGCTCCTG-3' (SEQ.ID.NO.:65; sense, 5' of initiation codon), 5'-TGTCCCTAATAAAGTCACATGAATGC-3' (SEQ.ID.NO.:66; antisense, 3' of stop codon) and human genomic DNA (Promega) as template. Advantage cDNA polymerase mix (Clonetech) was used for the amplification with 5% DMSO by the following cycle with

10

step 2 to 4 repeated 35 times: 94°C for 1 min; 94°C for 15 sec; 60°C for 20 sec; 72°C for 1 min and 30 sec; and 72°C for 5 min.

A 1.0 kp PCR fragment was isolated from 1% agarose gel and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Termiantor Kit (P.E. Biosystem). *See*, SEQ.ID.NO.:25 for nucleic acid sequence and SEQ.ID.NO.:26 for deduced amino acid sequence.

n. hRUP21 (Seq. Id. Nos. 27 & 28)

The disclosed human RUP21 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession Number AC026756 was identified as a human genomic sequence from chromosome 13. The full length RUP21 was cloned by PCR using primers:

- 5'-GGAGACAACCATGAATGAGCCAC-3' (SEQ.ID.NO.:67; sense)
- 5'- TATTTCAAGGGTTGTTTGAGTAAC -3' (SEQ.ID.NO.:68; antisense)

and human genomic DNA (Promega) as template. Taq Plus Precision polymerase (Stratagene) was used for the amplification in a 100ul reaction with 5% DMSO by the following cycle with step 2 to 4 repeated 30 times; 94°C for 1 min; 94°C for 15 sec; 55°C for 20 sec; 72°C for 1 min and 30 sec; and 72°C for 5 min.

A 1,014 bp PCR fragment was isolated from 1% agarose gel and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Termiantor Kit (P.E. Biosystem). *See*, SEQ.ID.NO.:27 for nucleic acid sequence and SEQ.ID.NO.:28 for deduced amino acid sequence.

o. hRUP22 (Seq. Id. Nos. 29 & 30)

The disclosed human RUP22 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession

5

10

15

Number AC027026 was identified as a human genomic sequence from chromosome 11. The full length RUP22 was cloned by PCR using primers:

- 5'- GGCACCAGTGGAGGTTTTCTGAGCATG -3' (SEQ.ID.NO.:69; sense, containing initiation codon)
- 5 5'-CTGATGGAAGTAGAGGCTGTCCATCTC-3' (SEQ.ID.NO.:70; antisense, 3' of stop codon)

and human genomic DNA (Promega) as template. TaqPlus Precision DNA polymerase (Stratagene) was used for the amplification in a 100ul reaction with 5% DMSO by the following cycle with step 2 to 4 repeated 30 times: 94°C, 1 minutes 94°C, 15 seconds 55°C, 20 seconds 72°C, 1.5 minute 72°C, 5 minutes.

A 970bp PCR fragment was isolated from 1% agarose gel and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Termiantor Kit (P.E. Biosystem). *See*, SEQ.ID.NO.:29 for nucleic acid sequence and SEQ.ID.NO.:30 for deduced amino acid sequence.

p. hRUP23 (Seq. Id. Nos. 31 & 32)

The disclosed human RUP23 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession Number AC007104 was identified as a human genomic sequence from chromosome 4. The full length RUP23 was cloned by PCR using primers:

- 5'-CCTGGCGAGCCGCTAGCGCCATG-3' (SEQ.ID.NO.:71; sense, ATG as the initiation codon),
 - 5'-ATGAGCCCTGCCAGGCCC<u>TCA</u>GT-3' (SEQ.ID.NO.:72; antisense, TCA as the stop codon)
- and human placenta Marathon-Ready cDNA (Clontech) as template. Advantage cDNA polymerase (Clontech) was used for the amplification in a 50ul reaction by the following

cycle with step 2 to 4 repeated 35 times: 95°C for 30 sec; 95°C for 15 sec; 66°C for 20 sec; 72°C for 1 min and 20 sec; and 72°C for 5 min.

A 1.0 kb PCR fragment was isolated and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Terminator Kit (P.E. Biosystem). See, SEQ.ID.NO.:31 for nucleic acid sequence and SEQ.ID.NO.:32 for deduced amino acid sequence.

q. hRUP24 (Seq. Id. Nos. 33 & 34)

The disclosed human RUP25 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession Number AC026331 was identified as a human genomic sequence from chromosome 12. The full length RUP25 was cloned by PCR using primers: 5'-GCTGGAGCATTCACTAGGCGAG-3' (SEQ.ID.NO.:73; sense, 5'of initiation codon), 5'-AGATCCTGGTTCTTGGTGACAATG-3' (SEQ.ID.NO.:74; antisense, 3' of stop codon) and human genomic DNA (Promega) as template. Advantage cDNA polymerase mix (Clontech) was used for the amplification with 5% DMSO by the following cycle with step 2 to 4 repeated 35 times: 94°C for 1 minute; 94°C for 15 seconds; 56°C for 20 seconds 72°C for 1 minute 30 seconds and 72°C for 5 minutes.

A 1.2kb PCR fragment was isolated from 1% agarose gel and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Termiantor Kit (P.E. Biosystem). *See*, SEQ.ID.NO.:33 for nucleic acid sequence and SEQ.ID.NO.:34 for deduced amino acid sequence.

r. hRUP25 (Seq. Id. Nos. 35 & 36)

The disclosed human RUP25 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession

5

10

15

Number AC026331 was identified as a human genomic sequence from chromosome 12. The full length RUP25 was cloned by PCR using primers:

- 5'-GCTGGAGCATTCACTAGGCGAG-3' (SEQ.ID.NO.:75; sense, 5'of initiation codon),
- 5'-AGATCCTGGTTCTTGGTGACAATG-3' (SEQ.ID.NO.:76; antisense, 3' of stop codon)
- and human genomic DNA (Promega) as template. Advantage cDNA polymerase mix (Clontech) was used for the amplification with 5% DMSO by the following cycle with step 2 to 4 repeated 35 times: 94°C for 1 minute; 94°C for 15 seconds; 56°C for 20 seconds 72°C for 1 minute 30 seconds and 72°C for 5 minutes.

A 1.2kb PCR fragment was isolated from 1% agarose gel and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Termiantor Kit (P.E. Biosystem). *See*, SEQ.ID.NO.:35 for nucleic acid sequence and SEQ.ID.NO.:36 for deduced amino acid sequence.

s. hRUP26 (Seq. Id. Nos. 37 & 38)

The disclosed human RUP26 was identified based upon the use of GeneBank

database information. While searching the database, a cDNA clone with Accession

Number AC023040 was identified as a human genomic sequence from chromosome

The full length RUP26 was cloned by RT-PCR using RUP26 specific primers:

5'-AGCCATCCCTGCCAGGAAGCATGG-3' (SEQ.ID.NO.:77; sense, containing initiation codon)

5'-CCAGACTGTGGACTCAAGAACTCTAGG-3' (SEQ.ID.NO.:78; antisense, containing stop codon)
 and human pancreas Marathon - Ready cDNA (Clontech) as template. Advantage cDNA polymerase mix (Clontech) was used for the amplification in a 100µl reaction with 5% DMSO by the following cycle with step 2 to 4 repeated 35 times: 94°C for 5 minute;
 95°C for 30 seconds; 65°C for 30 seconds 72°C for 2 minute and 72°C for 5 minutes.

A 1.1kb PCR fragment was isolated from 1% agarose gel and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Termiantor Kit (P.E. Biosystem). *See*, SEQ.ID.NO.:37 for nucleic acid sequence and SEQ.ID.NO.:38 for deduced amino acid sequence.

t. hRUP27 (Seq. Id. Nos. 39 & 40)

The disclosed human RUP27 was identified based upon the use of GeneBank database information. While searching the database, a cDNA clone with Accession Number AC027643 was identified as a human genomic sequence from chromosome 12. The full length RUP27 was cloned by PCR using RUP27 specific primers:

5'-AGTCCACGAACAATGAATCCATTTCATG-3' (SEQ.ID.NO.:79; sense, containing initiation codon),

5'-ATCATGTCTAGACTCATGGTGATCC-3' (SEQ.ID.NO.:80; antisense, 3' of stop codon) and the human adult brain Marathon-Ready cDNA (Clontech) as template. Advantage cDNA polymerase mix (Clontech) was used for the amplification in a 50µl reaction with 5% DMSO by the following cycle with step 2 to 4 repeated 35 times: 94°C for 1 minute; 94°C for 10 seconds; 58°C for 20 seconds 72°C for 1 minute 30 seconds and 72°C for 5 minutes.

A 1.1kb PCR fragment was isolated from 1% agarose gel and cloned into the pCRII-TOPO vector (Invitrogen) and completely sequenced using the ABI Big Dye Termiantor Kit (P.E. Biosystem). *See*, SEQ.ID.NO.:35 for nucleic acid sequence and SEQ.ID.NO.:36 for deduced amino acid sequence. The sequence of RUP27 cDNA clone isolated from human brain was determined to match with five unordered segments of AC027643, indicating that the RUP27 cDNA is composed of 5 exons.

5

15

Example 2 PREPARATION OF NON-ENDOGENOUS, CONSTITUTIVELY ACTIVATED GPCRS

Those skilled in the art are credited with the ability to select techniques for mutation of a nucleic acid sequence. Presented below are approaches utilized to create non-endogenous versions of several of the human GPCRs disclosed above. The mutations disclosed below are based upon an algorithmic approach whereby the 16th amino acid (located in the IC3 region of the GPCR) from a conserved proline (or an endogenous, conservative substitution therefore) residue (located in the TM6 region of the GPCR, near the TM6/IC3 interface) is mutated, preferably to an alanine, histidine, arginine or lysine amino acid residue, most preferably to a lysine amino acid residue.

1. Transformer Site-Directed TM Mutagenesis

Preparation of non-endogenous human GPCRs may be accomplished on human GPCRs using Transformer Site-DirectedTM Mutagenesis Kit (Clontech) according to the manufacturer instructions. Two mutagenesis primers are utilized, most preferably a lysine mutagenesis oligonucleotide that creates the lysine mutation, and a selection marker oligonucleotide. For convenience, the codon mutation to be incorporated into the human GPCR is also noted, in standard form (Table D):

20

5

10

TABLE D

| Receptor Identifier | Codon Mutation |
|---------------------|----------------|
| hRUP8 | V274K |
| hRUP9 | T249K |
| hRUP10 | R232K |
| hRUP11 | M294K |
| hRUP12 | F220K |
| hRUP16 | A238K |

| Y215K | | |
|-------|--|--|
| L294K | | |
| T219K | | |
| K248A | | |
| K248H | | |
| K248R | | |
| R240K | | |
| Y222K | | |
| A245K | | |
| I230K | | |
| V285K | | |
| T248K | | |
| | | |

2. QuikChange™ Site-Directed™ Mutagenesis

Preparation of non-endogenous human GPCRs can also be accomplished by

using QuikChangeTM Site-DirectedTM Mutagenesis Kit (Stratagene, according to
manufacturer's instructions). Endogenous GPCR is preferably used as a template and
two mutagenesis primers utilized, as well as, most preferably, a lysine mutagenesis
oligonucleotide and a selection marker oligonucleotide (included in kit). For
convenience, the codon mutation incorporated into the novel human GPCR and the
respective oligonucleotides are noted, in standard form (Table E):

TABLE E

| Receptor Identifier | Codon Mutation | 5'-3' orientation (sense), (SEQ.ID.NO.) mutation underlined | 5'-3' orientation (antisense) (SEQ.ID.NO.) | Cycle Conditions Min ('), Sec (") Cycles 2-4 repeated 16 times |
|------------------------|-------------------|---|---|--|
| hRUP13 | A268K | GGGGAGGGAAAGCAA AGGTGGTCCTCCTGG (81) | CCAGGAGAACCACCT TTGCTTTCCCTCCCC (82) | 98° for 2' 98° for 30" 56°C for 30" 72° for 11' 40" 72° for 5' |
| hRUP14 | L246K | CAGGAAGGCAAAGAC CACCATCATCATC (85) | GATGATGATGGTGGT CTTTGCCTTCCTG (86) | 98° for 2' 98° for 30" 55°C for 30" 72° for 11' 40" 72° for 5' |

| hRUP15 | A398K | CCAGTGCAAAGCTAAG AAAGTGATCTTC (89) | GAAGATCACTTTCTTA GCTTTGCACTGG (90) | 98° for 2' 98° for 30" 55°C for 30" 72° for 11' 40" 72° for 5' |
|--------|-------|---------------------------------------|---|--|
| hRUP23 | W275K | GCCGCCACCGCCCAA GAGGAAGATTGGC (93) | GCCAATCTTCCTCTTG GCGCGGTGGCGGC (94) | 98° for 2' 98° for 30" 56°C for 30" 72° for 11' 40" 72° for 5' |

The non-endogenous human GPCRs were then sequenced and the derived and verified nucleic acid and amino acid sequences are listed in the accompanying "Sequence Listing" appendix to this patent document, as summarized in Table F below:

TABLE F

| Non Endogenous Human GPCR | Nucleic Acid Sequence Listing | Amino Acid Sequence Listing |
|------------------------------|-------------------------------|--------------------------------|
| hRUP13 | SEQ.ID.NO.:83 | SEQ.ID.NO.:84 |
| hRUP14 | SEQ.ID.NO.:87 | SEQ.ID.NO.:88 |
| hRUP15 | SEQ.ID.NO.:91 | SEQ.ID.NO.:92 |
| hRUP23 | SEQ.ID.NO.:95 | SEQ.ID.NO.:96 |

Example 3 RECEPTOR EXPRESSION

Although a variety of cells are available to the art for the expression of proteins, it is most preferred that mammalian cells be utilized. The primary reason for this is predicated upon practicalities, *i.e.*, utilization of, *e.g.*, yeast cells for the expression of a GPCR, while possible, introduces into the protocol a non-mammalian cell which may not (indeed, in the case of yeast, does not) include the receptor-coupling, genetic-mechanism and secretary pathways that have evolved for mammalian systems — thus, results obtained in non-mammalian cells, while of

5

10

potential use, are not as preferred as that obtained from mammalian cells. Of the mammalian cells, COS-7, 293 and 293T cells are particularly preferred, although the specific mammalian cell utilized can be predicated upon the particular needs of the artisan.

a. Transient Transfection

On day one, 6x10⁶/10 cm dish of 293 cells well were plated out. On day two, two reaction tubes were prepared (the proportions to follow for each tube are per plate): tube A was prepared by mixing 4µg DNA (e.g., pCMV vector; pCMV vector with receptor cDNA, etc.) in 0.5 ml serum free DMEM (Gibco BRL); tube B was prepared by mixing 24µl lipofectamine (Gibco BRL) in 0.5ml serum free DMEM. Tubes A and B were admixed by inversions (several times), followed by incubation at room temperature for 30-45min. The admixture is referred to as the "transfection mixture". Plated 293 cells were washed with 1XPBS, followed by addition of 5 ml serum free DMEM. 1 ml of the transfection mixture were added to the cells, followed by incubation for 4hrs at 37°C/5% CO₂. The transfection mixture was removed by aspiration, followed by the addition of 10ml of DMEM/10% Fetal Bovine Serum. Cells were incubated at 37°C/5% CO₂. After 48hr incubation, cells were harvested and utilized for analysis.

b. Stable Cell Lines: Gs Fusion Protein

Approximately 12x10⁶ 293 cells are plated on a 15cm tissue culture plate. Grown in DME High Glucose Medium containing ten percent fetal bovine serum and one percent sodium pyruvate, L-glutamine, and anti-biotics. Twenty-four hours following plating of 293 cells to ~80% confluency, the cells are transfected using 12μg of DNA. The 12μg of DNA is combined with 60ul of lipofectamine and 2mL of DME High Glucose Medium without serum. The medium is aspirated from the plates and the cells are washed once with medium without serum. The DNA, lipofectamine, and

25

5

10

15

medium mixture is added to the plate along with 10mL of medium without serum. Following incubation at 37 degrees Celsius for four to five hours, the medium is aspirated and 25ml of medium containing serum is added. Twenty-four hours following transfection, the medium is aspirated again, and fresh medium with serum is added. Forty-eight hours following transfection, the medium is aspirated and medium with serum is added containing geneticin (G418 drug) at a final concentration of 500µg/m½. The transfected cells now undergo selection for positively transfected cells containing the G418 resistant gene. The medium is replaced every four to five days as selection occurs. During selection, cells are grown to create stable pools, or split for stable clonal selection.

Example 4 ASSAYS FOR DETERMINATION OF CONSTITUTIVE ACTIVITY OF NON-ENDOGENOUS GPCRS

A variety of approaches are available for assessment of constitutive activity of the non-endogenous human GPCRs. The following are illustrative; those of ordinary skill in the art are credited with the ability to determine those techniques that are preferentially beneficial for the needs of the artisan.

1. Membrane Binding Assays: [35S]GTPγS Assay

When a G protein-coupled receptor is in its active state, either as a result of ligand binding or constitutive activation, the receptor couples to a G protein and stimulates the release of GDP and subsequent binding of GTP to the G protein. The alpha subunit of the G protein-receptor complex acts as a GTPase and slowly hydrolyzes the GTP to GDP, at which point the receptor normally is deactivated. Constitutively activated receptors continue to exchange GDP for GTP. The non-hydrolyzable GTP analog, [35 S]GTP γ S, can be utilized to demonstrate enhanced binding of [35 S]GTP γ S to membranes expressing constitutively activated receptors. The advantage of using

5

10

15

20

[35S]GTPγS binding to measure constitutive activation is that: (a) it is generically applicable to all G protein-coupled receptors; (b) it is proximal at the membrane surface making it less likely to pick-up molecules which affect the intracellular cascade.

The assay utilizes the ability of G protein coupled receptors to stimulate [35S]GTPγS binding to membranes expressing the relevant receptors. The assay can, therefore, be used in the direct identification method to screen candidate compounds to known, orphan and constitutively activated G protein-coupled receptors. The assay is generic and has application to drug discovery at all G protein-coupled receptors.

The [35S]GTPγS assay was incubated in 20 mM HEPES and between 1 and about 20mM MgCl₂ (this amount can be adjusted for optimization of results, although 20mM is preferred) pH 7.4, binding buffer with between about 0.3 and about 1.2 nM [35S]GTPγS (this amount can be adjusted for optimization of results, although 1.2 is preferred) and 12.5 to 75 μg membrane protein (e.g., 293 cells expressing the Gs Fusion Protein; this amount can be adjusted for optimization) and 10 μM GDP (this amount can be changed for optimization) for 1 hour. Wheatgerm agglutinin beads (25 μl; Amersham) were then added and the mixture incubated for another 30 minutes at room temperature. The tubes were then centrifuged at 1500 x g for 5 minutes at room temperature and then counted in a scintillation counter.

2. Adenylyl Cyclase

A Flash PlateTM Adenylyl Cyclase kit (New England Nuclear; Cat. No. SMP004A) designed for cell-based assays can be modified for use with crude plasma membranes. The Flash Plate wells can contain a scintillant coating which also contains a specific antibody recognizing cAMP. The cAMP generated in the wells can be quantitated by a direct competition for binding of radioactive cAMP tracer to the cAMP

5

10

15

antibody. The following serves as a brief protocol for the measurement of changes in cAMP levels in whole cells that express the receptors.

Transfected cells were harvested approximately twenty four hours after transient transfection. Media is carefully aspirated off and discarded. 10ml of PBS is gently added to each dish of cells followed by careful aspiration. 1ml of Sigma cell dissociation buffer and 3ml of PBS are added to each plate. Cells were pipeted off the plate and the cell suspension was collected into a 50ml conical centrifuge tube. Cells were then centrifuged at room temperature at 1,100 rpm for 5 min. The cell pellet was carefully re-suspended into an appropriate volume of PBS (about 3ml/plate). The cells were then counted using a hemocytometer and additional PBS was added to give the appropriate number of cells (with a final volume of about 50 μl/well).

cAMP standards and Detection Buffer (comprising 1 μCi of tracer [125 I cAMP (50 μl] to 11 ml Detection Buffer) was prepared and maintained in accordance with the manufacturer's instructions. Assay Buffer was prepared fresh for screening and contained 50μl of Stimulation Buffer, 3ul of test compound (12uM final assay concentration) and 50μl cells, Assay Buffer was stored on ice until utilized. The assay was initiated by addition of 50μl of cAMP standards to appropriate wells followed by addition of 50ul of PBSA to wells H-11 and H12. 50μl of Stimulation Buffer was added to all wells. DMSO (or selected candidate compounds) was added to appropriate wells using a pin tool capable of dispensing 3μl of compound solution, with a final assay concentration of 12μM test compound and 100μl total assay volume. The cells were then added to the wells and incubated for 60 min at room temperature. 100μl of Detection Mix containing tracer cAMP was then added to the wells. Plates were then incubated additional 2 hours followed by counting in a Wallac MicroBeta scintillation

5

10

15

counter. Values of cAMP/well were then extrapolated from a standard cAMP curve which was contained within each assay plate.

3. Cell-Based cAMP for Gi Coupled Target GPCRs

TSHR is a Gs coupled GPCR that causes the accumulation of cAMP upon activation. TSHR will be constitutively activated by mutating amino acid residue 623 (i.e., changing an alanine residue to an isoleucine residue). A Gi coupled receptor is expected to inhibit adenylyl cyclase, and, therefore, decrease the level of cAMP production, which can make assessment of cAMP levels challenging. An effective technique for measuring the decrease in production of cAMP as an indication of constitutive activation of a Gi coupled receptor can be accomplished by co-transfecting, most preferably, non-endogenous, constitutively activated TSHR (TSHR-A623I) (or an endogenous, constitutively active Gs coupled receptor) as a "signal enhancer" with a Gi linked target GPCR to establish a baseline level of cAMP. Upon creating a nonendogenous version of the Gi coupled receptor, this non-endogenous version of the target GPCR is then co-transfected with the signal enhancer, and it is this material that can be used for screening. We will utilize such approach to effectively generate a signal when a cAMP assay is used; this approach is preferably used in the direct identification of candidate compounds against Gi coupled receptors. It is noted that for a Gi coupled GPCR, when this approach is used, an inverse agonist of the target GPCR will increase the cAMP signal and an agonist will decrease the cAMP signal.

On day one, 2X10⁴ 293 and 293 cells/well will be plated out. On day two, two reaction tubes will be prepared (the proportions to follow for each tube are per plate): tube A will be prepared by mixing 2µg DNA of each receptor transfected into the mammalian cells, for a total of 4µg DNA (e.g., pCMV vector; pCMV vector with mutated THSR (TSHR-A623I); TSHR-A623I and GPCR, etc.) in 1.2ml serum free

25

5

10

15

DMEM (Irvine Scientific, Irvine, CA); tube B will be prepared by mixing 120μl lipofectamine (Gibco BRL) in 1.2ml serum free DMEM. Tubes A and B will then be admixed by inversions (several times), followed by incubation at room temperature for 30-45min. The admixture is referred to as the "transfection mixture". Plated 293 cells will be washed with 1XPBS, followed by addition of 10ml serum free DMEM. 2.4ml of the transfection mixture will then be added to the cells, followed by incubation for 4hrs at 37°C/5% CO₂. The transfection mixture will then be removed by aspiration, followed by the addition of 25ml of DMEM/10% Fetal Bovine Serum. Cells will then be incubated at 37°C/5% CO₂. After 24hr incubation, cells will then be harvested and utilized for analysis.

A Flash PlateTM Adenylyl Cyclase kit (New England Nuclear; Cat. No. SMP004A) is designed for cell-based assays, however, can be modified for use with crude plasma membranes depending on the need of the skilled artisan. The Flash Plate wells will contain a scintillant coating which also contains a specific antibody recognizing cAMP. The cAMP generated in the wells can be quantitated by a direct competition for binding of radioactive cAMP tracer to the cAMP antibody. The following serves as a brief protocol for the measurement of changes in cAMP levels in whole cells that express the receptors.

Transfected cells will be harvested approximately twenty four hours after transient transfection. Media will be carefully aspirated off and discarded. 10ml of PBS will be gently added to each dish of cells followed by careful aspiration. 1ml of Sigma cell dissociation buffer and 3ml of PBS will be added to each plate. Cells will be pipeted off the plate and the cell suspension will be collected into a 50ml conical centrifuge tube. Cells will then be centrifuged at room temperature at 1,100 rpm for 5 min. The cell pellet will be carefully re-suspended into an appropriate volume of PBS (about

25

5

10

15

3ml/plate). The cells will then be counted using a hemocytometer and additional PBS is added to give the appropriate number of cells (with a final volume of about 50µl/well).

cAMP standards and Detection Buffer (comprising 1 μCi of tracer [125] cAMP (50 μl] to 11 ml Detection Buffer) will be prepared and maintained in accordance with the manufacturer's instructions. Assay Buffer should be prepared fresh for screening and contained 50μl of Stimulation Buffer, 3ul of test compound (12uM final assay concentration) and 50μl cells, Assay Buffer can be stored on ice until utilized. The assay can be initiated by addition of 50μl of cAMP standards to appropriate wells followed by addition of 50μl of PBSA to wells H-11 and H12. 50ul of Stimulation Buffer will be added to all wells. Selected compounds (e.g., TSH) will be added to appropriate wells using a pin tool capable of dispensing 3μl of compound solution, with a final assay concentration of 12μM test compound and 100μl total assay volume. The cells will then be added to the wells and incubated for 60 min at room temperature. 100μl of Detection Mix containing tracer cAMP will then be added to the wells. Plates were then incubated additional 2 hours followed by counting in a Wallac MicroBeta scintillation counter. Values of cAMP/well will then be extrapolated from a standard cAMP curve which is contained within each assay plate.

4. Reporter-Based Assays

a. CRE-LUC Reporter Assay (Gs-associated receptors)

293 and 293T cells are plated-out on 96 well plates at a density of 2 x 10⁴ cells per well and were transfected using Lipofectamine Reagent (BRL) the following day according to manufacturer instructions. A DNA/lipid mixture is prepared for each 6-well transfection as follows: 260ng of plasmid DNA in 100µl of DMEM were gently mixed with 2µl of lipid in 100µl of DMEM (the 260ng of plasmid DNA consisted of

25

5

10

15

200ng of a 8xCRE-Luc reporter plasmid, 50ng of pCMV comprising endogenous receptor or non-endogenous receptor or pCMV alone, and 10ng of a GPRS expression plasmid (GPRS in pcDNA3 (Invitrogen)). The 8XCRE-Luc reporter plasmid was prepared as follows: vector SRIF-β-gal was obtained by cloning the rat somatostatin promoter (-71/+51) at BglV-HindIII site in the pβgal-Basic Vector (Clontech). Eight (8) copies of cAMP response element were obtained by PCR from an adenovirus template AdpCF126CCRE8 (see, 7 Human Gene Therapy 1883 (1996)) and cloned into the SRIF-β-gal vector at the Kpn-BglV site, resulting in the 8xCRE-β-gal reporter vector. The 8xCRE-Luc reporter plasmid was generated by replacing the beta-galactosidase gene in the 8xCRE-β-gal reporter vector with the luciferase gene obtained from the pGL3-basic vector (Promega) at the HindIII-BamHI site. Following 30 min. incubation at room temperature, the DNA/lipid mixture was diluted with 400 µl of DMEM and 100µl of the diluted mixture was added to each well. 100 μ l of DMEM with 10% FCS were added to each well after a 4hr incubation in a cell culture incubator. The following day the transfected cells were changed with 200 µl/well of DMEM with 10% FCS. Eight (8) hours later, the wells were changed to 100 µl /well of DMEM without phenol red, after one wash with PBS. Luciferase activity were measured the next day using the LucLite™ reporter gene assay kit (Packard) following manufacturer instructions and read on a 1450 MicroBeta™ scintillation and luminescence counter (Wallac).

b. AP1 reporter assay (Gq-associated receptors)

A method to detect Gq stimulation depends on the known property of Gq-dependent phospholipase C to cause the activation of genes containing AP1 elements in their promoter. A PathdetectTM AP-1 cis-Reporting System (Stratagene, Catalogue # 219073) can be utilized following the protocol set forth above with respect to the

5

10

15

20

CREB reporter assay, except that the components of the calcium phosphate precipitate were 410 ng pAP1-Luc, 80 ng pCMV-receptor expression plasmid, and 20 ng CMV-SEAP.

c. SRF-LUC Reporter Assay (Gq- associated receptors)

One method to detect Gq stimulation depends on the known property of Gqdependent phospholipase C to cause the activation of genes containing serum response factors in their promoter. A PathdetectTM SRF-Luc-Reporting System (Stratagene) can be utilized to assay for Gq coupled activity in, e.g., COS7 cells. Cells are transfected with the plasmid components of the system and the indicated expression plasmid encoding endogenous or non-endogenous GPCR using a Mammalian Transfection™ Kit (Stratagene, Catalogue #200285) according to the Briefly, 410 ng SRF-Luc, 80 ng pCMV-receptor manufacturer's instructions. expression plasmid and 20 ng CMV-SEAP (secreted alkaline phosphatase expression plasmid; alkaline phosphatase activity is measured in the media of transfected cells to control for variations in transfection efficiency between samples) are combined in a calcium phosphate precipitate as per the manufacturer's instructions. Half of the precipitate is equally distributed over 3 wells in a 96-well plate, kept on the cells in a serum free media for 24 hours. The last 5 hours the cells are incubated with $1\mu M$ Angiotensin, where indicated. Cells are then lysed and assayed for luciferase activity using a Luclite™ Kit (Packard, Cat. # 6016911) and "Trilux 1450 Microbeta" liquid scintillation and luminescence counter (Wallac) as per the manufacturer's instructions. The data can be analyzed using GraphPad Prism™ 2.0a (GraphPad Software Inc.).

20

5

10

d. Intracellular IP₃ Accumulation Assay (Gq-associated receptors)

On day 1, cells comprising the receptors (endogenous and/or non-endogenous) can be plated onto 24 well plates, usually 1x10⁵ cells/well (although his umber can be optimized. On day 2 cells can be transfected by firstly mixing $0.25\mu g$ DNA in 50 μl serum free DMEM/well and 2 µl lipofectamine in 50 µl serumfree DMEM/well. The solutions are gently mixed and incubated for 15-30 min at room temperature. Cells are washed with 0.5 ml PBS and 400 µl of serum free media is mixed with the transfection media and added to the cells. The cells are then incubated for 3-4 hrs at 37°C/5%CO₂ and then the transfection media is removed and replaced with 1ml/well of regular growth media. On day 3 the cells are labeled with ³H-myo-inositol. Briefly, the media is removed and the cells are washed with 0.5 ml PBS. Then 0.5 ml inositol-free/serum free media (GIBCO BRL) is added/well with 0.25 μ Ci of ³H-myo-inositol/ well and the cells are incubated for 16-18 hrs o/n at 37°C/5%CO2. On Day 4 the cells are washed with 0.5 ml PBS and 0.45 ml of assay medium is added containing inositol-free/serum free media 10 μM pargyline 10 mM lithium chloride or 0.4 ml of assay medium and 50 μl of 10x ketanserin (ket) to final concentration of $10\mu M$. The cells are then incubated for 30 min at 37°C. The cells are then washed with 0.5 ml PBSand 200µl of fresh/icecold stop solution (1M KOH; 18 mM Na-borate; 3.8 mM EDTA) is added/well. The solution is kept on ice for 5-10 min or until cells were lysed and then neutralized by 200 μl of fresh/ice cold neutralization sol. (7.5 % HCL). The lysate is then transferred into 1.5 ml eppendorf tubes and 1 ml of chloroform/methanol (1:2) is added/tube. The solution is vortexed for 15 sec and the upper phase is applied to a Biorad AG1-X8TM anion exchange resin (100-200 mesh). Firstly, the resin is washed with water at 1:1.25 W/V and 0.9 ml of upper phase is loaded onto the column. The column is washed with 10 mls of 5 mM myo-inositol and 10 ml of 5 mM Na-borate/60mM Na-formate. The inositol

5

10

15

20

tris phosphates are eluted into scintillation vials containing 10 ml of scintillation cocktail with 2 ml of 0.1 M formic acid/1 M ammonium formate. The columns are regenerated by washing with 10 ml of 0.1 M formic acid/3M ammonium formate and rinsed twice with dd $\rm H_2O$ and stored at $\rm 4^oC$ in water.

Exemplary results are presented below in Table G:

TABLE G

| Receptor | Mutation | Assay Utilized Figure No.) | Signal Generated: CMV | Signal Generated: Endogenous Version (Relative Light Units) | Signal Generated: Non- Endogenous Version (Relative Light Units) | Difference (<=⟨) Between CMV v. Wild-type Wild-type v. Mutant |
|----------|----------|----------------------------------|-----------------------------------|---|--|---|
| hRUP12 | N/A | IP ₃ (Figure 1) | 317.03 cpm/mg protein | 3463.29 cpm/mg protein | | 1. 11 Fold ← |
| hRUP13 | N/A | cAMP (Figure 2) | 8.06 pmol/cAMP/mg protein | 19.10 pmol/cAMP/mg protein | | 1. 2.4 Fold ← |
| | A268K | 8XCRE- LUC (Figure 3) | 3665.43 LCPS | 83280.17 LPCS | 61713.6 LCPS | 1. 23 Fold ← 2. 26 % ⟨ |
| hRUP14 | L246K | 8XCRE- LUC (Figure 5) | 86.07 LCPS | . 1962.87 LCPS | 789.73 LCPS | 23 Fold ← 60% ⟨ |
| hRUP15 | A398K | 8XCRE- LUC (Figure 6) | 86.07 LCPS | 18286.77 LCPS | 17034.83 LCPS | 1. 212 Fold ← 2. 1%⟨ |
| | A398K | cAMP (Figure 7) | 15.00 pmol/cAMP/mg protein | 164.4 pmol/cAMP/mg protein | 117.5 pmol/cAMP/ mg protein | 1. 11 Fold ← 2. 29% ⟨ |
| hRUP17 | N/A | IP ₃ (Figure 9) | 317.03 cpm/mg protein | 741.07 cpm/mg protein | | 1. 2.3 Fold |
| hRUP21 | N/A | IP ₃ (Figure 10) | 730.5 cpm/mg protein | 1421.9 cpm/mg protein | | 1. 2 Fold ⇐ |
| hRUP23 | W275K | 8XCRE- LUC (Figure 11) | 311.73 pmol/cAMP/mg protein | 13756.00 pmol/cAMP/mg protein | 9756.87 pmol/cAMP/ mg protein | 1. 44 Fold ← 2. 30% ⟨ |

N/A = not applied

Exemplary results of GTPγS assay for detecting constitutive activation, as disclosed in Example 4(1) above, was accomplished utilizing Gs:Fusion Protein Constructs on human RUP13 and RUP15. Table H below lists the signals generated from this assay and the difference in signals as indicated:

5

TABLE H

| Receptor: Gs Fusion Protein | Assay Utilized | Signal Generated: CMV (cpm bound GTP) | Signal Generated: Fusion Protein (cpm bound GTP) | Signal Generated: CMV+ 10µMGDP (cpm bound GTP) | Signal Generated: Fusion Protein + 10µM GDP (cpm bound GTP) | Difference Between: 1. CMV v. Fusion Protein 2. CMV+GDP vs. Fusion+GDP 3. Fusion vs. Fusion+GDP (cpm bound GTP) |
|-----------------------------------|---------------------|---------------------------------------|--|---|---|---|
| hRUP13-Gs | GTPγS (Figure 4) | 32494.0 | 49351.30 | 11148.30 | 28834.67 | 1. 1.5 Fold ← 2. 2.6 Fold ← 3. 42% ⟨ |
| hRUP15-Gs | GTPγS (Figure 8) | 30131.67 | 32493.67 | 7697.00 | 14157.33 | 1. 1.1 Fold ← 2. 1.8 Fold ← 3. 56% ⟨ |

Example 5
FUSION PROTEIN PREPARATION

a. GPCR:Gs Fusion Constuct

The design of the constitutively activated GPCR-G protein fusion construct was accomplished as follows: both the 5' and 3' ends of the rat G protein Gsα (long form; Itoh, H. et al., 83 *PNAS* 3776 (1986)) were engineered to include a HindIII (5'-AAGCTT-3') sequence thereon. Following confirmation of the correct sequence (including the flanking HindIII sequences), the entire sequence was shuttled into pcDNA3.1(-) (Invitrogen, cat. no. V795-20) by subcloning using the HindIII restriction site of that vector. The correct orientation for the Gsα sequence was determined after

10

subcloning into pcDNA3.1(-). The modified pcDNA3.1(-) containing the rat Gsα gene at HindIII sequence was then verified; this vector was now available as a "universal" Gsα protein vector. The pcDNA3.1(-) vector contains a variety of well-known restriction sites upstream of the HindIII site, thus beneficially providing the ability to insert, upstream of the Gs protein, the coding sequence of an endogenous, constitutively active GPCR. This same approach can be utilized to create other "universal" G protein vectors, and, of course, other commercially available or proprietary vectors known to the artisan can be utilized – the important criteria is that the sequence for the GPCR be upstream and in-frame with that of the G protein.

RUP13 couples via Gs. For the following exemplary GPCR Fusion Proteins, fusion to Gsα was accomplished.

A RUP13-Gs α Fusion Protein construct was made as follows: primers were designed as follows:

5'-gatc[TCTAGAAT]GGAGTCCTCACCCATCCCCAG -3' (SEQ.ID.NO.:97; sense)

 $5'-gatc[GATATC]CGTGACTCCAGCCGGGGTGAGGCGGC-3' \ (SEQ.ID.NO.:98; \ antisense).$

Nucleotides in lower caps are included as spacers in the restriction sites (designated in brackets) between the G protein and RUP13. The sense and anti-sense primers included the restriction sites for XbaI and EcoRV, respectively, such that spacers (attributed to the restriction sites) exists between the G protein and RUP15.

PCR was then utilized to secure the respective receptor sequences for fusion within the Gsα universal vector disclosed above, using the following protocol for each: 100ng cDNA for RUP15 was added to separate tubes containing 2μl of each primer (sense and anti-sense), 3μL of 10mM dNTPs, 10μL of 10XTaqPlusTM Precision buffer, 1μL of TaqPlusTM Precision polymerase (Stratagene: #600211), and 80μL of water. Reaction temperatures and cycle times for RUP15 were as follows with cycle steps 2

25

5

10

15

through 4 were repeated 35 times: 94°C for 1 min; 94°C for 30 seconds; 62°C for 20 sec; 72°C 1 min 40sec; and 72° C 5 min . PCR product for was run on a 1% agarose gel and then purified (data not shown). The purified product was digested with XbaI and EcoRV and the desired inserts purified and ligated into the Gs universal vector at the respective restriction site. The positive clones was isolated following transformation and determined by restriction enzyme digest; expression using 293 cells was accomplished following the protocol set forth infra. Each positive clone for RUP15-Gs Fusion Protein was sequenced to verify correctness. (See, SEQ.ID.NO.:99 for nucleic acid sequence and SEQ.ID.NO.:100 for amino acid sequence).

RUP15 couples via Gs. For the following exemplary GPCR Fusion Proteins, fusion to Gsα was accomplished.

A RUP15-Gsα Fusion Protein construct was made as follows: primers were designed as follows:

5'-TCTAGAATGACGTCCACCTGCACCAACAGC-3' (SEQ.ID.NO.:101; sense)

5'-gatatcGCAGGAAAAGTAGCAGAATCGTAGGAAG-3' (SEQ.ID.NO.:102; antisense). 15

Nucleotides in lower caps are included as spacers in the restriction sites between the G protein and RUP15. The sense and anti-sense primers included the restriction sites for EcoRV and Xba1, respectively, such that spacers (attributed to the restriction sites) exists between the G protein and RUP15.

PCR was then utilized to secure the respective receptor sequences for fusion 20 within the Gsa universal vector disclosed above, using the following protocol for each: 100ng cDNA for RUP15 was added to separate tubes containing 2µl of each primer (sense and anti-sense), 3μL of 10mM dNTPs, 10μL of 10XTaqPlus™ Precision buffer, 1 uL of TaqPlus™ Precision polymerase (Stratagene: #600211), and 80μL of water.

Reaction temperatures and cycle times for RUP15 were as follows with cycle steps 2 25

5

through 4 were repeated 35 times: 94°C for 1 min; 94°C for 30 seconds; 62°C for 20 sec; 72°C 1 min 40sec; and 72° C 5 min. PCR product for was run on a 1% agarose gel and then purified (data not shown). The purified product was digested). The purified product was digested with EcoRV and Xba1 and the desired inserts purified and ligated into the Gs universal vector at the respective restriction site. The positive clones was isolated following transformation and determined by restriction enzyme digest; expression using 293 cells was accomplished following the protocol set forth *infra*. Each positive clone for RUP15-Gs Fusion Protein was sequenced to verify correctness. (See, SEQ.ID.NO.:103 for nucleic acid sequence and SEQ.ID.NO.:104 for amino acid sequence).

b. Gq(6 amino acid deletion)/Gi Fusion Construct

The design of a Gq (del)/Gi fusion construct can be accomplished as follows: the N-terminal six (6) amino acids (amino acids 2 through 7, having the sequence of TLESIM (SEQ.ID.NO.: 129) Gαq-subunit will be deleted and the C-terminal five (5) amino acids, having the sequence EYNLV (SEQ.ID.NO.:130) will be replace with the corresponding amino acids of the Gαi Protein, having the sequence DCGLF (SEQ.ID.NO.:131). This fusion construct will be obtained by PCR using the following primers:

5'-gatcaagcttcCATGGCGTGCTGCCTGAGCGAGGAG-3' (SEQ.ID.NO.:132) and

5'-gatcggatccTTAGAACAGGCCGCAGTCCTTCAGGTTCAGCTGCAGGATGGTG-3' (SEQ.ID.NO.:133)

and Plasmid 63313 which contains the mouse Gaq-wild type version with a hemagglutinin tag as template. Nucleotides in lower caps are included as spacers.

TaqPlus Precision DNA polymerase (Stratagene) will be utilized for the amplification by the following cycles, with steps 2 through 4 repeated 35 times: 95°C

5

10

15

20

for 2 min; 95°C for 20 sec; 56°C for 20 sec; 72°C for 2 min; and 72°C for 7 min. The PCR product will be cloned into a pCRII-TOPO vector (Invitrogen) and sequenced using the ABI Big Dye Terminator kit (P.E. Biosystem). Inserts from a TOPO clone containing the sequence of the fusion construct will be shuttled into the expression vector pcDNA3.1(+) at the HindIII/BamHI site by a 2 step cloning process.

Example 6 TISSUE DISTRIBUTION OF THE DISCLOSED HUMAN GPCRS: RT-PCR

RT-PCR was applied to confirm the expression and to determine the tissue distribution of several novel human GPCRs. Oligonucleotides utilized were GPCR-10 specific and the human multiple tissue cDNA panels (MTC, Clontech) as templates. Taq DNA polymerase (Stratagene) were utilized for the amplification in a 40µl reaction according to the manufacturer's instructions. 20µl of the reaction will be loaded on a 1.5% agarose gel to analyze the RT-PCR products. Table J below lists the receptors, the cycle conditions and the primers utizilized.

TABLE J

| Receptor Identifier | Cycle Conditions Min ('), Sec (") Cycles 2-4 repeated 30 times | 5' Primer (SEQ.ID.NO.) | 3' Primer (SEQ.ID.NO.) | DNA Fragment | Tissue Expression |
|------------------------|--|--|---|--------------|---|
| hRUP10 | 94° for 30" 94° for 10" 62°C for 20" 72° for 1' 72° for 7' *cycles 2-4 repeated 35 times | CATGTATGC CAGCGTCCT GCTCC (105) | GCTATGCCTG AAGCCAGTC TTGTG (106) | 730bp | Kidney, leukocyte, liver, placenta and spleen |
| hRUP11 | 94° for 2' 94° for 15" 67°C for 15" 72° for 45" 72° for 5' | GCACCTGCT CCTGAGCAC CTTCTCC (107) | CACAGCGCT GCAGCCCTG CAGCTGGC (108) | 630bp | Liver, kidney, pancreas, colon, small intestinal, spleen and prostate |

5

| hRUP12 | 94° for 2' 94° for 15" 66°C for 15" 72° for 45" 72° for 5' | CCAGTGATG ACTCTGTCC AGCCTG (109) | CAGACACTT GGCAGGGAC GAGGTG (110) | 490bp | Brain, colon, heart, kidney, leukocyte, pancreas, prostate, small intestinal, spleen, testis, and thymus |
|--------|--|--|--|-------|---|
|--------|--|--|--|-------|---|

| hRUP13 | 94° for 1' | CTTGTGGTC | CATATCCCTC | 7001 | |
|----------|-------------------|--------------|--|----------|--------------------|
| | 94° for 15" | ACTGCAGCA | | 700bp | Placenta and |
| | 68°C for 20" | TGTTCCG | AGCGGC (112) | | lung |
| | 72° for 1' 45" | (111) | // // // // // // // // // // // // // | | |
| 1 | 72° for 5' | () | | |] |
| hRUP14 | 94° for 1' | ATGGATCCT | 044040 | | |
| Intel 14 | 94° for 15" | TATCATGGC | CAAGAACAG | 700bp | Not yet |
| | 1 | | GTCTCATCTA | İ | determined |
| | 68°C for 20" | TTCCTC (113) | AGAGCTCC | | |
| | 72° for 1' 45" | Ì | (114) | | |
| hRUP16 | 72° for 5' | | | <u> </u> | |
| nKUPIO | 94° for 30" | CTCTGATGC | GTAGTCCACT | 370bp | Fetal brain, fetal |
| | 94° for 5" | CATCTGCTG | GAAAGTCCA | | kidney and fetal |
| | 69°C for 15" | GATTCCTG | GTGATCC | | skeletal muscle |
| | 72° for 30" | (115) | (116) | | 1 |
| | 72° for 5' | | | | |
| hRUP18 | 94° for 2' | TGGTGGCGA | GTTGCGCCTT | 330bp | Pancreas |
| | 94° for 15" | TGGCCAACA | AGCGACAGA | • | |
| j | 60°C for 20" | GCGCTC (117) | TGACC (118) | 1 | |
| | 72° for 1' | | 1 | | |
| | 72° for 5' | | | | |
| hRUP21 | 94° for 1' | TCAACCTGT | AAGGAGTAG | · | Kidney, lung |
| | 94° for 15" | ATAGCAGCA | CAGAATGGT | | and testis |
| ļ | 56°C for 20" | TCCTC (119) | TAGCC (120) | | and testis |
| | 72° for 40" | | | | |
| | *cycles 2-3 | | | | |
| | repeated 30 times | | | | |
| hRUP22 | 94° for 30" | GACACCTGT | CTGATGGAA | | Testis, thymus |
| I | 94° for 15" | CAGCGGTCG | GTAGAGGCT | | and spleen |
| | 69°C for 20" | TGTGTG (121) | GTCCATCTC | | und spicen |
| | 72° for 40" | | (122) | | |
| | *cycles 2-3 | | | | |
| | repeated 30 times | | | | |
| hRUP23 | 94° for 2' | GCGCTGAGC | CACGGTGAC | 520bp | Placenta |
| | 94° for 15" | GCAGACCAG | GAAGGGCAC | | 1 lacenta |
| | 60°C for 20" | TGGCTG (123) | GAGCTC (124) | | |
| | 72° for 1' | | | | |
| | 72° for 5' | | } | Ì | |
| hRUP26 | 94° for 2' | AGCCATCCC | CCAGGTAGG | 470bp | Pancreas |
| | 94° for 15" | TGCCAGGAA | TGTGCAGCA | 1700p | 1 ancicas |
| | 65°C for 20" | GCATGG (125) | CAATGGC | | ł |
| | 72° for 1' | | (126) | | ł |
| | 72° for 5' | | | | |
| | | | | | ; |
| hRUP27 | 94° for 30" | CTGTTCAAC | ATCATGTCTA | 890bp | Brain |
| | 94° for 10" | AGGGCTGGT | GACTCATGGT | олоор | DIALII |
| i | 55°C for 20" | TGGCAAC | GATCC (128) | | |
| } | 72° for 1' | (127) | - () | | |
| ļ | 72° for 3' | | | | |
| | *cycles 2-4 | | 1 | l | |
| | repeated 35 times | | | | |
| L | | | | | |

Example 7

5

10

15

20

Protocol: Direct Identification of Inverse Agonists and Agonists

A. [35S]GTPγS Assay

Although we have utilized endogenous, constitutively active GPCRs for the direct identification of candidate compounds as, e.g., inverse agonists, for reasons that are not altogether understood, intra-assay variation can become exacerbated. Preferably, then, a GPCR Fusion Protein, as disclosed above, is also utilized with a non-endogenous, constitutively activated GPCR. We have determined that when such a protein is used, intra-assay variation appears to be substantially stabilized, whereby an effective signal-to-noise ratio is obtained. This has the beneficial result of allowing for a more robust identification of candidate compounds. Thus, it is preferred that for direct identification, a GPCR Fusion Protein be used and that when utilized, the following assay protocols be utilized.

1. Membrane Preparation

Membranes comprising the constitutively active orphan GPCR Fusion Protein of interest and for use in the direct identification of candidate compounds as inverse agonists, agonists or partial agonists are preferably prepared as follows:

a. Materials

"Membrane Scrape Buffer" is comprised of 20mM HEPES and 10mM EDTA, pH 7.4; "Membrane Wash Buffer" is comprised of 20 mM HEPES and 0.1 mM EDTA, pH 7.4; "Binding Buffer" is comprised of 20mM HEPES, 100 mM NaCl, and 10 mM MgCl₂, pH 7.4

b. Procedure

All materials will be kept on ice throughout the procedure. Firstly, the media will be aspirated from a confluent monolayer of cells, followed by rinse with 10ml cold

PBS, followed by aspiration. Thereafter, 5ml of Membrane Scrape Buffer will be added to scrape cells; this will be followed by transfer of cellular extract into 50ml centrifuge tubes (centrifuged at 20,000 rpm for 17 minutes at 4°C). Thereafter, the supernatant will be aspirated and the pellet will be resuspended in 30ml Membrane Wash Buffer followed by centrifuge at 20,000 rpm for 17 minutes at 4°C. The supernatant will then be aspirated and the pellet resuspended in Binding Buffer. This will then be homogenized using a Brinkman polytron™ homogenizer (15-20 second bursts until the all material is in suspension). This is referred to herein as "Membrane Protein".

2. **Bradford Protein Assay**

Following the homogenization, protein concentration of the membranes will 10 be determined using the Bradford Protein Assay (protein can be diluted to about 1.5mg/ml, aliquoted and frozen (-80°C) for later use; when frozen, protocol for use will be as follows: on the day of the assay, frozen Membrane Protein is thawed at room temperature, followed by vortex and then homogenized with a polytron at about 12 x 1,000 rpm for about 5-10 seconds; it was noted that for multiple preparations, the homogenizor should be thoroughly cleaned between homoginezation of different preparations).

a. **Materials**

Binding Buffer (as per above); Bradford Dye Reagent; Bradford Protein Standard will be utilized, following manufacturer instructions (Biorad, cat. no. 500-20 0006).

b. Procedure

Duplicate tubes will be prepared, one including the membrane, and one as a control "blank". Each contained 800ul Binding Buffer. Thereafter, 10µl of Bradford Protein Standard (1mg/ml) will be added to each tube, and 10µl of membrane Protein

25

5

will then be added to just one tube (not the blank). Thereafter, 200ul of Bradford Dye Reagent will be added to each tube, followed by vortex of each. After five (5) minutes, the tubes will be re-vortexed and the material therein will be transferred to cuvettes. The cuvettes will then be read using a CECIL 3041 spectrophotometer, at wavelength 595.

3. Direct Identification Assay

a. Materials

GDP Buffer consisted of 37.5 ml Binding Buffer and 2mg GDP (Sigma, cat. no. G-7127), followed by a series of dilutions in Binding Buffer to obtain 0.2 μM GDP (final concentration of GDP in each well was 0.1 μM GDP); each well comprising a candidate compound, has a final volume of 200ul consisting of 100μl GDP Buffer (final concentration, 0.1μM GDP), 50ul Membrane Protein in Binding Buffer, and 50μl [³⁵S]GTPγS (0.6 nM) in Binding Buffer (2.5 μl [³⁵S]GTPγS per 10ml Binding Buffer).

b. Procedure

Candidate compounds will be preferably screened using a 96-well plate format (these can be frozen at -80°C). Membrane Protein (or membranes with expression vector excluding the GPCR Fusion Protein, as control), will be homogenized briefly until in suspension. Protein concentration will then be determined using the Bradford Protein Assay set forth above. Membrane Protein (and control) will then be diluted to 0.25mg/ml in Binding Buffer (final assay concentration, 12.5μg/well). Thereafter, 100 μl GDP Buffer was added to each well of a Wallac ScintistripTM (Wallac). A 5ul pintool will then be used to transfer 5 μl of a candidate compound into such well (*i.e.*, 5μl in total assay volume of 200 μl is a 1:40 ratio such that the final screening concentration of the candidate compound is 10μM). Again, to avoid contamination, after each transfer step the pin tool should be rinsed in three reservoirs comprising water (1X), ethanol (1X)

25

5

10

15

and water (2X) – excess liquid should be shaken from the tool after each rinse and dried with paper and kimwipes. Thereafter, 50 μl of Membrane Protein will be added to each well (a control well comprising membranes without the GPCR Fusion Protein was also utilized), and pre-incubated for 5-10 minutes at room temperature. Thereafter, 50μl of [³⁵S]GTPγS (0.6 nM) in Binding Buffer will be added to each well, followed by incubation on a shaker for 60 minutes at room temperature (again, in this example, plates were covered with foil). The assay will then be stopped by spinning of the plates at 4000 RPM for 15 minutes at 22°C. The plates will then be aspirated with an 8 channel manifold and sealed with plate covers. The plates will then be read on a Wallacc 1450 using setting "Prot. #37" (as per manufacturer instructions).

B. Cyclic AMP Assay

Another assay approach to directly identified candidate compound was accomplished by utilizing a cyclase-based assay. In addition to direct identification, this assay approach can be utilized as an independent approach to provide confirmation of the results from the $[^{35}S]GTP\gamma S$ approach as set forth above.

A modified Flash PlateTM Adenylyl Cyclase kit (New England Nuclear; Cat. No. SMP004A) was preferably utilized for direct identification of candidate compounds as inverse agonists and agonists to constitutively activated orphan GPCRs in accordance with the following protocol.

Transfected cells were harvested approximately three days after transfection. Membranes were prepared by homogenization of suspended cells in buffer containing 20mM HEPES, pH 7.4 and 10mM MgCl₂. Homogenization was performed on ice using a Brinkman Polytron[™] for approximately 10 seconds. The resulting homogenate is centrifuged at 49,000 X g for 15 minutes at 4°C. The resulting pellet was then resuspended in buffer containing 20mM HEPES, pH 7.4 and 0.1 mM EDTA,

25

5

10

15

homogenized for 10 seconds, followed by centrifugation at 49,000 X g for 15 minutes at 4°C. The resulting pellet was then stored at -80°C until utilized. On the day of direct identification screening, the membrane pellet as slowly thawed at room temperature, resuspended in buffer containing 20mM HEPES, pH 7.4 and 10mM MgCL2, to yield a final protein concentration of 0.60mg/ml (the resuspended membranes are placed on ice until use).

cAMP standards and Detection Buffer (comprising 2 μCi of tracer [¹²⁵I cAMP (100 μl] to 11 ml Detection Buffer) were prepared and maintained in accordance with the manufacturer's instructions. Assay Buffer was prepared fresh for screening and contained 20mM HEPES, pH 7.4, 10mM MgCl₂, 20mM phospocreatine (Sigma), 0.1 units/ml creatine phosphokinase (Sigma), 50 μM GTP (Sigma), and 0.2 mM ATP (Sigma); Assay Buffer was then stored on ice until utilized.

Candidate compounds identified as per above (if frozen, thawed at room temperature) were added, preferably, to 96-well plate wells (3µl/well; 12µM final assay concentration), together with 40 µl Membrane Protein (30µg/well) and 50µl of Assay Buffer. This admixture was then incubated for 30 minutes at room temperature, with gentle shaking.

Following the incubation, 100µl of Detection Buffer was added to each well, followed by incubation for 2-24 hours. Plates were then counted in a Wallac MicroBetaTM plate reader using "Prot. #31" (as per manufacturer instructions).

A representative screening assay plate (96 well format) result is presented in Figure 12. Each bar represents the results for a different compound in each well, plus RUP13-Gsα Fusion Protein construct, as prepared in Example 5(a) above. The representative results presented in Figure 12 also provide standard deviations based upon the mean results of each plate ("m") and the mean plus two arbitrary preference for

25

5

10

15

selection of inverse agonists as "leads" from the primary screen involves selection of candidate compounds that that reduce the per cent response by at least the mean plate response, minus two standard deviations. Conversely, an arbitrary preference for selection of an agonists as "leads" from the primary screen involves selection of candidate compounds that increase the per cent response by at least the mean plate response, plus the two standard deviations. Based upon these selection processes, the candidate compounds in the following wells were directly identified as putative inverse agonist (Compound A) and agonist (Compound B) to RUP13 in wells A2 and G9, respectively. See, Figure 12. It is noted for clarity: these compounds have been directly identified without any knowledge of the endogenous ligand for this GPCR. By focusing on assay techniques that are based upon receptor function, and not compound binding affinity, we are able to ascertain compounds that are able to reduce the functional activity of this receptor (Compound A) as well as increase the functional activity of the receptor (Compound B). Based upon the location of these receptor in lung tissue (see, for example, hRUP13 and hRUP21 in Example 6), pharmaceutical agents can be developed for potential therapeutic treatment of lung cancer.

References cited throughout this patent document, including co-pending and related patent applications, unless otherwise indicated, are fully incorporated herein by reference. Modifications and extension of the disclosed inventions that are within the purview of the skilled artisan are encompassed within the above disclosure and the claims that follow.

Although a variety of expression vectors are available to those in the art, for purposes of utilization for both the endogenous and non-endogenous human GPCRs, it is most preferred that the vector utilized be pCMV. This vector was deposited with the American Type Culture Collection (ATCC) on October 13, 1998 (10801 University

25

5

10

15

Blvd., Manassas, VA 20110-2209 USA) under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure. The DNA was tested by the ATCC and determined to be viable. The ATCC has assigned the following deposit number to pCMV: ATCC #203351.

5 //

//

//

//

//

10 //

//

//

//

//

15 //

//

//

//

//

20 //

//

//

//

//

25 //

CLAIMS

What is claimed is:

5

1. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:2.

- 2. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 1.
- 3. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:1.
- 4. A host cell comprising the plasmid of claim 3.
- A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:4.
 - 6. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 5.
 - 7. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:3.
- 15 8. A host cell comprising the plasmid of claim 7.
 - A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:6.
 - 10. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 9.
- 20 11. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:5.
 - 12. A host cell comprising the plasmid of claim 11.
 - 13. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:8.
 - 14. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 13.

15. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:7.

- 16. A host cell comprising the plasmid of claim 15.
- 17. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:10.
- 5 18. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 17.
 - 19. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:9.
 - 20. A host cell comprising the plasmid of claim 19.
 - 21. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:12.
 - 22. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 21 comprising an amino acid sequence of SEQ.ID.NO.84.
 - 23. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:11.
 - 24. A host cell comprising the plasmid of claim 23.
- 25. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:14.
 - 26. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 25 comprising an amino acid sequence of SEQ.ID.NO.88.
 - 27. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:13.
- 20 28. A host cell comprising the plasmid of claim 27.
 - 29. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:16.
 - 30. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 29 comprising an amino acid sequence of SEQ.ID.NO.:92.
- 25 31. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:15.

- 32. A host cell comprising the plasmid of claim 31.
- 33. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:18.
- 34. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 33.
- 35. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:17.
- 36. A host cell comprising the plasmid of claim 35.
- 37. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:20.
- 38. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 37.
 - 39. A plasmid comprising a vector and the cDNA of SE.ID.NO.:19.
 - 40. A host cell comprising the plasmid of claim 39.
 - 41. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:22.
 - 42. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 41.
 - 43. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:21.
 - 44. A host cell comprising the plasmid of claim 43.
- 45. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:24.
 - 46. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 45.
 - 47. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:23.
- 48. A host cell comprising the plasmid of claim 47.

5

49. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:26.

- 50. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 49.
- 5 51. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:25.
 - 52. A host cell comprising the plasmid of claim 51.
 - 53. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:28.
 - 54. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 53.
 - 55. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:27.
 - 56. A host cell comprising the plasmid of claim 55.
 - 57. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:30.
- 15 58. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 57.
 - 59. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:29.
 - 60. A host cell comprising the plasmid of claim 59.
 - 61. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:32.
 - 62. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 61 comprising an amino acid sequence of SEQ.ID.NO.:96.
 - 63. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:95.
 - 64. A host cell comprising the plasmid of claim 63.

20


65. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:34.


- 66. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 65.
- 5 67. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:33.
 - 68. A host cell comprising the plasmid of claim 67.
 - 69. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:36.
 - 70. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 69.
 - 71. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:35.
 - 72. A host cell comprising the plasmid of claim 71.
 - 73. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:38.
- 74. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 73.
 - 75. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:37.
 - 76. A host cell comprising the plasmid of claim 75.
 - 77. A G protein-coupled receptor encoded by an amino acid sequence of SEQ.ID.NO.:40.
 - 78. A non-endogenous, constitutively activated version of the G protein-coupled receptor of claim 77.
 - 79. A plasmid comprising a vector and the cDNA of SEQ.ID.NO.:39.
 - 80. A host cell comprising the plasmid of claim 79.


25


20

10









8XCRE Reporter Assay in 293 Cells




Figure 5







Figure 6


6/12



-igure 7











Figure 11



-igure 12

SEQUENCE LISTING

<110> Arena Pharmaceuticals, Inc. Chen, Rupong Dang, Huong T. Lowitz, Kevin P. <120> Non-Endogenous, Constitutively Activated Human G Protein-Coupled Receptors <130> AREN0087 <150> 60/166,088 <151> 1999-11-17 <150> 60/166,369 1999-11-17 <151> 60/166,099 <150> 1999-11-17 <151> <150> 61/171,902 1999-12-23 <151> <150> 60/171,901 1999-12-23 <151> <150> 60/171,900 1999-12-23 <151> 60/181,749 <150> 2000-02-11 <151> 60/189,258 <150> 2000-03-14 <151> 60/189,259 <150> 2000-03-14 <151> 60/195,899 <150> 2000-04-10 <151> 60/196,078 <150> 2000-04-10 <151> 60/195,898 <150> 2000-04-10 <151> 60/200,419 <150> 2000-04-28 <151> <150> 60/203,630 2000-05-12 <151> 60/210,741 <150> 2000-06-12 <151> <150> 60/210,982 2000-06-12 <151> 60/226,760 <150> 2000-08-21 <151> <150> 60/235,779 <151> 2000-09-26

Page 1

```
60/235,418
 <150>
 <151>
       2000-09-26
 <150> 60/242,332
 <151> 2000-10-20
 <150> 60/242,343
 <151> 2000-10-20
<150> 60/243,019
<151> 2000-10-24
<160> 133
<170> PatentIn version 3.0
<210>
<211>
       1155
      DNA
<212>
<213> Homo sapiens
<400> 1
atggcagccc agaatggaaa caccagtttc acacccaact ttaatccacc ccaagaccat
                                                                       60
gcctcctccc tctcctttaa cttcagttat ggtgattatg acctccctat ggatgaggat
                                                                      120
gaggacatga ccaagacccg gaccttcttc gcagccaaga tcgtcattgg cattgcactg
                                                                      180
gcaggcatca tgctggtctg cggcatcggt aactttgtct ttatcgctgc cctcacccgc
                                                                      240
tataagaagt tgcgcaacct caccaatctg ctcattgcca acctggccat ctccgacttc
                                                                      300
ctggtggcca tcatctgctg ccccttcgag atggactact acgtggtacg gcagctctcc
                                                                      360
tgggagcatg gccacgtgct ctgtgcctcc gtcaactacc tgcgcaccgt ctccctctac
                                                                      420
gtctccacca atgccttgct ggccattgcc attgacagat atctcgccat cgttcacccc
                                                                      480
ttgaaaccac ggatgaatta tcaaacggcc tccttcctga tcgccttggt ctgjatggtg
                                                                     540
tccattctca ttgccatccc atcggcttac tttgcaacag aaacggtcct ctttattgtc
                                                                     600
aagagccagg agaagatett etgtggccag atetggeetg tggatcagea getetaetae
                                                                     660
aagteetact teetetteat etttggtgte gagttegtgg geeetgtggt caccatgace
                                                                     720
ctgtgctatg ccaggatctc ccgggagctc tggttcaagg cagtccctgg gttccagacg
                                                                     780
gagcagattc gcaagcggct gcgctgccgc aggaagacgg tcctggtgct catgtgcatt
                                                                     840
ctcacggcct atgtgctgtg ctgggcaccc ttctacggtt tcaccatcgt tcgtgacttc
                                                                     900
ttccccactg tgttcgtgaa ggaaaagcac tacctcactg ccttctacgt ggtcgagtgc
                                                                     960
atcgccatga gcaacagcat gatcaacacc gtgtgcttcg tgacggtcaa gaacaacacc
                                                                    1020
atgaagtact tcaagaagat gatgctgctg cactggcgtc cctcccagcg ggggagcaag
                                                                    1080
tccagtgctg accttgacct cagaaccaac ggggtgccca ccacagaaga ggtggactgt
                                                                    1140
atcaggctga agtga
                                                                    1155
```

<210> 2 <211> 384

Page 2

<212> PRT

<213> Homo sapiens

<400> 2

Met Ala Ala Gln Asn Gly Asn Thr Ser Phe Thr Pro Asn Phe Asn Pro 1 5 10 15

Pro Gln Asp His Ala Ser Ser Leu Ser Phe Asn Phe Ser Tyr Gly Asp 20 25 30

Tyr Asp Leu Pro Met Asp Glu Asp Glu Asp Met Thr Lys Thr Arg Thr 35 40 45

Phe Phe Ala Ala Lys Ile Val Ile Gly Ile Ala Leu Ala Gly Ile Met 50 55 60

Leu Val Cys Gly Ile Gly Asn Phe Val Phe Ile Ala Ala Leu Thr Arg 70 75 80

Tyr Lys Lys Leu Arg Asn Leu Thr Asn Leu Leu Ile Ala Asn Leu Ala 85 90 95

Ile Ser Asp Phe Leu Val Ala Ile Ile Cys Cys Pro Phe Glu Met Asp 100 105 110

Tyr Tyr Val Val Arg Gln Leu Ser Trp Glu His Gly His Val Leu Cys 115 120 125

Ala Ser Val Asn Tyr Leu Arg Thr Val Ser Leu Tyr Val Ser Thr Asn

Ala Leu Leu Ala Ile Ala Ile Asp Arg Tyr Leu Ala Ile Val His Pro 145 150 160

Leu Lys Pro Arg Met Asn Tyr Gln Thr Ala Ser Phe Leu Ile Ala Leu 165 170 175

Val Trp Met Val Ser Ile Leu Ile Ala Ile Pro Ser Ala Tyr Phe Ala 180 185 190

Thr Glu Thr Val Leu Phe Ile Val Lys Ser Gln Glu Lys Ile Phe Cys 195 200 205

Gly Gln Ile Trp Pro Val Asp Gln Gln Leu Tyr Tyr Lys Ser Tyr Phe 210 215 220

Leu Phe Ile Phe Gly Val Glu Phe Val Gly Pro Val Val Thr Met Thr 225 230 235 235

Leu Cys Tyr Ala Arg Ile Ser Arg Glu Leu Trp Phe Lys Ala Val Pro 245 250 255

Gly Phe Gln Thr Glu Gln Ile Arg Lys Arg Leu Arg Cys Arg Arg Lys 260 265 270

Thr Val Leu Val Leu Met Cys Ile Leu Thr Ala Tyr Val Leu Cys Trp 275 280 285

Ala Pro Phe Tyr Gly Phe Thr Ile Val Arg Asp Phe Phe Pro Thr Val 290 295 300

Phe Val Lys Glu Lys His Tyr Leu Thr Ala Phe Tyr Val Val Glu Cys 305 310 315 320

Ile Ala Met Ser Asn Ser Met Ile Asn Thr Val Cys Phe Val Thr Val Lys Asn Asn Thr Met Lys Tyr Phe Lys Lys Met Met Leu Leu His Trp Arg Pro Ser Gln Arg Gly Ser Lys Ser Ser Ala Asp Leu Asp Leu Arg Thr Asn Gly Val Pro Thr Thr Glu Glu Val Asp Cys Ile Arg Leu Lys 370 <210> 3 <211> 1260 <212> DNA <213> Homo sapiens <400> 3 atgotggcag ctgcctttgc agactotaac tocagcagca tgaatgtgtc otttgctcac 60 ctccactttg ccggagggta cctgccctct gattcccagg actggagaac catcatcccg 120 gctctcttgg tggctgtctg cctggtgggc ttcgtgggaa acctgtgtgt gattggcatc 180 ctccttcaca atgcttggaa aggaaagcca tccatgatcc actccctgat tctgaatctc 240 agectggetg atetetecet cetgetgttt tetgeaceta teegagetae ggegtaetee 300 aaaagtgttt gggatctagg ctggtttgtc tgcaagtcct ctgactggtt tatccacaca 360 tgcatggcag ccaagagcct gacaatcgtt gtggtggcca aagtatgctt catgtatgca 420 agtgacccag ccaagcaagt gagtatccac aactacacca tctggtcagt gctggtggcc 480 atctggactg tggctagcct gttacccctg ccggaatggt tctttagcac catcaggcat 540 catgaaggtg tggaaatgtg cctcgtggat gtaccagctg tggctgaaga gtttatgtcg 600 atgtttggta agctctaccc actcctggca tttggccttc cattattttt tgccagcttt 660 tatttctgga gagcttatga ccaatgtaaa aaacgaggaa ctaagactca aaatcttaga 720 aaccagatac gctcaaagca agtcacagtg atgctgctga gcattgccat catctctgct 780 ctcttgtggc tccccgaatg ggtagcttgg ctgtgggtat ggcatctgda ggctgcaggc 840 ccggccccac cacaaggttt catagccctg tctcaagtct tgatgttttc catctcttca 900 gcaaatcctc tcatttttct tgtgatgtcg gaagagttca gggaaggctt gaaaggtgta 960 tggaaatgga tgataaccaa aaaacctcca actgtctcag agtctcagga aacaccagct 1020 ggcaactcag agggtcttcc tgacaaggtt ccatctccag aatccccagc atccatacca 1080 gaaaaagaga aacccagctc tecetectet ggcaaaggga aaactgagaa ggcagagatt 1140 cccatccttc ctgacgtaga gcagttttgg catgagaggg acacagtccc ttctgtacag 1200

gacaatgacc ctatcccctg ggaacatgaa gatcaagaga caggggaagg tgttaaatag

<210> 4 <211> 419 <212> PRT <213> Homo sapiens

Page 4

1260

<400> 4

Met Leu Ala Ala Ala Phe Ala Asp Ser Asn Ser Ser Ser Met Asn Val Ser Phe Ala His Leu His Phe Ala Gly Gly Tyr Leu Pro Ser Asp Ser 20 25 30Gln Asp Trp Arg Thr Ile Ile Pro Ala Leu Leu Val Ala Val Cys Leu Val Gly Phe Val Gly Asn Leu Cys Val Ile Gly Ile Leu Leu His Asn 50 55 Ala Trp Lys Gly Lys Pro Ser Met Ile His Ser Leu Ile Leu Asn Leu 65 70 75 80 Ser Leu Ala Asp Leu Ser Leu Leu Leu Phe Ser Ala Pro Ile Arg Ala Thr Ala Tyr Ser Lys Ser Val Trp Asp Leu Gly Trp Phe Val Cys Lys Ser Ser Asp Trp Phe Ile His Thr Cys Met Ala Ala Lys Ser Leu Thr Ile Val Val Val Ala Lys Val Cys Phe Met Tyr Ala Ser Asp Pro Ala Lys Gln Val Ser Ile His Asn Tyr Thr Ile Trp Ser Val Leu Val Ala Ile Trp Thr Val Ala Ser Leu Leu Pro Leu Pro Glu Trp Phe Phe Ser Thr Ile Arg His His Glu Gly Val Glu Met Cys Leu Val Asp Val Pro Ala Val Ala Glu Glu Phe Met Ser Met Phe Gly Lys Leu Tyr Pro Leu 200 Leu Ala Phe Gly Leu Pro Leu Phe Phe Ala Ser Phe Tyr Phe Trp Arg Ala Tyr Asp Gln Cys Lys Lys Arg Gly Thr Lys Thr Gln Asn Leu Arg 225 230 235 240 Asn Gln Ile Arg Ser Lys Gln Val Thr Val Met Leu Leu Ser Ile Ala Ile Ile Ser Ala Leu Leu Trp Leu Pro Glu Trp Val Ala Trp Leu Trp Val Trp His Leu Lys Ala Ala Gly Pro Ala Pro Pro Gln Gly Phe Ile Ala Leu Ser Gln Val Leu Met Phe Ser Ile Ser Ser Ala Asn Pro Leu Ile Phe Leu Val Met Ser Glu Glu Phe Arg Glu Gly Leu Lys Gly Val 315 Trp Lys Trp Met Ile Thr Lys Lys Pro Pro Thr Val Ser Glu Ser Gln

```
Glu Thr Pro Ala Gly Asn Ser Glu Gly Leu Pro Asp Lys Val Pro Ser
  Pro Glu Ser Pro Ala Ser Ile Pro Glu Lys Glu Lys Pro Ser Ser Pro
  Ser Ser Gly Lys Gly Lys Thr Glu Lys Ala Glu Ile Pro Ile Leu Pro
                          375
 Asp Val Glu Gln Phe Trp His Glu Arg Asp Thr Val Pro Ser Val Gln
 Asp Asn Asp Pro Ile Pro Trp Glu His Glu Asp Gln Glu Thr Gly Glu
                                      410
 Gly Val Lys
 <210>
        5
 <211>
        1014
 <212>
        DNA
 <213>
        Homo sapiens
 <400> 5
 atggggaacg attctgtcag ctacgagtat ggggattaca gcgacctctc ggaccgccct
                                                                        60
 gtggactgcc tggatggcgc ctgcctggcc atcgacccgc tgcgcgtggc cccgctccca
                                                                       120
 ctgtatgccg ccatcttcct ggtgggggtg ccgggcaatg ccatggtggc ctgggtggct
                                                                       180
gggaaggtgg cccgccggag ggtgggtgcc acctggttgc tccacctggc cgtggcggat
                                                                       240
ttgctgtgct gtttgtctct gcccatcctg gcagtgccca ttgcccgtgg aggccactgg
                                                                       300
ccgtatggtg cagtgggctg tcgggcgctg ccctccatca tcctgctgac catgtatgcc
                                                                       360
agegteetge teetggeage teteagtgee gaeetetget teetggetet egggeetgee
                                                                       420
tggtggtcta cggttcagcg ggcgtgcggg gtgcaggtgg cctgtggggc agcctggaca
                                                                       480
ctggccttgc tgctcaccgt gccctccgcc atctaccgcc ggctgcacca ggagcacttc
                                                                       540
ccagcccggc tgcagtgtgt ggtggactac ggcggctcct ccagcaccga gaatgcggtg
                                                                       600
actgccatcc ggtttctttt tggcttcctg gggcccctgg tggccgtggc cagctgccac
                                                                       660
agtgccctcc tgtgctgggc agcccgacgc tgccggccgc tgggcacagc cattgtggtg
                                                                      720
gggttttttg tctgctgggc accctaccac ctgctggggc tggtgctcac tgtggcggcc
                                                                      780
ccgaactccg cactcctggc cagggccctg cgggctgaac ccctcatcgt gggccttgcc
                                                                      840
ctegeteaca getgeeteaa teecatgete tteetgtatt ttgggaggge teaacteege
                                                                      900
cggtcactgc cagctgcctg tcactgggcc ctgagggagt cccagggcca ggacgaaagt
                                                                      960
gtggacagca agaaatccac cagccatgac ctggtctcgg agatggaggt gtag
                                                                     1014
<210>
       6
```

Page 6

<211>

337 <212> PRT

<213> Homo sapiens

<400> 6

Met Gly Asn Asp Ser Val Ser Tyr Glu Tyr Gly Asp Tyr Ser Asp Leu
1 5 10 15 Ser Asp Arg Pro Val Asp Cys Leu Asp Gly Ala Cys Leu Ala Ile Asp Pro Leu Arg Val Ala Pro Leu Pro Leu Tyr Ala Ala Ile Phe Leu Val Gly Val Pro Gly Asn Ala Met Val Ala Trp Val Ala Gly Lys Val Ala Arg Arg Arg Val Gly Ala Thr Trp Leu Leu His Leu Ala Val Ala Asp
70 75 80 Leu Leu Cys Cys Leu Ser Leu Pro Ile Leu Ala Val Pro Ile Ala Arg Gly Gly His Trp Pro Tyr Gly Ala Val Gly Cys Arg Ala Leu Pro Ser Ile Ile Leu Leu Thr Met Tyr Ala Ser Val Leu Leu Ala Ala Leu Ser Ala Asp Leu Cys Phe Leu Ala Leu Gly Pro Ala Trp Trp Ser Thr Val Gln Arg Ala Cys Gly Val Gln Val Ala Cys Gly Ala Ala Trp Thr 145 150 155 Leu Ala Leu Leu Thr Val Pro Ser Ala Ile Tyr Arg Arg Leu His Gln Glu His Phe Pro Ala Arg Leu Gln Cys Val Val Asp Tyr Gly Gly Ser Ser Ser Thr Glu Asn Ala Val Thr Ala Ile Arg Phe Leu Phe Gly 200 Phe Leu Gly Pro Leu Val Ala Val Ala Ser Cys His Ser Ala Leu Leu Cys Trp Ala Ala Arg Arg Cys Arg Pro Leu Gly Thr Ala Fle Val Val 225 230 235 240 Gly Phe Phe Val Cys Trp Ala Pro Tyr His Leu Leu Gly Leu Val Leu

Met Leu Phe Leu Tyr Phe Gly Arg Ala Gln Leu Arg Arg Ser Leu Pro 290 295 300

Ala Ala Cys His Trp Ala Leu Arg Glu Ser Gln Gly Gln Asp Glu Ser 305 310 315 320

Thr Val Ala Ala Pro Asn Ser Ala Leu Leu Ala Arg Ala Leu Arg Ala

Glu Pro Leu Ile Val Gly Leu Ala Leu Ala His Ser Cys Leu Asn Pro

Val Asp Ser Lys Lys Ser Thr Ser His Asp Leu Val Ser Glu Met Glu 325 330 335

Val

<210>

7

```
<211>
        1272
 <212>
        DNA
 <213> Homo sapiens
 <400> 7
 atgttgtgtc accgtggtgg ccagctgata gtgccaatca tcccactttg ccctgagcac
                                                                        60
 tcctgcaggg gtagaagact ccagaacctt ctctcaggcc catggcccaa gcagcccatg
                                                                       120
 gaacttcata acctgagete tecatetece teteteteet cetetgttet ecetecetee
                                                                       180
 ttctctccct caccctcctc tgctccctct gcctttacca ctgtgggggg gtcctctgga
                                                                       240
 gggccctgcc accccacctc ttcctcgctg gtgtctgcct tcctggcacc aatcctggcc
                                                                       300
 ctggagtttg tcctgggcct ggtggggaac agtttggccc tcttcatctt ctgcatccac
                                                                       360
 acgcggccct ggacctccaa cacggtgttc ctggtcagcc tggtggccgc tgacttcctc
                                                                       420
ctgatcagca acctgcccct ccgcgtggac tactacctcc tccatgagac ctggcgcttt
                                                                       480
ggggctgctg cctgcaaagt caacctcttc atgctgtcca ccaaccgcac ggccagcgtt
                                                                       540
gtcttcctca cagccatcgc actcaaccgc tacctgaagg tggtgcagcc ccaccacgtg
                                                                      600
ctgagccgtg cttccgtggg ggcagctgcc cgggtggccg ggggactctg ggtgggcatc
                                                                      660
ctgctcctca acgggcacct gctcctgagc accttctccg gcccctcctg cctcagctac
                                                                      720
agggtgggca cgaagccctc ggcctcgctc cgctggcacc aggcactgta cctgctggag
                                                                      780
ttetteetge cactggeget cateetettt getattgtga geattggget caccateegg
                                                                      840
aaccgtggtc tgggcgggca ggcaggcccg cagagggcca tgcgtgtgct ggccatggtg
                                                                      900
gtggccgtct acaccatctg cttcttgccc agcatcatct ttggcatggc ttccatggtg
                                                                      960
getttetgge tgteegeetg ecgatecetg gaeetetgea caeagetett ecatggetee
                                                                     1020
ctggccttca cctacctcaa cagtgtcctg gaccccgtgc tctactgctt ctctagcccc
                                                                     1080
aactteetee accagageeg ggeettgetg ggeeteaege ggggeeggea gggeeeagtg
                                                                     1140
agcgacgaga gctcctacca accctccagg cagtggcgct accgggaggc ctctaggaag
                                                                     1200
gcggaggcca tagggaagct gaaagtgcag ggcgaggtct ctctggaaaa ggaaggctcc
                                                                    1260
tcccagggct ga
                                                                     1272
```

<210> 8 <211> 423 <212> PRT <213> Homo sapiens <400> 8

Met Leu Cys His Arg Gly Gly Gln Leu Ile Val Pro Ile Ile Pro Leu 1 5 10 15

Cys Pro Glu His Ser Cys Arg Gly Arg Arg Leu Gln Asn Leu Leu Ser Page 8

| | | | 20 | | | | | 25 | | | | | 30 | | |
|------------|------------|------------|--------------|------------|------------|------------|--------------|------------|------------|------------|------------|------------|------------|------------|------------|
| Gly | Pro | Trp 35 | Pro | Lys | Gln | Pro | Met 40 | Glu | Leu | His | Asn | Leu 45 | Ser | Ser | Pro |
| Ser | Pro 50 | Ser | Leu | Ser | Ser | Ser 55 | Val | Leu | Pro | Pro | Ser 60 | Phe | Ser | Pro | Ser |
| Pro 65 | Ser | Ser | Ala | Pro | Ser 70 | Ala | Phe | Thr | Thr | Val 75 | Gly | Gly | Ser | Ser | Gly 80 |
| Gly | Pro | Cys | His | Pro 85 | Thr | Ser | Ser | Ser | Leu 90 | Val | Ser | Ala | Phe | Leu 95 | Ala |
| Pro | Ile | Leu | Ala 100 | Leu | Glu | Phe | Val | Leu 105 | Gly | Leu | Val | Gly | Asn 110 | Ser | Leu |
| Ala | Leu | Phe 115 | Ile | Phe | Cys | Ile | His 120 | Thr | Arg | Pro | Trp | Thr 125 | Ser | Asn | Thr |
| Val | Phe 130 | Leu | Val | Ser | Leu | Val 135 | Ala | Ala | Asp | Phe | Leu 140 | Leu | Ile | Ser | Asn |
| Leu 145 | Pro | Leu | Arg | Val | Asp 150 | Tyr | Tyr | Leu | Leu | His 155 | Glu | Thr | Trp | Arg | Phe 160 |
| Gly | Ala | Ala | Ala | Cys 165 | Lys | Val | Asn | Leu | Phe 170 | Met | Leu | Ser | Thr | Asn 175 | Arg |
| Thr | Ala | Ser | Val 180 | Val | Phe | Leu | Thr | Ala 185 | Ile | Ala | Leu | Asn | Arg 190 | Tyr | Leu |
| Lys | Val | Val 195 | Gln | Pro | His | His | Val 200 | Leu | Ser | Arg | Ala | Ser 205 | Val | Gly | Ala |
| Ala | Ala 210 | Arg | Val | Ala | Gly | Gly 215 | Leu | Trp | Val | Gly | 11e 220 | Leu | Leu | Leu | Asn |
| Gly 225 | His | Leu | Leu | Leu | Ser 230 | | Phe | Ser | Gly | Pro 235 | Ser | Суѕ | Leu | Ser | Tyr 240 |
| Arg | Val | Gly | Thr | Lys 245 | Pro | Ser | Ala | Ser | Leu 250 | Arg | Trp | His | Gln | Ala 255 | Leu |
| Tyr | Leu | Leu | Glu 260 | | Phe | Leu | Pro | Leu 265 | Ala | Leu | Ile | Leu | Phe 270 | Ala | Ile |
| Val | Ser | 11e 275 | | Leu | Thr | Ile | 280 | Asn | Arg | Gly | Leu | Gly 285 | Gly | Gln | Ala |
| Gly | Pro 290 | | Arg | Ala | Met | Arg 295 | Val | Leu | Ala | Met | Val 300 | Val | Ala | Val | Tyr |
| Thr 305 | | Cys | Phe | Leu | 310 | | Ile | Ile | Phe | Gly 315 | Met | Ala | Ser | Met | Val 320 |
| Ala | Phe | Trp | Leu | 325 | Ala | Cys | s Arg | Ser | 330 | ı Asp | Leu | Cys | Thr | Gln 335 | Leu |
| Phe | His | Gly | / Ser 340 | | n Ala | Phe | e Thr | Tyr 345 | Leu b | ı Ası | ser | Val | Leu 350 | Asp | Pro |
| Val | . Let | ту: 359 | | s Phe | e Ser | Ser | 7 Pro 360 | Asr | n Phe | e Lev | ı His | 365 | Ser | Arg | Ala |
| | | | | | | | | | | | | | | | |

Page 9

Leu Leu Gly Leu Thr Arg Gly Arg Gln Gly Pro Val Ser Asp Glu Ser Ser Tyr Gln Pro Ser Arg Gln Trp Arg Tyr Arg Glu Ala Ser Arg Lys Ala Glu Ala Ile Gly Lys Leu Lys Val Gln Gly Glu Val Ser Leu Glu Lys Glu Gly Ser Ser Gln Gly 420 <210> 9 966 <211> <212> DNA <213> Homo sapiens <400> atgaaccaga ctttgaatag cagtgggacc gtggagtcag ccctaaacta ttccagaggg 60 agcacagtgc acacggccta cctggtgctg agctccctgg ccatgttcac ctgcctgtgc 120 gggatggcag gcaacagcat ggtgatctgg ctgctgggct ttcgaatgca caggaacccc 180 ttctgcatct atatcctcaa cctggcggca gccgacctcc tcttcctctt cagcatggct 240 tocacgetca geotogaaac ecageceetg gteaatacea etgacaaggt ecaegagetg 300 atgaagagac tgatgtactt tgcctacaca gtgggcctga gcctgctgac ggccatcagc 360 acceageget gtetetetgt cetetteeet atetggttea agtgteaceg geeeaggeac 420 ctgtcagect gggtgtgtgg cetgetgtgg acaetetgte teetgatgaa egggttgaee 480 tetteettet geageaagtt ettgaaatte aatgaagate ggtgetteag ggtggaeatg 540 gtccaggccg ccctcatcat gggggtctta accccagtga tgactctgtc cagcctgacc 600 ctetttgtet gggtgcggag gageteceag cagtggcgge ggcageeeac aeggetgtte 660 gtggtggtcc tggcctctgt cctggtgttc ctcatctgtt ccctgcctct gagcatctac 720 tggtttgtgc tctactggtt gagcctgccg cccgagatgc aggtcctgtg cttcagcttg 780 tcacgcctct cctcgtccgt aagcagcagc gccaaccccg tcatctactt cctggtgggc 840 agccggagga gccacaggct gcccaccagg tccctgggga ctgtgctcca acaggcgctt 900 cgcgaggagc ccgagctgga aggtggggag acgcccaccg tgggcaccaa tgagatgggg 960 gcttga 966 <210> 10 321 <212> PRT <213> Homo sapiens

<211>

<400> 10

Met Asn Gln Thr Leu Asn Ser Ser Gly Thr Val Glu Ser Ala Leu Asn

Tyr Ser Arg Gly Ser Thr Val His Thr Ala Tyr Leu Val Leu Ser Ser 20 25 30

| Leu . | Ala | Met 35 | Phe | Thr | Суѕ | Leu | Cys 40 | Gly | Met | Ala | Gly | Asn 45 | Ser | Met | Val |
|--------------------------|------------|---------------------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| | Trp 50 | Leu | Leu | Gly | Phe | Arg 55 | Met | His | Arg | Asn | Pro 60 | Phe | Cys | Ile | Tyr |
| Ile 65 | Leu | Asn | Leu | Ala | Ala 70 | Ala | Asp | Leu | Leu | Phe 75 | Leu | Phe | Ser | Met | Ala 80 |
| Ser | Thr | Leu | Ser | Leu 85 | Glu | Thr | Gln | Pro | Leu 90 | Val | Asn | Thr | Thr | Asp 95 | Lys |
| Val | His | Glu | Leu 100 | Met | Lys | Arg | Leu | Met 105 | Tyr | Phe | Ala | Tyr | Thr 110 | Val | Gly |
| Leu | Ser | Leu 115 | Leu | Thr | Ala | Ile | Ser 120 | Thr | Gln | Arg | Суз | Leu 125 | Ser | Val | Leu |
| Phe | Pro 130 | Ile | Trp | Phe | Lys | Cys 135 | His | Arg | Pro | Arg | His 140 | Leu | Ser | Ala | Trp |
| Val 145 | Cys | Gly | Leu | Leu | Trp 150 | Thr | Leu | Cys | Leu | Leu 155 | Met | Asn | Gly | Leu | Thr 160 |
| Ser | Ser | Phe | Cys | Ser 165 | Lys | Phe | Leu | Lys | Phe 170 | Asn | Glu | Asp | Arg | Cys 175 | Phe |
| | | | 180 | | | | | 185 | | | | | Leu 190 | | |
| | | 195 | | | | | 200 | | | | | 205 | Arg | | |
| | 210 | | | | | 215 | | | | | 220 | | Val | | |
| 225 | | | | | 230 | | | | | 235 | | | Ser | | 240 |
| | | | | 245 | | | | | 250 | | | | Gln | 255 | |
| - | | | 260 | | | | | 265 | | | | | 2-70 | | Asn |
| | | 275 | | | | | 280 | | | | | 285 | | | Pro |
| | 290 |) | | | | 295 | | | | | 300 | | | | Pro |
| Glu 305 | Leu | ı Glu | Gly | Gly | Glu 310 | Thr | Pro | Thr | Val | Gly 315 | Thr | Asn | Glu | Met | Gly 320 |
| Ala | | | | | | | | | | | | | | | |
| <21 <21 <21 <21 | 1> 2> | 11 1356 DNA Homo | | oiens | 5 | | | | | | | | | | |
| <40 | 0> | 11 | | | | | casto | ~ | 70220 | -t-ct1 | | actti | aaa | gagg | gtccct |
| atg | gagī | LCCT | cacc | Juan | (| .cay | | ou gy | ,,,,,,,, | | | e 11 | 223 | , , | |

60

| caaacccca | g gtccctcta | c tgccagtgg | g gtcccggagg | tggggctace | g ggatgttgct | 120 |
|------------|-------------|-------------|--------------|------------|--------------|------|
| | | | | | ggctggcaat | 180 |
| | | | | | cttcgtcttc | 240 |
| | | | | | : catgctctcc | 300 |
| | | | ggggaggtgg | | | 360 |
| | | | ctctcggtgt | | | 420 |
| | | | gaggtgcgca | | | 480 |
| | | | ttggccatgg | | | 540 |
| | | | gtcccccag | | | 600 |
| | | | gtctttgctg | | | 660 |
| | | | atgttccgag | | | 720 |
| | | | acaccccggc | | | 780 |
| | | | gccccccaga | | | 840 |
| | | | gctgtggggg | | | 900 |
| | | | gccctgagtg | | | 960 |
| | | | tacttttgct | | | 1020 |
| tatggatgtc | tcaaccggca | gatccggggg | gageteagea | agcagtttgt | ctgcttcttc | 1080 |
| aagccagctc | cagaggagga | gctgaggctg | cctagccggg | agggctccat | tgaggagaac | 1140 |
| ttcctgcagt | tccttcaggg | gactggctgt | ccttctgagt | cctgggtttc | ccgaccccta | 1200 |
| ccagcccca | agcaggagcc | acctgctgtt | gactttcgaa | tcccaggcca | gatagctgag | 1260 |
| gagacctctg | agttcctgga | gcagcaactc | accagcgaca | tcatcatgtc | agacagctac | 1320 |
| ctccgtcctg | ccgcctcacc | ccggctggag | tcatga | | | 1356 |

<210> 12 <211> 451

<212> PRT

<213> Homo sapiens

<400> 12

Met Glu Ser Ser Pro Ile Pro Gln Ser Ser Gly Asn Ser Ser Thr Leu 1 5 10 15

Gly Arg Val Pro Gln Thr Pro Gly Pro Ser Thr Ala Ser Gly Val Pro 20 25 30

Glu Val Gly Leu Arg Asp Val Ala Ser Glu Ser Val Ala Leu Phe Phe 35 40 45

Met Leu Leu Asp Leu Thr Ala Val Ala Gly Asn Ala Ala Val Met 50 60

| Ala 65 | Val | Ile | Ala | Lys | Thr 70 | Pro | Ala | Leu | Arg | Lys 75 | Phe | Val | Phe | Val | Phe 80 |
|------------|--------------|------------|------------|------------------|--------------|------------|--------------|------------|--------------|------------|------------|------------|------------|------------|------------|
| His | Leu | Cys | Leu | Val 85 | Asp | Leu | Leu | Ala | Ala 90 | Leu | Thr | Leu | Met | Pro 95 | Leu |
| Ala | Met | Leu | Ser 100 | Ser | Ser | Ala | Leu | Phe 105 | Asp | His | Ala | Leu | Phe 110 | Gly | Glu |
| Val | Ala | Cys 115 | Arg | Leu | Tyr | Leu | Phe 120 | Leu | Ser | Val | Cys | Phe 125 | Val | Ser | Leu |
| Ala | Ile 130 | Leu | Ser | Val | Ser | Ala 135 | Ile | Asn | Val | Glu | Arg 140 | Tyr | Tyr | Tyr | Val |
| Val 145 | His | Pro | Met | Arg | Tyr 150 | Glu | Val | Arg | Met | Thr 155 | Leu | Gly | Leu | Val | Ala 160 |
| Ser | Val | Leu | Val | Gly 165 | Val | Trp | Val | Lys | Ala 170 | Leu | Ala | Met | Ala | Ser 175 | Val |
| Pro | Val | Leu | Gly 180 | Arg | Val | Ser | Trp | Glu 185 | Glu | Gly | Ala | Pro | Ser 190 | Val | Pro |
| Pro | Gly | Cys 195 | Ser | Leu | Gln | Trp | Ser 200 | His | Ser | Ala | Туr | Cys 205 | Gln | Leu | Phe |
| Val | Val 210 | Val | Phe | Ala | Val | Leu 215 | Tyr | Phe | Leu | Leu | Pro 220 | Leu | Leu | Leu | Ile |
| Leu 225 | | Val | Tyr | Cys | Ser 230 | Met | Phe | Arg | Val | Ala 235 | Arg | Val | Ala | Ala | Met 240 |
| Gln | His | Gly | Pro | Leu 245 | Pro | Thr | Trp | Met | Glu 250 | Thr | Pro | Arg | Gln | Arg 255 | Ser |
| Glu | Ser | Leu | Ser 260 | | Arg | Ser | Thr | Met 265 | Val | Thr | Ser | Ser | Gly 270 | Ala | Pro |
| Gln | Thr | Thr 275 | | His | Arg | Thr | Phe 280 | Gly | Gly | Gly | Lys | Ala 285 | Ala | Val | Val |
| Leu | Leu 290 | | Val | Gly | Gly | Gln 295 | | Leu | Leu | Суѕ | Trp 300 | Leu | Pro | Tyr | Phe |
| Ser 305 | | His | Leu | Tyr | Val 310 | | Leu | Ser | Ala | Gln 315 | Pro | Ile | Ser | Thr | Gly 320 |
| Glr | Val | . Glu | Ser | Val 325 | Val | Thr | Trp | Ile | Gly 330 | Tyr | Phe | Cys | Phe | Thr 335 | Ser |
| Asr | Pro | Phe | Phe 340 | | Gly | Cys | Leu | Asr 345 | a Arg | , Gln | Ile | Arg | Gly 350 | Glu | Leu |
| Sei | Lys | 355 | | · Val | Cys | Phe | 2 Phe 360 | Lys | Pro | Ala | Pro | Glu 365 | Glu | Glu | Leu |
| Arg | g Lei 370 | | Ser | Arg | g Glu | Gly 375 | y Ser | : Ile | e Glu | ı Glu | Asn 380 | Phe | Leu | Gln | Phe |
| Le: | | n Gly | y Thi | Gly | 7 Cys 390 | | Se 1 | Glu | ı Sei | Trp 395 | Val | . Ser | Arç | Pro | Leu 400 |
| Pr | o Se | r Pro | b Lys | 3 Glr 40 | | ı Pro | o Pro | Ala | a Vai 410 | l Asp | Phe | e Arg | ıle | Pro 415 | Gly |
| | | | | | | | | | | | Page | e 13 | | | |

Gln Ile Ala Glu Glu Thr Ser Glu Phe Leu Glu Gln Gln Leu Thr Ser Asp Ile Ile Met Ser Asp Ser Tyr Leu Arg Pro Ala Ala Ser Pro Arg 435 Leu Glu Ser 450 <210> 13 <211> 1041 <212> DNA <213> Homo sapiens <400> 13 atggagagaa aatttatgtc cttgcaacca tccatctccg tatcagaaat ggaaccaaat 60 ggcaccttca gcaataacaa cagcaggaac tgcacaattg aaaacttcaa gagagaattt 120 ttcccaattg tatatctgat aatattttc tggggagtct tgggaaatgg gttgtccata 180 tatgttttcc tgcagcctta taagaagtcc acatctgtga acgttttcat gctaaatctg 240 gccatttcag atctcctgtt cataagcacg cttcccttca gggctgacta ttatcttaga 300 ggctccaatt ggatatttgg agacctggcc tgcaggatta tgtcttattc cttgtatgtc 360 aacatgtaca gcagtattta tttcctgacc gtgctgagtg ttgtgcgttt cctggcaatg 420 gttcacccct ttcggcttct gcatgtcacc agcatcagga gtgcctggat cctctgtggg 480 atcatatgga teettateat ggetteetea ataatgetee tggaeagtgg etetgageag 540 aacggcagtg tcacatcatg cttagagctg aatctctata aaattgctaa gctgcagacc 600 atgaactata ttgccttggt ggtgggctgc ctgctgccat ttttcacact cagcatctgt 660 tatctgctga tcattcgggt tctgttaaaa gtggaggtcc cagaatcggg gctgcgggtt 720 totcacagga aggcactgac caccatcatc atcaccttga tcatcttctt cttgtgtttc 780 ctgccctatc acacactgag gaccgtccac ttgacgacat ggaaagtggg tttatgcaaa 840 gacagactgc ataaagcttt ggttatcaca ctggccttgg cagcagccaa tgcctgcttc 900 aatcctctgc tctattactt tgctggggag aattttaagg acagactaaa gtctgcactc 960 agaaaaggcc atccacagaa ggcaaagaca aagtgtgttt tccctgttag tgtgtggttg 1020 agaaaggaaa caagagtata a 1041 <210> 14 346 PRT <213> Homo sapiens

<211>

<212>

<400> 14

Met Glu Arg Lys Phe Met Ser Leu Gln Pro Ser Ile Ser Val Ser Glu

Met Glu Pro Asn Gly Thr Phe Ser Asn Asn Asn Ser Arg Asn Cys Thr 25 30

| Ile | Glu | Asn 35 | Phe | Lys | Arg | Glu | Phe 40 | Phe | Pro | Ile | Val | Tyr 45 | Leu | Ile | Ile |
|------------|--------------------------|--------------------------|------------|------------|------------|------------|------------|------------|------------|--------------------|------------|------------|------------|------------|------------|
| Phe | Phe 50 | Trp | Gly | Val | Leu | Gly 55 | Asn | Gly | Leu | Ser | Ile 60 | Tyr | Val | Phe | Leu |
| Gln 65 | Pro | Tyr | Lys | Lys | Ser 70 | Thr | Ser | Val | Asn | Val 75 | Phe | Met | Leu | Asn | Leu 80 |
| Ala | Ile | Ser | Asp | Leu 85 | Leu | Phe | Ile | Ser | Thr 90 | Leu | Pro | Phe | Arg | Ala 95 | Asp |
| Tyr | Tyr | Leu | Arg 100 | Gly | Ser | Asn | Trp | Ile 105 | Phe | Gly | Asp | Leu | Ala 110 | Cys | Arg |
| Ile | Met | Ser 115 | Tyr | Ser | Leu | Tyr | Val 120 | Asn | Met | Tyr | Ser | Ser 125 | Ile | Tyr | Phe |
| Leu | Thr 130 | Val | Leu | Ser | Val | Val 135 | Arg | Phe | Leu | Ala | Met 140 | Val | His | Pro | Phe |
| Arg 145 | Leu | Leu | His | Val | Thr 150 | Ser | Ile | Arg | Ser | Al a 155 | Trp | Ile | Leu | Суз | Gly 160 |
| Ile | Ile | Trp | Ile | Leu 165 | Ile | Met | Ala | Ser | Ser 170 | Ile | Met | Leu | Leu | Asp 175 | Ser |
| Gly | Ser | Glu | Gln 180 | Asn | Gly | Ser | Val | Thr 185 | Ser | Cys | Leu | Glu | Leu 190 | Asn | Leu |
| Tyr | Lys | Ile 195 | | Lys | Leu | Gln | Thr 200 | Met | Asn | Tyr | Ile | Ala 205 | Leu | Val | Val |
| Gly | Cys 210 | | Leu | Pro | Phe | Phe 215 | Thr | Leu | Ser | Ile | Cys 220 | Tyr | Leu | Leu | Ile |
| 11e 225 | Arg | Val | Leu | Leu | Lys 230 | Val | Glu | Val | Pro | Glu 235 | Ser | Gly | Leu | Arg | Val 240 |
| Ser | His | Arg | Lys | Ala 245 | | Thr | Thr | Ile | 11e 250 | Ile | Thr | Leu | Ile | Ile 255 | Phe |
| Phe | Leu | Cys | Phe 260 | | Pro | Tyr | His | Thr 265 | Leu | Arg | Thr | Val | His 270 | Leu | Thr |
| Thr | Trp | Lys 275 | | Gly | Leu | Cys | Lys 280 | Asp | Arg | Leu | His | Lys 285 | Ala | Leu | Val |
| Ile | Thr 290 | | Ala | Lev | ı Ala | Ala 295 | Ala | Asn | Ala | Cys | Phe 300 | Asn | Pro | Leu | Leu |
| Туг 305 | | Phe | e Ala | Gly | Glu 310 | Asr | Phe | Lys | Asp | Arg 315 | Leu | Lys | Ser | Ala | Leu 320 |
| Arç | , Lys | 61) | / His | 325 | | Lys | s Ala | Lys | 330 | Lys | Cys | Val | . Phe | 9rc 335 | Val |
| Sei | val | Trp | Let 340 | | g Lys | s Glu | ı Thi | 345 | Val | L | | | | | |
| <2: <2: | 10> 11> 12> 13> | 15 152 DNA Home | 7 o saj | pien | 5 | | | | | | | | | | |

Page 15

```
<400> 15
 atgacgtcca cctgcaccaa cagcacgcgc gagagtaaca gcagccacac gtgcatgccc
                                                                        60
 ctctccaaaa tgcccatcag cctggcccac ggcatcatcc gctcaaccgt gctggttatc
                                                                       120
 ttcctcgccg cctctttcgt cggcaacata gtgctggcgc tagtgttgca gcgcaagccg
                                                                       180
 cagetgetge aggtgaceaa cegttttate tttaacetee tegteacega cetgetgeag
                                                                       240
 atttcgctcg tggccccctg ggtggtggcc acctctgtgc ctctcttctg gcccctcaac
                                                                       300
 agccacttct gcacggccct ggttagcctc acccacctgt tcgccttcgc cagcgtcaac
                                                                       360
 accattgtcg tggtgtcagt ggatcgctac ttgtccatca tccaccctct ctcctacccg
                                                                       420
 tocaagatga cocagogoog oggttacotg otcototatg goacotggat tgtggccato
                                                                       480
 ctgcagagca ctcctccact ctacggctgg ggccaggctg cctttgatga gcgcaatgct
                                                                       540
 ctctgctcca tgatctgggg ggccagcccc agctacacta ttctcagcgt ggtgtccttc
                                                                       600
atcgtcattc cactgattgt catgattgcc tgctactccg tggtgttctg tgcagcccgg
                                                                       660
aggcagcatg ctctgctgta caatgtcaag agacacagct tggaagtgcg agtcaaggac
                                                                      720
tgtgtggaga atgaggatga agagggagca gagaagaagg aggagttcca ggatgagagt
                                                                      780
gagtttcgcc gccagcatga aggtgaggtc aaggccaagg agggcagaat ggaagccaag
                                                                      840
gacggcagcc tgaaggccaa ggaaggaagc acggggacca gtgagagtag tgtagaggcc
                                                                      900
aggggcagcg aggaggtcag agagagcagc acggtggcca gcgacggcag catggagggt
                                                                      960
aaggaaggca gcaccaaagt tgaggagaac agcatgaagg cagacaaggg tcgcacagag
                                                                     1020
gtcaaccagt gcagcattga cttgggtgaa gatgacatgg agtttggtga agacgacatc
                                                                     1080
aatttcagtg aggatgacgt cgaggcagtg aacatcccgg agagcctccc acccagtcgt
                                                                     1140
cgtaacagca acagcaaccc tcctctgccc aggtgctacc agtgcaaagc tgctaaagtg
                                                                     1200
atcttcatca tcattttctc ctatgtgcta tccctggggc cctactgctt tttagcagtc
                                                                     1260
ctggccgtgt gggtggatgt cgaaacccag gtaccccagt gggtgatcac cataatcatc
                                                                     1320
tggcttttct tcctgcagtg ctgcatccac ccctatgtct atggctacat gcacaagacc
                                                                     1380
attaagaagg aaatccagga catgctgaag aagttcttct gcaaggaaaa gcccccgaaa
                                                                    1440
gaagatagcc acccagacct gcccggaaca gagggtggga ctgaaggcaa gattgtccct
                                                                    1500
tcctacgatt ctgctacttt tccttga
                                                                    1527
```

<210> 16 <211> 508 <212> PRT <213> Homo sapiens

<400> 16

Met Thr Ser Thr Cys Thr Asn Ser Thr Arg Glu Ser Asn Ser Ser His $1 \ 5 \ 10 \ 15$

Page 16

| Thr | Cys | Met | Pro 20 | Leu | Ser | Lys | Met | Pro 25 | Ile | Ser | Leu | Ala | 30 | GIĀ | 11e |
|------------|------------|------------|-------------|--------------|------------|------------|------------|------------|------------|------------|------------|------------------|------------|------------|------------|
| Ile | Arg | Ser 35 | Thr | Val | Leu | Val | Ile 40 | Phe | Leu | Ala | Ala | Ser 45 | Phe | Val | Gly |
| Asn | Ile 50 | Val | Leu | Ala | Leu | Val 55 | Leu | Gln | Arg | Lys | Pro 60 | Gln [.] | Leu | Leu | Gln |
| Val 65 | Thr | Asn | Arg | Phe | Ile 70 | Phe | Asn | Leu | Leu | Val 75 | Thr | Asp | Leu | Leu | Gln 80 |
| Ile | Ser | Leu | Val | Ala 85 | Pro | Trp | Val | Val | Ala 90 | Thr | Ser | Val | Pro | Leu 95 | Phe |
| Trp | Pro | Leu | Asn 100 | Ser | His | Phe | Cys | Thr 105 | Ala | Leu | Val | Ser | Leu 110 | Thr | His |
| Leu | Phe | Ala 115 | | Ala | Ser | Val | Asn 120 | Thr | Ile | Val | Val | Val 125 | Ser | Val | Asp |
| Arg | Tyr 130 | | Ser | Ile | Ile | His 135 | Pro | Leu | Ser | Tyr | Pro 140 | Ser | Lys | Met | Thr |
| Gln 145 | | Arg | Gly | Tyr | Leu 150 | Leu | Leu | туr | Gly | Thr 155 | Trp | Ile | Val | Ala | Ile 160 |
| Leu | Gln | Ser | Thr | Pro 165 | Pro | Leu | Tyr | Gly | Trp 170 | Gly | Gln | Ala | Ala | Phe 175 | Asp |
| Glu | Arg | Asr | Ala 180 | Leu) | Cys | Ser | Met | Ile 185 | Trp | Gly | Ala | Ser | Pro 190 | Ser | Tyr |
| Thr | Ile | Leu 195 | | · Val | Val | Ser | Phe 200 | Ile | Val | Ile | Pro | Leu 205 | Ile | Val | Met |
| Ile | Ala 210 | | 5 Туг | Ser | Val | Val 215 | Phe | Cys | Ala | Ala | Arg 220 | Arg | Gln | His | Ala |
| Leu 225 | | ту: | c Ası | n Val | Lys 230 | Arg | His | Ser | Leu | Glu 235 | Val | Arg | Val | Lys | Asp 240 |
| Cys | va: | l Gl | u Ası | n Glu 245 | Asp | Glu | ı Glu | Gly | Ala 250 | Glu | Lys | Lys | Glu , | Glu 255 | Phe |
| | | | 26 | 0 | | | | 265 | • | | | | 270 | , | Ala |
| | | 27 | 5 | | | | 280 | , | | | | 20. | , | | Glu |
| Gl | y Se 29 | | r Gl | y Th | r Ser | Gl: 29 | ı Sen 5 | Sei | · Val | l Gl | Ala 300 | a Arg | g Gl) | / Ser | Glu |
| 30 | 5 | | | | 310 |) | | | | 31 | 5 | | | | 320 |
| Ly | s Gl | u Gl | y Se | r Th 32 | r Lys 5 | s Va | 1 G1: | ı Glı | 33 | n Se O | r Me | t Ly | s Ala | 33! | b Lys |
| Gl | y Ar | g Th | r Gl 34 | | l Ası | n Gl | n Cy | s Se 34 | r Il | e As | p Le | u Gl | y Gl | u Asj O | p Asp |
| Me | t Gl | | ne G1 55 | .y G1 | u Ası | p As | p Il | e As O | n Ph | e Se | | 30 | 3 | p Va | l Glu |
| | | | | | | | | | | | Pag | re 17 | • | | |

| Ala Val 1 370 | Asn Ile | Pro Glu | Ser L 375 | eu Pro | Pro | Ser | Arg 380 | Arg | Asn | Ser | Asn | |
|----------------------|---------------|----------------|-----------------|---------------|------------|------------|------------|------------|------------|------------|------------|-----|
| Ser Asn B 385 | Pro Pro | Leu Pro | Arg C | ys Tyr | Gln | Cys 395 | Lys | Ala | Ala | Lys | Val 400 | |
| Ile Phe 1 | le Ile | Ile Phe 405 | Ser T | yr Val | Leu 410 | Ser | Leu | Gly | Pro | Tyr 415 | Cys | |
| Phe Leu A | la Val 420 | Leu Ala | Val T | rp Val 425 | Asp | Val | Glu | Thr | Gln 430 | Val | Pro | |
| Gln Trp V | al Ile 35 | Thr Ile | Ile II | le Trp 10 | Leu | Phe | Phe | Leu 445 | Gln | Cys | Cys | |
| Ile His P 450 | ro Tyr | Val Tyr | Gly Ty 455 | /r Met | His | Lys | Thr 460 | Ile | Lys | Lys | Glu | |
| Ile Gln A 465 | sp Met | Leu Lys 470 | Lys P | e Phe | Cys | Lys 475 | Glu | Lys | Pro | Pro | Lys 480 | |
| Glu Asp S | er His | Pro Asp 485 | Leu Pr | o Gly | Thr 490 | Glu | Gly | Gly | Thr | Glu 495 | Gly | |
| Lys Ile V | al Pro 500 | Ser Tyr | Asp Se | r Ala 505 | Thr | Phe | Pro | | | | | |
| <210> 17 <211> 10 | 60 | | | | | | | | | | | |
| <212> DNA | P | | | | | | | | | | | |
| <400> 17 | no sapie | ens | | | | | | | | | | |
| atgcccttga | a cggac | ggcat tt | cttcat | tt gag | gacc | tct | tggci | taaca | aa t | atcc | tcaga | 60 |
| atatttgtct | gggtta | atagc tt | tcatta | cc tgc | tttg | gaa | atcti | tttt | gt c | attg | gcatg | 120 |
| agatctttca | ttaaaq | gctga aa | atacaa | ct cac | gctai | tgt (| ccato | caaaa | at c | cttt | gttgc | 180 |
| gctgattgcc | tgatgo | ggtgt tt | acttgt | tc ttt | gttg | gca 1 | ttttc | gata | at aa | aaata | accga | 240 |
| gggcagtato | | | | | | | | | | | _ | 300 |
| ttcctggcca | | | | | | | | | | | _ | 360 |
| aagttcctgg | | | | | | | | | | | | 420 |
| gtcatcctca | tttgca | itctg ga | tggcggg | ga ttt | ttaat | ag c | ctgta | atto | c at | tttc | ıgaat | 480 |
| aaggattatt | ttggaa | actt tt | atgggaa | a aat | ggagt | at ç | gtttc | ccac | t tt | atta | tgac | 540 |
| caaacagaag | atattg | gaag caa | aagggt <i>a</i> | t tct | cttgg | jaa t | tttc | ctag | g tg | ıtgaa | cttg | 600 |
| ctggcttttc | tcatca | ttgt gt | ttccta | t atta | actat | gt t | ctgt | tcca | t to | aaaa | aacc | 660 |
| gccttgcaga | | | | | | | | | | | | 720 |
| ttctttttta | | | | | | | | | | | | 780 |
| ctttccctct | | | | | | | | | | | | 840 |
| cttccagtta | | | | | | | | | | | | 900 |
| gacaagttga | | | | | | | | | | | | 960 |
| | | | ٠ | | | | ge 1 | | | | | - |

1020 1068

ttgaacaaaa taacacttgg agacagtata atgaaaccag tttcctag <210> 18 <211> 355 <212> PRT <213> Homo sapiens <400> 18 Met Pro Leu Thr Asp Gly Ile Ser Ser Phe Glu Asp Leu Leu Ala Asn Asn Ile Leu Arg Ile Phe Val Trp Val Ile Ala Phe Ile Thr Cys Phe Gly Asn Leu Phe Val Ile Gly Met Arg Ser Phe Ile Lys Ala Glu Asn Thr Thr His Ala Met Ser Ile Lys Ile Leu Cys Cys Ala Asp Cys Leu Met Gly Val Tyr Leu Phe Phe Val Gly Ile Phe Asp Ile Lys Tyr Arg Gly Gln Tyr Gln Lys Tyr Ala Leu Leu Trp Met Glu Ser Val Gln Cys 85 90 95 Arg Leu Met Gly Phe Leu Ala Met Leu Ser Thr Glu Val Ser Val Leu Leu Leu Thr Tyr Leu Thr Leu Glu Lys Phe Leu Val Ile Val Phe Pro Phe Ser Asn Ile Arg Pro Gly Lys Arg Gln Thr Ser Val Ile Leu Ile Cys Ile Trp Met Ala Gly Phe Leu Ile Ala Val Ile Pro Phe Trp Asn Lys Asp Tyr Phe Gly Asn Phe Tyr Gly Lys Asn Gly Val Cys Phe Pro 165 170 175 Leu Tyr Tyr Asp Gln Thr Glu Asp Ile Gly Ser Lys Gly Tyr Ser Leu 180 185 190 Gly Ile Phe Leu Gly Val Asn Leu Leu Ala Phe Leu Ile Ile Val Phe Ser Tyr Ile Thr Met Phe Cys Ser Ile Gln Lys Thr Ala Leu Gln Thr 210 215 220 Thr Glu Val Arg Asn Cys Phe Gly Arg Glu Val Ala Val Ala Asn Arg Phe Phe Phe Ile Val Phe Ser Asp Ala Ile Cys Trp Ile Pro Val Phe Val Val Lys Ile Leu Ser Leu Phe Arg Val Glu Ile Pro Asp Thr Met Thr Ser Trp Ile Val Ile Phe Phe Leu Pro Val Asn Ser Ala Leu Asn Page 19

aaaagtttat ctacatccat tgtgtggata gaggactcct cttccctgaa acttggggtt

| 275 | 280 | 285 |
|--|----------------------------|------------------------|
| Pro Ile Leu Tyr Thr Leu Thr 290 295 | Thr Asn Phe Phe Lys 300 | Asp Lys Leu Lys |
| Gln Leu Leu His Lys His Gln 305 310 | Arg Lys Ser Ile Phe 315 | Lys Ile Lys Lys 320 |
| Lys Ser Leu Ser Thr Ser Ile 325 | Val Trp Ile Glu Asp 330 | Ser Ser Ser Leu 335 |
| Lys Leu Gly Val Leu Asn Lys 340 | Ile Thr Leu Gly Asp 345 | Ser Ile Met Lys 350 |
| Pro Val Ser 355 | | |
| <210> 19 <211> 969 <212> DNA <213> Homo sapiens | | |
| <400> 19 atggatecaa ecateteaae ettggad | caca daactdacac caat | |
| actetttget acaageagae ettgage | | |
| gggctgacag gaaacgcagt tgtgctc | | |
| ttctccatct acatcctcaa cttggcc | | |
| atatattccc tgttaagctt catcagt | | |
| gtgatgatgt tttcctactt tgcaggc | | |
| tgcctgtccg tcctgtggcc catctgg | | |
| | | |
| gtggtgtgt teetgetetg ggeeetg | | |
| tgtggcttcc tgttcagtgg tgctgat | | |
| gtcgcgtggc tgatttttt atgtgtg | | |
| aggattetet gtggateeeg gaagata | | |
| acagtactgg tetteeteet etgtgge | | |
| tggatccacg tggacaggga agtctta | | |
| teegetetta acageagtge caaceee. | | |
| cgtcaaaata ggcagaacct gaagctg | gtt ctccagaggg ctctgd | cagga cgcgtctgag 900 |
| gtggatgaag gtggagggca gcttcctg | gag gaaatcctgg agctgt | cggg aagcagattg 960 |
| gagcagtga | | 969 |
| <210> 20 <211> 322 <212> PRT <213> Homo sapiens | | |

Page 20

<400> 20

Met Asp Pro Thr Ile Ser Thr Leu Asp Thr Glu Leu Thr Pro Ile Asn 1 5 10 15 Gly Thr Glu Glu Thr Leu Cys Tyr Lys Gln Thr Leu Ser Leu Thr Val 20 25 30 Leu Thr Cys Ile Val Ser Leu Val Gly Leu Thr Gly Asn Ala Val Val 35 40 45Leu Trp Leu Leu Gly Cys Arg Met Arg Arg Asn Ala Phe Ser Ile Tyr 50 55 60 Ile Leu Asn Leu Ala Ala Asp Phe Leu Phe Leu Ser Gly Arg Leu 65 70 75 80 Ile Tyr Ser Leu Leu Ser Phe Ile Ser Ile Pro His Thr Ile Ser Lys Ile Leu Tyr Pro Val Met Met Phe Ser Tyr Phe Ala Gly Leu Ser Phe Leu Ser Ala Val Ser Thr Glu Arg Cys Leu Ser Val Leu Trp Pro Ile Trp Tyr Arg Cys His Arg Pro Thr His Leu Ser Ala Val Val Cys Val Leu Leu Trp Ala Leu Ser Leu Leu Arg Ser Ile Leu Glu Trp Met Leu Cys Gly Phe Leu Phe Ser Gly Ala Asp Ser Ala Trp Cys Gln Thr Ser Asp Phe Ile Thr Val Ala Trp Leu Ile Phe Leu Cys Val Val Leu Cys Gly Ser Ser Leu Val Leu Leu Ile Arg Ile Leu Cys Gly Ser Arg Lys 195 200 205 Ile Pro Leu Thr Arg Leu Tyr Val Thr Ile Leu Leu Thr Val Leu Val Phe Leu Cys Gly Leu Pro Phe Gly Ile Gln Phe Phe Leu Phe Leu Trp Ile His Val Asp Arg Glu Val Leu Phe Cys His Val His Leu Val Ser Ile Phe Leu Ser Ala Leu Asn Ser Ser Ala Asn Pro Ile Ile Tyr Phe Phe Val Gly Ser Phe Arg Gln Arg Gln Asn Arg Gln Asn Leu Lys Leu Val Leu Gln Arg Ala Leu Gln Asp Ala Ser Glu Val Asp Glu Gly Gly Gly Gln Leu Pro Glu Glu Ile Leu Glu Leu Ser Gly Ser Arg Leu Glu Gln

Page 21

<210> 21 <211> 1305

<212> DNA <213> Homo sapiens atggaggate tetttagece etcaattetg eegeeggege ecaacattte egtgeeeate 60 ttgctgggct ggggtctcaa cctgaccttg gggcaaggag cccctgcctc tgggccgccc 120 ageegeegeg teegeetggt gtteetgggg gteateetgg tggtggeggt ggeaggeaae 180 240 aagatggact tcctgctggt gcagctggcc ctggcggacc tgtacgcgtg cgggggcacg 300 gegetgteae agetggeetg ggaactgetg ggegageeec gegeggeeae gggggaeetg 360 gcgtgccgct tcctgcagct gctgcaggca tccgggcggg gcgcctcggc ccacctcgtg 420 gtgctcatcg ccctcgagcg ccggcgcgcg gtgcgtcttc cgcacggccg gccgctgccc 480 gegegtgeec tegeegeect gggetggetg etggeactge tgetggeget geeceeggee 540 ttegtggtge geggggaete eccetegeeg etgeegeege egeegeegee aacgteeetg 600 cagecaggeg egeceegge egecegegee tggeeggggg agegtegetg ecaegggate 660 ttcgcgcccc tgccgcgctg gcacctgcag gtctacgcgt tctacgaggc cgtcgcgggc 720 ttcgtcgcgc ctgttacggt cctgggcgtc gcttgcggcc acctactctc cgtctggtgg 780 eggeacegge egeaggeece egeggetgea gegeectggt eggegageec aggtegagee 840 cctgcgccca gcgcgctgcc ccgcgccaag gtgcagagcc tgaagatgag cctgctgctg 900 gegetgetgt tegtgggetg egagetgeee tactttgeeg eeeggetgge ggeegegtgg 960 tegteeggge cegegggaga etgggaggga gagggeetgt eggeggeget gegegtggtg 1020 gcgatggcca acagcgctct caatcccttc gtctacc+ct tcttccaggc gggcgactgc 1080 cggctccggc gacagctgcg gaagcggctg ggctctctgt gctgcgcgcc gcagggaggc 1140 gcggaggacg aggagggcc ccggggccac caggcgctct accgccaacg ctggccccac 1200 cctcattatc accatgctcg gcgggaaccg ctggacgagg gcggcttgcg cccacccct 1260 ccgcgcccca gacccctgcc ttgctcctgc gaaagtgcct tctag 1305

<210> 22 <211> 434

<212> PRT

<213> Homo sapiens

<400> 22

Met Glu Asp Leu Phe Ser Pro Ser Ile Leu Pro Pro Ala Pro Asn Ile 1 5 10 15

Ser Val Pro Ile Leu Leu Gly Trp Gly Leu Asn Leu Thr Leu Gly Gln 20 25 30

Gly Ala Pro Ala Ser Gly Pro Pro Ser Arg Arg Val Arg Leu Val Phe 35 40 . 45

| Leu | Gly 50 | Val | Ile | Leu | Val | Val 55 | Ala | Val | Ala | Gly | Asn 60 | Thr | Thr | Val | Leu |
|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| Cys 65 | Arg | Leu | Cys | Gly | Gly 70 | Gly | Gly | Pro | Trp | Ala 75 | Gly | Pro | Lys | Arg | Arg 80 |
| Lys | Met | Asp | Phe | Leu 85 | Leu | Val | Gln | Leu | Ala 90 | Leu | Ala | Asp | Leu | Tyr 95 | Ala |
| Cys | Gly | Gly | Thr 100 | Ala | Leu | Ser | Gln | Leu 105 | Ala | Trp | Glu | Leu | Leu 110 | Gly | Glu |
| Pro | Arg | Ala 115 | Ala | Thr | Gly | Asp | Leu 120 | Ala | Cys | Arg | Phe | Leu 125 | Gln | Leu | Leu |
| Gln | Ala 130 | Ser | Gly | Arg | Gly | Ala 135 | Ser | Ala | His | Leu | Val 140 | Val | Leu | Ile | Ala |
| Leu 145 | Glu | Arg | Arg | Arg | Ala 150 | Val | Arg | Leu | Pro | His 155 | Gly | Arg | Pro | Leu | Pro 160 |
| Ala | Arg | Ala | Leu | Ala 165 | Ala | Leu | Gly | Trp | Leu 170 | Leu | Ala | Leu | Leu | Leu 175 | Ala |
| | | | 180 | | | | Arg | 185 | | | | | 190 | - | |
| | | 195 | | | | | Leu 200 | | | | | 205 | | | |
| | 210 | | | | | 215 | Arg | | | | 220 | | | | |
| 225 | | | | | 230 | | Tyr | | | 235 | | | | | 240 |
| | | | | 245 | | | Leu | | 250 | | | | | 255 | |
| | | • | 260 | | | | Pro | 265 | | | | | 270 | | |
| | | 275 | | | | | Ala 280 | | | | | 285 | · | | |
| | 290 | | | | | 295 | Met | | | | 300 | | | | |
| 305 | | | | | 310 | | Phe | | | 315 | | | | | 320 |
| | | | | 325 | | | | | 330 | | | | | 335 | Ala |
| | | | 340 | | | | Asn | 345 | | | | | 350 | | |
| | | 355 | | | | | 360 | | | | | 365 | | | Lys |
| | 370 | | | | | 375 | | | | | 380 | | | | Glu |
| Glu 385 | | Pro | Arg | Gly | His 390 | | Ala | Leu | Tyr | Arg 395 | | | Trp | Pro | His 400 |
| | | | | | | | | | | | Page | | | | |

540

600

Pro His Tyr His His Ala Arg Arg Glu Pro Leu Asp Glu Gly Gly Leu 405 410 415

Arg Pro Pro Pro Pro Arg Pro Arg Pro Leu Pro Cys Ser Cys Glu Ser 420 425 430

Ala Phe

| <210> <211> <212> <213> | 23 104 DNA Hom | _ | | | | | |
|----------------------------------|-------------------------|------------|------------|------------|------------|------------|--|
| <400> | 23 | | | | | | |
| atgtaca | acg | ggtcgtgctg | ccgcatcgag | ggggacacca | tctcccaggt | gatgccgccg | |
| ctgctca | ittg | tggcctttgt | gctgggcgca | ctaggcaatg | gggtcgccct | gtgtggtttc | |
| tgcttcc | aca | tgaagacctg | gaagcccagc | actgtttacc | ttttcaattt | ggccgtggct | |
| gatttcc | tcc | ttatgatctg | cctgcctttt | cggacagact | attacctcag | acgtagacac | |
| tgggctt | ttg | gggacattcc | ctgccgagtg | gggctcttca | cgttggccat | gaacagggcc | |
| gggagca | tcg | tgttccttac | ggtggtggct | gcggacaggt | atttcaaagt | ggtccacccc | |
| caccacg | cgg | tgaacactat | ctccacccgg | gtggcggctg | gcatcgtctg | caccctgtgg | |
| gccctgg | tca | tcctgggaac | agtgtatctt | ttgctggaga | accatctctg | cgtgcaagag | |

tggagcetga ggcggaggca gcagetggce agacaggete ggatgaagaa ggcgaceegg 660 tteateatgg tggtggcaat tgtgtteate acatgetace tgcccagegt gtetgetaga 720

acggccgtct cctgtgagag cttcatcatg gagtcggcca atggctggca tgacatcatg

ttccagctgg agttctttat gcccctcggc atcatcttat tttgctcctt caagattgtt

ctctatttcc tctggacggt gcctcgagt gcctgcgatc cctctgtcca tggggccctg 780

cacataacce tcagettcae ctacatgaac agcatgetgg atccetggt gtattatttt 840 tcaageeeet eetteesaa attetacaac aageteaaaa tetgeagtet gaaacceaag 900

cagccaggac actcaaaaac acaaaggccg gaagagatgc caatttcgaa cctcggtcgc 960

aggagttgca tcagtgtggc aaatagtttc caaagccagt ctgatgggca atgggatccc 1020

cacattgttg agtggcactg a 1041

<210> 24 <211> 346

<212> PRT

<213> Homo sapiens

<400> 24

Met Tyr Asn Gly Ser Cys Cys Arg Ile Glu Gly Asp Thr Ile Ser Gln $1 \hspace{1cm} 5 \hspace{1cm} 10 \hspace{1cm} 15$

Val Met Pro Pro Leu Leu Ile Val Ala Phe Val Leu Gly Ala Leu Gly 20 25 30

Asn Gly Val Ala Leu Cys Gly Phe Cys Phe His Met Lys Thr Trp Lys Pro Ser Thr Val Tyr Leu Phe Asn Leu Ala Val Ala Asp Phe Leu Leu Met Ile Cys Leu Pro Phe Arg Thr Asp Tyr Tyr Leu Arg Arg Arg His 65 70 75 80 Trp Ala Phe Gly Asp Ile Pro Cys Arg Val Gly Leu Phe Thr Leu Ala Met Asn Arg Ala Gly Ser Ile Val Phe Leu Thr Val Val Ala Ala Asp Arg Tyr Phe Lys Val Val His Pro His His Ala Val Asn Thr Ile Ser Thr Arg Val Ala Ala Gly Ile Val Cys Thr Leu Trp Ala Leu Val Ile Leu Gly Thr Val Tyr Leu Leu Leu Glu Asn His Leu Cys Val Gln Glu Thr Ala Val Ser Cys Glu Ser Phe Ile Met Glu Ser Ala Asn Gly Trp 170 His Asp Ile Met Phe Gln Leu Glu Phe Phe Met Pro Leu Gly Ile Ile Leu Phe Cys Ser Phe Lys Ile Val Trp Ser Leu Arg Arg Gln Gln Leu Ala Arg Gln Ala Arg Met Lys Lys Ala Thr Arg Phe Ile Met Val Val Ala Ile Val Phe Ile Thr Cys Tyr Leu Pro Ser Val Ser Ala Arg Leu Tyr Phe Leu Trp Thr Val Pro Ser Ser Ala Cys Asp Pro Ser Val His Gly Ala Leu His Ile Thr Leu Ser Phe Thr Tyr Met Asn Ser Met Leu Asp Pro Leu Val Tyr Tyr Phe Ser Ser Pro Ser Phe Pro Lys Phe Tyr Asn Lys Leu Lys Ile Cys Ser Leu Lys Pro Lys Gln Pro Gly His 290 295 300 Ser Lys Thr Gln Arg Pro Glu Glu Met Pro Ile Ser Asn Leu Gly Arg Arg Ser Cys Ile Ser Val Ala Asn Ser Phe Gln Ser Gln Ser Asp Gly Gln Trp Asp Pro His Ile Val Glu Trp His <210> 25 1011 DNA <212>

Page 25

<213> Homo sapiens

| <400> 25 | | | | | | |
|------------|--------------|--------------|--------------|------------|--------------|------|
| | | | | | c tttaccaatc | 60 |
| atttacatc | c tcctttgtat | tgttggtgtt | : tttggaaaca | ctctctctc | a atggatattt | 120 |
| ttaacaaaa | a taggtaaaaa | aacatcaacg | cacatctacc | tgtcacacct | tgtgactgca | 180 |
| aacttactt | g tgtgcagtgc | catgcctttc | atgagtatct | atttcctgaa | aggtttccaa | 240 |
| tgggaatato | aatctgctca | atgcagagtg | gtcaattttc | tgggaactct | atccatgcat | 300 |
| gcaagtatgt | ttgtcagtct | cttaatttta | agttggattg | ccataagccg | ctatgctacc | 360 |
| ttaatgcaaa | aggattcctc | gcaagagact | acttcatgct | atgagaaaat | attttatggc | 420 |
| catttactga | aaaaatttcg | ccagcccaac | tttgctagaa | aactatgcat | ttacatatgg | 480 |
| ggagttgtac | tgggcataat | cattccagtt | accgtatact | actcagtcat | agaggctaca | 540 |
| | | | | | gatctctcag | 600 |
| attgcaggtc | tcattggaac | cacatttatt | ggattttcct | ttttagtagt | actaacatca | 660 |
| | ttgtaagcca | | | | | 720 |
| | acagttctgt | | | | | 780 |
| tgcttccttc | cttatagtat | ttttaaaccc | attttttatg | ttctacacca | aagagataac | 840 |
| tgtcagcaat | tgaattattt | aatagaaaca | aaaaacattc | tcacctgtct | tgcttcggcc | 900 |
| | cagaccccat | | | | | 960 |
| tataatctct | ttacaaagtc | taattcagca ' | catatgcaat . | catatggttg | a | 1011 |

<210> 26 <211> 336

<211> 336 <212> PRT

<213> Homo sapiens

<400> 26

Met Asn Asn Asn Thr Thr Cys Ile Gln Pro Ser Met Ile Ser Ser Met 1 5 10 15

Ala Leu Pro Ile Ile Tyr Ile Leu Leu Cys Ile Val Gly Val Phe Gly 20 25 30

Asn Thr Leu Ser Gln Trp Ile Phe Leu Thr Lys Ile Gly Lys Lys Thr 35 40 45

Ser Thr His Ile Tyr Leu Ser His Leu Val Thr Ala Asn Leu Leu Val 50 60

Cys Ser Ala Met Pro Phe Met Ser Ile Tyr Phe Leu Lys Gly Phe Gln 70 75 80

Trp Glu Tyr Gln Ser Ala Gln Cys Arg Val Val Asn Phe Leu Gly Thr 85 90 95

Leu Ser Met His Ala Ser Met Phe Val Ser Leu Leu Ile Leu Ser Trp 100 105 110

| Ile | Ala | 11e 115 | Ser | Arg | Туr | Ala | Thr 120 | Leu | Met | Gln | Lys | Asp 125 | Ser | Ser | Gln | |
|--------------------------|-------------|---------------------------|------------|------------|-------------------|------------|----------------|------------|------------|------------|------------|------------|------------|------------|------------|-----|
| Glu | Thr 130 | Thr | Ser | Cys | Tyr | Glu 135 | Lys | Ile | Phe | Tyr | Gly 140 | His | Leu | Leu | Lys | |
| Lys 145 | Phe | Arg | Gln | Pro | Asn 150 | Phe | Ala | Arg | Lys | Leu 155 | Cys | Ile | Tyr | Ile | Trp 160 | |
| Gly | Val | Val | Leu | Gly 165 | Ile | Ile | ·Ile | Pro | Val 170 | Thr | Val | Tyr | Tyr | Ser 175 | Val | |
| Ile | Glu | Ala | Thr 180 | Glu | Gly | Glu | Glu | Ser 185 | Leu | Суз | Tyr | Asn | Arg 190 | Gln | Met | |
| Glu | Leu | Gly 195 | Ala | Met | Ile | Ser | Gln 200 | Ile | Ala | Gly | Leu | 11e 205 | Gly | Thr | Thr | |
| Phe | Ile 210 | Gly | Phe | Ser | Phe | Leu 215 | Val | Val | Leu | Thr | Ser 220 | Tyr | Tyr | Ser | Phe | |
| Val 225 | Ser | His | Leu | Arg | Lys 230 | Ile | Arg | Thr | Cys | Thr 235 | Ser | Ile | Met | Glu | Lys 240 | |
| Asp | Leu | Thr | Tyr | Ser 245 | Ser | Val | Lys | Arg | His 250 | Leu | Leu | Val | Ile | Gln 255 | Ile | |
| Leu | Leu | Ile | Val 260 | Cys | Phe | Leu | Pro | Tyr 265 | Ser | Ile | Phe | Lys | Pro 270 | Ile | Phe | |
| Tyr | Val | Leu 275 | His | Gln | Arg | Asp | Asn 280 | Cys | Gln | Gln | Leu | Asn 285 | Tyr | Leu | Ile | |
| Glu | Thr 290 | | Asn | Ile | Leu | Thr 295 | Cys | Leu | Ala | Ser | .Ala | Arg | Ser | Ser | Thr | |
| 305 | | | | | 310 | | | | | 315 | • | | | | Leu 320 | |
| Tyr | Asn | Leu | ı Ph∈ | 325 | Lys | Ser | Asn | ser | 330 | His | Met | Gln | Ser | 335 | Gly | |
| <21 <21 <21 <21 | 1> 2> | 27 1014 DNA Homo | i sap | oiens | 5 | | | | | | | | • | | | |
| <40 ato | 0> Jaato | 27 gagc | cact | aga | cta 1 | ttag | jca a a | at go | cttct | gati | t tc | ccga | atta | tgca | igctgct | 60 |
| ttt | ggaa | att | gcad | ctga | tga a | aaaca | atcc | ca c | tcaaq | gatgo | act | tacci | ccc | tgti | atttat | 120 |
| | | | | | | | | | | | | | | | cattttc | 180 |
| | | | | | | | | | | | | | | | agatctg | 240 |
| | | | | | | | | | | | | | | | ctggatc | 300 |
| | | | | | | | | | | | | | | | tagcagc | 360 |
| | | | | | | | | | | | | | | | aatgagc | 420 |
| | | | | | | | | | | | | | | | gatcatt | 480 |
| | | | | | | | | | | | | | | | caacaga | 540 |
| | | | | | | | | | | | | je 27 | | | | |

| tcagcctgtc | tcgacctcac | cagttcggat | gaactcaata | ctattaagtg | gtacaacctg | 600 |
|------------|------------|------------|------------|------------|------------|-------|
| attttgactg | caactacttt | ctgcctcccc | ttggtgatag | tgacactttg | ctataccacg | 660 |
| attatccaca | ctctgaccca | tggactgcaa | actgacagct | gccttaagca | gaaagcacga | 720 |
| aggctaacca | ttctgctact | ccttgcattt | tacgtatgtt | ttttaccctt | ccatatcttg | 780 |
| agggtcattc | ggatcgaatc | tcgcctgctt | tcaatcagtt | gttccattga | gaatcagatc | 840 |
| catgaagctt | acatcgtttc | tagaccatta | gctgctctga | acacctttgg | taacctgtta | 900 |
| ctatatgtgg | tggtcagcga | caactttcag | caggctgtct | gctcaacagt | gagatgcaaa | . 960 |
| gtaagcggga | accttgagca | agcaaagaaa | attagttact | caaacaaccc | ttga | 1014 |
| | | | | | | |

<210> 28 <211> 337

<212> PRT

<213> Homo sapiens

<400> 28

Met Asn Glu Pro Leu Asp Tyr Leu Ala Asn Ala Ser Asp Phe Pro Asp 1 5 10 15

Tyr Ala Ala Ala Phe Gly Asn Cys Thr Asp Glu Asn Ile Pro Leu Lys 20 25 30

Met His Tyr Leu Pro Val Ile Tyr Gly Ile Ile Phe Leu Val Gly Phe 35 40 45

Pro Gly Asn Ala Val Val Ile Ser Thr Tyr Ile Phe Lys Met Arg Pro 50 60

Trp Lys Ser Ser Thr Ile Ile Met Leu Asn Leu Ala Cys Thr Asp Leu 65 70 75 80

Leu Tyr Leu Thr Ser Leu Pro Phe Leu Ile His Tyr Tyr Ala Ser Gly 85 90 95

Glu Asn Trp Ile Phe Gly Asp Phe Met Cys Lys Phe Ile Arg Phe Ser 100 105 140

Phe His Phe Asn Leu Tyr Ser Ser Ile Leu Phe Leu Thr Cys Phe Ser 115 120 125

Ile Phe Arg Tyr Cys Val Ile Ile His Pro Met Ser Cys Phe Ser Ile 130 135 140

His Lys Thr Arg Cys Ala Val Val Ala Cys Ala Val Val Trp Ile Ile
150 155 160

Ser Leu Val Ala Val Ile Pro Met Thr Phe Leu Ile Thr Ser Thr Asn 165 170 175

Arg Thr Asn Arg Ser Ala Cys Leu Asp Leu Thr Ser Ser Asp Glu Leu 180 185 190

Asn Thr Ile Lys Trp Tyr Asn Leu Ile Leu Thr Ala Thr Thr Phe Cys 195 200 205

Leu Pro Leu Val Ile Val Thr Leu Cys Tyr Thr Thr Ile Ile His Thr Page $28\,$

| | 210 | | | | | 215 | | | | | 220 | | | | | |
|--------------------------|------------|--------------------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|---------------------|-----|
| Leu 225 | Thr | His | Gly | Leu | Gln 230 | Thr | Asp | Ser | Суѕ | Leu 235 | Lys | Gln | Lys | Ala | Arg 2 4 0 | |
| Arg | Leu | Thr | Ile | Leu 245 | Leu | Leu | Leu | Ala | Phe 250 | Tyr | Val | Cys | Phe | Leu 255 | Pro | |
| Phe | His | Ile | Leu 260 | Arg | Val | Ile | Arg | Ile 265 | Glu | Ser | Arg | Leu | Leu 270 | Ser | Ile | |
| Ser | Cys | Ser 275 | Ile | Glu | Asn | Gln | 11e 280 | His | Glu | Ala | Tyr | Ile 285 | Val | Ser | Arg | |
| Pro | Leu 290 | Ala | Ala | Leu | Asn | Thr 295 | Phe | Gly | Asn | Leu | Leu 300 | Leu | Tyr | Val | Val | |
| Val 305 | Ser | Asp | Asn | Phe | Gln 310 | Gln | Ala | Val | Cys | Ser 315 | Thr | Val | Arg | Cys | Lys 320 | |
| Val | Ser | Gly | Asn | Leu 325 | Glu | Gln | Ala | Lys | Lys 330 | Ile | Ser | Tyr | Ser | Asn 335 | Asn | |
| Pro | | | | | | | | | | | | | | | | |
| <21 <21 <21 <21 | 1> 2> | 29 993 DNA Homo | sap | iens | | | | | | | | | | | | |
| <40 atg | 0> gatc | 29 caa | ccac | :cccg | gc c | tggg | gaac | a ga | aagt | acaa | cag | tgaa | tgg | aaat | gaccaa | 60 |
| acc | cttc | ttc | tgct | ttgt | .gg c | aagg | agac | c ct | gato | ccgg | tct | tcct | gat | cctt | ttcatt | 120 |
| | | | | | | | | | | | | | | | atgcgc | 180 |
| | | | | | | | | | | | | | | | ctctgc | |
| | | | | | | | | | | | | | | | atcaat | |
| | | | | | | | | | | | | | | | atgctg | |
| | | | | | | | | | | | | | | | tgccgc | |
| | | | | | | | | | | | | | | | actgctg | |
| | | | | | | | | | | | | | | | tggtgt | |
| | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | etgtggg daccag | |
| | | | | | | | | | | | | | | | gaccago | |
| | | | | | | | | | | | | | | | ctttggo | |
| | | | | | | | | | | | | | | | tcatatt | |
| | | | | | | | | | | | | | | | ttactto | |
| | | | | | | | | | | | | | | | ggctct | |
| | | | | | | | | | | cagt | g aa | .ggat | gctt | . ccg | tcaggg | |
| ac | cccg | gaga | tgt | cgag | aag | cagt | ctgg | jtg t | ag | | Pag | je 29 | • | | | 99: |

<210> 30 <211> 330 <212> PRT <213> Homo sapiens <400> 30 Met Asp Pro Thr Thr

Met Asp Pro Thr Thr Pro Ala Trp Gly Thr Glu Ser Thr Thr Val Asn 1 5 10 15

Gly Asn Asp Gln Ala Leu Leu Leu Cys Gly Lys Glu Thr Leu Ile 20 25 30

Pro Val Phe Leu Ile Leu Phe Ile Ala Leu Val Gly Leu Val Gly Asn 35 40 45

Gly Phe Val Leu Trp Leu Leu Gly Phe Arg Met Arg Arg Asn Ala Phe 50 60

Ser Val Tyr Val Leu Ser Leu Ala Gly Ala Asp Phe Leu Phe Leu Cys 70 75 80

Phe Gln Ile Ile Asn Cys Leu Val Tyr Leu Ser Asn Phe Phe Cys Ser 85 90 95

Ile Ser Ile Asn Phe Pro Ser Phe Phe Thr Thr Val Met Thr Cys Ala 100 105 110

Tyr Leu Ala Gly Leu Ser Met Leu Ser Thr Val Ser Thr Glu Arg Cys

Leu Ser Val Leu Trp Pro Ile Trp Tyr Arg Cys Arg Arg Pro Arg His

Leu Ser Ala Val Val Cys Val Leu Leu Trp Ala Leu Ser Leu Leu Leu 145 150 155 160

Ser Ile Leu Glu Gly Lys Phe Cys Gly Phe Leu Phe Ser Asp Gly Asp 165 170

Ser Gly Trp Cys Gln Thr Phe Asp Phe Ile Thr Ala Ala Trp Leu Ile 180 185 190

Phe Leu Phe Met Val Leu Cys Gly Ser Ser Leu Ala Leu Leu Val Arg 195 200 205

Ile Leu Cys Gly Ser Arg Gly Leu Pro Leu Thr Arg Leu Tyr Leu Thr 210 220

Ile Leu Leu Thr Val Leu Val Phe Leu Leu Cys Gly Leu Pro Phe Gly 235 235 240

Ile Gln Trp Phe Leu Ile Leu Trp Ile Trp Lys Asp Ser Asp Val Leu 245 250 255

Phe Cys His Ile His Pro Val Ser Val Val Leu Ser Ser Leu Asn Ser 260 265 270

Ser Ala Asn Pro Ile Ile Tyr Phe Phe Val Gly Ser Phe Arg Lys Gln 275 280 285

Trp Arg Leu Gln Gln Pro Ile Leu Lys Leu Ala Leu Gln Arg Ala Leu 290 295 300

| Gln 305 | Asp | Ile | Ala | Glu | Val 310 | Asp | His | Ser | Glu | Gly 315 | Cys | Phe | Arg | Gln | Gly 320 | |
|------------------------------|--------------------------|---------------------------|------|------------|------------|-------|-------|--------------|------------|------------|------|------|------|------|------------|------|
| Thr | Pro | Glu | Met | Ser 325 | Arg | Ser | Ser | Leu | Val 330 | | | | | | | |
| <210 <211 <212 <213 | 1> : 2> ! | 31 1092 DNA Homo | sap | iens | | | | | | | | | | | | |
| <400 atgg |)> ggcc | 31 ccg (| gcga | ggcg | ct g | ctgg | cgggi | t ct | cctg | gtga | tggt | tact | ggc | cgtg | gcgctg | 60 |
| | | | | | | | | | | | | | | | cgagcc | 120 |
| tca | ggcg | tcc | tcct | ggtg | aa t | ctgt | cgct | g g g | ccac | ctgc | tgc | tggc | ggc | gctg | gacatg | 180 |
| ccc | ttca | cgc | tgct | cggt | gt g | atgc | gcgg | g cg | gaca | ccgt | cgg | cgcc | cgg | cgca | tgccaa | 240 |
| gtc | attg | gct | tcct | ggac | ac c | ttcc | tggc | g tc | caac | gcgg | cgc | tgag | cgt | ggcg | gcgctg | 300 |
| agc | gcag | acc | agtg | gctg | gc a | gtgg | gctt | c cc | actg | cgct | acg | ccgg | acg | cctg | cgaccg | 360 |
| cgc | tatg | ccg | gcct | gctg | ct g | ggct | gtgc | c tg | ggga | cagt | cgc | tggc | ctt | ctca | ggcgct | 420 |
| gca | cttg | gct | gctc | gtgg | ct t | ggct | acag | c ag | cgcc | ttcg | cgt | cctg | ttc | gctg | cgcctg | 480 |
| ccg | cccg | agc | ctga | gcgt | cc g | cgct | tcgc | a go | ctto | accg | сçа | cgct | cca | tgcc | gtgggc | 540 |
| ttc | gtgc | tgc | cgct | ggcg | ıgt ç | ctct | gcct | c ac | ctcg | ctcc | agg | tgca | ccg | ggtg | gcacgc | 600 |
| agc | cact | gcc | agcg | catg | ıga o | cacco | tcac | c at | gaag | gcgc | tcg | cgct | gct | cgcc | gacctg | 660 |
| cac | ссса | igtg | tgcg | gcag | jcg (| ctgcc | tcat | c ca | gcaç | jaagc | ggc | gccg | cca | ccgc | gccacc | 720 |
| agg | aaga | ittg | gcat | tgct | at t | gcga | acctt | c ct | cato | etget | ttg | cccc | gta | tgto | atgacc | 780 |
| agg | ctgg | ıcgg | agct | cgtç | jcc (| cttc | gtcac | c gt | gaad | gccc | agt | gggg | cat | cct | agcaag | 840 |
| tgo | ctga | cct | acag | caaç | ggc (| ggtgg | geega | ic c | gtto | cacgt | act | ctct | gct | ccg | cggccg | 900 |
| | | | | | | | | | | | | | | | ccagca | |
| | | | | | | | | | | | | | • | | gagaacc | |
| ccç | gege | ccag | cgto | ccac | cca (| caac | ggcto | t g | tggad | cacag | aga | atga | attc | ctg | cctgcaç | 1080 |
| caç | gacad | cact | ga | | | | | | | | | | | | | 1092 |
| <21 <23 <23 | 10> 11> 12> 13> | 32 363 PRT | o sa | pien | s | | | | | | | | | | | |

<400> 32

Met Gly Pro Gly Glu Ala Leu Leu Ala Gly Leu Leu Val Met Val Leu l $1 \\ 5 \\ 10 \\ 15$

Ala Val Ala Leu Leu Ser Asn Ala Leu Val Leu Leu Cys Cys Ala Tyr 20 25 30

Ser Ala Glu Leu Arg Thr Arg Ala Ser Gly Val Leu Leu Val Asn Leu 35 40 45

Ser Leu Gly His Leu Leu Leu Ala Ala Leu Asp Met Pro Phe Thr Leu 50 55 60

Leu Gly Val Met Arg Gly Arg Thr Pro Ser Ala Pro Gly Ala Cys Gln 65 70 75 80

Val Ile Gly Phe Leu Asp Thr Phe Leu Ala Ser Asn Ala Ala Leu Ser 85 90 95

Val Ala Ala Leu Ser Ala Asp Gln Trp Leu Ala Val Gly Phe Pro Leu 100 105 110

Arg Tyr Ala Gly Arg Leu Arg Pro Arg Tyr Ala Gly Leu Leu Gly 115 120 125

Cys Ala Trp Gly Gln Ser Leu Ala Phe Ser Gly Ala Ala Leu Gly Cys 130 135 140

Ser Trp Leu Gly Tyr Ser Ser Ala Phe Ala Ser Cys Ser Leu Arg Leu 145 150 155 160

Pro Pro Glu Pro Glu Arg Pro Arg Phe Ala Ala Phe Thr Ala Thr Leu 165 170 175

His Ala Val Gly Phe Val Leu Pro Leu Ala Val Leu Cys Leu Thr Ser 180 185 190

Leu Gln Val His Arg Val Ala Arg Ser His Cys Gln Arg Met Asp Thr 195 200 205

Val Thr Met Lys Ala Leu Ala Leu Leu Ala Asp Leu His Pro Ser Val 210 215 220

Arg Gln Arg Cys Leu Ile Gln Gln Lys Arg Arg Arg His Arg Ala Thr 225 230 235 240

Arg Lys Ile Gly Ile Ala Ile Ala Thr Phe Leu Ile Cys Phe Ala Pro 245 250

Tyr Val Met Thr Arg Leu Ala Glu Leu Val Pro Phe Val Thr Val Asn 260 265 270

Ala Gln Trp Gly Ile Leu Ser Lys Cys Leu Thr Tyr Ser Lys Ala Val 275 280 285

Ala Asp Pro Phe Thr Tyr Ser Leu Leu Arg Arg Pro Phe Arg Gln Val 290 295 300

Leu Ala Gly Met Val His Arg Leu Leu Lys Arg Thr Pro Arg Pro Ala 305 310 315 320

Ser Thr His Asp Ser Ser Leu Asp Val Ala Gly Met Val His Gln Leu 325 330 335

Leu Lys Arg Thr Pro Arg Pro Ala Ser Thr His Asn Gly Ser Val Asp 340 345 350

Thr Glu Asn Asp Ser Cys Leu Gln Gln Thr His 355

<210> 33 <211> 1125

| <212> DNA <213> Homo | sapiens | | | | | |
|-------------------------|--------------|------------|--------------------|------------|------------|------|
| <400> 33 atgcccacac | tcaatacttc | tgcctctcca | cccacattct | tctgggccaa | tgcctccgga | 60 |
| | | tgatgctccg | | | | 120 |
| atggttgccc | tggcctatgg | gcttgtgggg | gccattggct | tgctgggaaa | tttggcggtg | 180 |
| ctgtgggtac | tgagtaactg | tgcccggaga | gcccctggcc | caccttcaga | caccttcgtc | 240 |
| ttcaacctgg | ctctggcgga | cctgggactg | gcactcactc | tccccttttg | ggcagccgag | 300 |
| tcggcactgg | actttcactg | gcccttcgga | ggtgccctct | gcaagatggt | tctgacggcc | 360 |
| actgtcctca | acgtctatgc | cagcatcttc | ctcatcacag | cgctgagcgt | tgctcgctac | 420 |
| tgggtggtgg | ccatggctgc | ggggccaggc | acccacctct | cactettetg | ggcccgaata | 480 |
| gccaccctgg | cagtgtgggc | ggcggctgcc | ctggtgacgg | tgcccacagc | tgtcttcggg | 540 |
| | | tgtgcgcctt | | | | 600 |
| | | gagggtggtg | | | | 660 |
| | | gctggccttc | | | | 720 |
| | | tgtccgcatc | | | | 780 |
| | | ctggggtgtc | | | | 840 |
| | | gacgtatgtc | | | | 900 |
| | | | | | gcaggctctg | 960 |
| | | | | | ctgggtgcaa | 1020 |
| caggtggccc | : taaagcaggt | aggcaggcgg | tg ggtcgcaa | gcaacccccg | ggagagscgc | 1080 |
| ccttctaccc | tgctcaccaa | cctggacaga | gggacaccco | g ggtga | | 112 |
| | | | | | | |

<210> 34 <211> 374 <212> PRT <213> Homo sapiens

<400> 34

Met Pro Thr Leu Asn Thr Ser Ala Ser Pro Pro Thr Phe Phe Trp Ala 1 5 10

Asn Ala Ser Gly Gly Ser Val Leu Ser Ala Asp Asp Ala Pro Met Pro 20 25 30

Val Lys Phe Leu Ala Leu Arg Leu Met Val Ala Leu Ala Tyr Gly Leu 35 40 45

Val Gly Ala Ile Gly Leu Leu Gly Asn Leu Ala Val Leu Trp Val Leu 50 60

Ser Asn Cys Ala Arg Arg Ala Pro Gly Pro Pro Ser Asp Thr Phe Val 65 70 75 80

| 211 | e As | II Te | eu A. | 8. | eu A 5 | la A | Asp | Leu | ı G1; | y Le 90 | u Al | a Le | u Tł | ır L∈ | u Pr 95 | o Phe |
|------------|------------|------------|-------------|-------------------|-----------|------------|-----------|------------|------------|------------|------|--------------|------------|--------------|------------|--------------|
| Trį | o Al | a Al | la G: 10 | lu Se 00 | er A | la I | Leu | Asp | Phe 105 | e Hi | s Tr | p Pr | o Pi | e Gl | | y Ala |
| Leu | 1 СА | s Ly 11 | s Me | et Va | al Le | eu T | hr | Ala 120 | Thi | va: | l Le | u As | n Va 12 | 1 Ту 5 | r Al | a Ser |
| Ile | 2 Ph | e Le O | u Il | .e Ti | nr Al | la I 1 | eu .35 | Ser | Va] | Ala | a Ar | g Ty | r Tr O. | p Va | l Va | l Ala |
| Met 145 | Ala | a Al | a Gl | y Pr | o GI | lу Т 50 | hr | His | Leu | Ser | Le: | u Phe 5 | ∋ Tr | p Al | a Ar | g Ile 160 |
| Ala | Thi | : Le | u Al | a Va 16 | 1 Tr | рА | la | Ala | Ala | Ala 170 | Let | u Val | L Th | r Va | l Pro | o Thr |
| Ala | Va] | . Ph | e Gl 18 | y V a 0 | 1 G1 | u G | lу | Glu | Val 185 | Суѕ | Gly | y Val | Ar | J Let 190 | | s Leu |
| Leu | Arg | 19! | e Pr | o Se | r Ar | g T | yr | Trp 200 | Leu | Gly | Ala | а Туг | Gl: 20 | ı Let | ı Glr | n Arg |
| Val | Val 210 | Let | ı Al | a Ph | e Me | t Va 2: | al 15 | Pro | Leu | Gly | Va 1 | . Ile 220 | Thi | Thr | Ser | Tyr |
| 223 | | | | | 23 | U | | | | | 235 | • | | | | Asp 240 |
| | | | | 24. | , | | | | | 250 | | | | | 255 | |
| | | | 200 | , | | | | | 265 | | | | | 270 | | Val |
| Lys | Phe | Asp 275 | Leu | ı Val | l Pro | o Tr | ъ Р 2 | Asn 280 | Ser | Thr | Phe | Tyr | Thr 285 | Ile | Gln | Thr |
| | | | | | | 29 | 5 | | | | | 300 | | | | Leu |
| 505 | | | | | 310 | ' | | | | | 315 | Pro | | J | | 320 |
| | | | | 323 | | | | | | 330 | | Trp | | | 335 | |
| | | | 340 | | | | | • | 345 | | | Gly | | 350 | | |
| Ala | Ser | Asn 355 | Pro | Arg | Glu | Se: | r A 3 | rg 1 60 | Pro | Ser | Thr | Leu | Leu 365 | Thr | Asn | Leu |
| | _ | | | | | | | | | | | | | | | |

Asp Arg Gly Thr Pro Gly 370

<210> 35 <211> 1092 <212> DNA <213> Homo sapiens

atgaatcggc accatctgca ggatcacttt ctggaaatag acaagaagaa ctgctgtgtg

60

Page 34

| ttccgagatg | acttcattgt | caaggtgttg | ccgccggtgt | tggggctgga | gtttatcttc | 120 |
|------------|------------|------------|------------|------------|------------|------|
| | gcaatggcct | | | | | 180 |
| | tttcctgtt | | | | | 240 |
| | tggacaacta | | | | | 300 |
| | tcttcatgtt | | | | | 360 |
| | acaggtattt | | | | | 420 |
| | cagccatcat | | | | | 480 |
| | agaagaagat | | | | | 540 |
| | ataccttcca | | | | | 600 |
| | tcctgttctg | | | | | 660 |
| | ccaagatcaa | | | | | 720 |
| | tccttcccag | | | | | 780 |
| | agaattgtga | | | | | 840 |
| | acatgaacag | | | | | 900 |
| | tcttctccac | | | | * | 960 |
| | | | | | aaccagaggc | 1020 |
| | | | | | tctgggccca | 1080 |
| acctctcctt | | | | | | 1092 |
| | | | | | | |

<210> 36

<211> 363 <212> PRT

<213> Homo sapiens

<400> 36

Met Asn Arg His His Leu Gln Asp His Phe Leu Glu Ile Asp Lys Lys 1 5 10 15

Asn Cys Cys Val Phe Arg Asp Asp Phe Ile Val Lys Val Leu Pro Pro 20 25 30

Val Leu Gly Leu Glu Phe Ile Phe Gly Leu Leu Gly Asn Gly Leu Ala 35 40 45

Leu Trp Ile Phe Cys Phe His Leu Lys Ser Trp Lys Ser Ser Arg Ile 50 55

Phe Leu Phe Asn Leu Ala Val Ala Asp Phe Leu Leu Ile Ile Cys Leu 65 70 75 80

Pro Phe Leu Met Asp Asn Tyr Val Arg Arg Trp Asp Trp Lys Phe Gly 85

Asp Ile Pro Cys Arg Leu Met Leu Phe Met Leu Ala Met Asn Arg Gln 100 105 110

| GI | у зе | 11 11 | .e 11 .5 | e Ph | e Le | u Th | r Val 120 | l Va D | l Al | a Va | l As | p Ar | g Ту 5 | r Ph | e Arg | |
|----------------------|------------|------------|--------------|--------------|--------------|--------------|--------------|------------|--------------|------------|----------------|------------|------------|--------------|--------------|--|
| Va | l Va 13 | 1 Hi | s Pr | o Hi | s Hi | s Ala 135 | a Lei | ı As | n Ly. | s Il | e Se 14 | r Ası O | n Ar | g Th | r Ala | |
| A1. | a Il 5 | e Il | e Se | r Cy | s Le: 150 | u Lei 0 | Trp | Gl; | y Ile | e Th. | r Ile 5 | e Gly | y Le | u Thi | r Val 160 | |
| Hi: | s Le | u Le | u Ly: | s Lys 165 | s Lys | s Met | Pro | 110 | ≘ Glr 17(| n Ası | n Gly | λ GJ? | / Ala | a Asr 175 | n Leu | |
| Суз | s Se | r Se | r Phe 180 | e Sei | : Ile | e Cys | His | Th: | Phe | e Glr | ı Trp | His | Glu 190 | ı Ala | Met | |
| Phe | e Lei | Lei 195 | u Glu 5 | ı Phe | Phe | e Leu | Pro 200 | Let | Gly | ' Ile | e Ile | Leu 205 | Phe | Cys | Ser | |
| Ala | 210 | ı Ile | ∋ Ile | Trp | Ser | Leu 215 | Arg | Gln | Arg | Gln | Met 220 | Asp | Arg | His | Ala | |
| Lys 225 | Ile | Lys | Arg | Ala | Ile 230 | Thr | Phe | Ile | Met | Val 235 | Val | Ala | Ile | Val | Phe 240 | |
| Val | Ile | Cys | Phe | Leu 245 | Pro | Ser | Val | Val | Val 250 | Arg | Ile | Arg | Ile | Phe 255 | Trp | |
| Leu | Leu | His | Thr 260 | Ser | Gly | Thr | Gln | Asn 265 | Cys | Glu | Val | Tyr | Arg 270 | Ser | Val | |
| Asp | Leu | Ala 275 | Phe | Phe | Ile | Thr | Leu 280 | Ser | Phe | Thr | Tyr | Met 285 | Asn | Ser | Met | |
| Leu | Asp 290 | Pro | Val | Val | Tyr | Tyr 295 | Phe | Ser | Ser | Pro | Ser 300 | Phe | Pro | Asn | Phe | |
| Phe 305 | Ser | Thr | Leu | Ile | Asn 310 | Arg | Cys | Leu | Gln | Arg 315 | Lys | Met | Thr | Gly | Glu 320 | |
| Pro | Asp | Asn | Asn | Arg 325 | Ser | Thr | Ser | Val | Glu 330 | Leu | Thr | Gly | Asp | Pro 335 | Asn | |
| Lys | Thr | Arg | Gly 340 | Ala | Pro | Glu . | Ala | Leu 345 | Met | Ala | Asn | Ser | Gly 350 | Glu | Pro | |
| Trp | Ser | Pro 355 | Ser | Tyr | Leu | Gly i | Pro : | Thr | Ser | Pro | | | • | | | |
| <210 <211 <212 | > 1 | 7 044 | | | | | | | | | | | | | | |
| <213 | _ | NA omo | sapi | ens | | | | | | | | | | | | |
| <400: atgg | _ | | gctg | gcaco | : ttg | gccct | gtg | ggca | actac | cao d | ettar | accaa | ור כי | rtas+ | tccag | |
| | | | | | | | | | | | | | | | gggg | |
| gacct | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | gctg | |
| cgaca | | | | | | | | | | | | | | | | |
| ctcct | ccac | a to | gctca | itctc | ctc | cago | agc · | ctga | gtaa | ct o | Iggan | ctaa | a cc | acet | aacc | |
| | | | | | | J - | - | - 55 | J-79 | | iggag age 3 | | y cc | ycat | yycc | |
| | | | | | | | | | | | | | | | | |

| | | | | | atacttcacc | 420 |
|------------|------------|------------|------------|------------|------------|------|
| tgtggcattc | tcactgatgc | tgtcttcgcc | gcctgcacca | geaecatect | gecereace | |
| gccattgtgc | tgcacaccta | cctggcagtc | atccatccac | tgcgctacct | ctccttcatg | 480 |
| | | | ctcatctggc | | | 540 |
| | | | gatgcccagc | | | 600 |
| | | | ccgggatgtg | | | 660 |
| | | | tgcacagctc | | | 720 |
| | | | atctgggggc | | | 780 |
| | | | acattgtacg | | | 840 |
| | | | cacattgact | | | 900 |
| | | | cttccccgtg | | | 960 |
| | | | | | caggaggcac | 1020 |
| | ttaccatttc | | | | | 1044 |
| caggecatet | | = | | | | |

<210> 38 <211> 347 <212> PRT

<213> Homo sapiens

<400> 38

Met Gly Asp Glu Leu Ala Pro Cys Pro Val Gly Thr Thr Ala Trp Pro 1 5 10 15

Ala Leu Ile Gln Leu Ile Ser Lys Thr Pro Cys Met Pro Gln Ala Ala 20 25 30

Ser Asn Thr Ser Leu Gly Leu Gly Asp Leu Arg Val Pro Ser Ser Met 35

Leu Tyr Trp Leu Phe Leu Pro Ser Ser Leu Leu Ala Ala Ala Thr Leu 50 60

Ala Val Ser Pro Leu Leu Leu Val Thr Ile Leu Arg Asn Gln Arg Leu 65 70 75 80

Arg Gln Glu Pro His Tyr Leu Leu Pro Ala Asn Ile Leu Leu Ser Asp 85 90 95

Leu Ala Tyr Ile Leu Leu His Met Leu Ile Ser Ser Ser Leu Gly 100 105

Gly Trp Glu Leu Gly Arg Met Ala Cys Gly Ile Leu Thr Asp Ala Val 115 120 125

Phe Ala Ala Cys Thr Ser Thr Ile Leu Ser Phe Thr Ala Ile Val Leu 130 135 140

His Thr Tyr Leu Ala Val Ile His Pro Leu Arg Tyr Leu Ser Phe Met 145 150 155 160

Ser His Gly Ala Ala Trp Lys Ala Val Ala Leu Ile Trp Leu Val Ala 165 170 175

| Cys | s Cys | Phe | Pro 180 | Thr | ? Phe | Leu | Ile | Trp 185 | Leu | Ser | Lys | Trp | Gln 190 | | Ala | |
|----------------------|------------|----------------|------------|------------|------------|------------|------------|------------|------------|------------|---------------|------------|------------|------------|------------|-----|
| Glr | Leu | Glu 195 | Glu | Gln | Gly | Ala | Ser 200 | Туг | Ile | Leu | Pro | Pro 205 | Ser | Met | Gly | |
| Thr | Gln 210 | Pro | Gly | Cys | Gly | Leu 215 | Leu | Val | Ile | Val | Thr 220 | Tyr | Thr | Ser | Ile | |
| Leu 225 | Суз | Val | Leu | Phe | Leu 230 | Cys | Thr | Ala | Leu | Ile 235 | Ala | Asn | Cys | Phe | Trp 240 | |
| Arg | Ile | Tyr | Ala | Glu 245 | Ala | Lys | Thr | Ser | Gly 250 | Ile | Trp | Gly | Gln | Gly 255 | Tyr | |
| Ser | Arg | Ala | Arg 260 | Gly | Thr | Leu | Leu | Ile 265 | His | Ser | Val | Leu | Ile 270 | Thr | Leu | |
| Tyr | Val | Ser 275 | Thr | Gly | Val | Val | Phe 280 | Ser | Leu | Asp | Met | Val 285 | Leu | Thr | Arg | |
| Туr | His 290 | His | Ile | Asp | Ser | Gly 295 | Thr | His | Thr | Trp | Leu 300 | Leu | Ala | Ala | Asn | |
| Ser 305 | Glu | Val | Leu | Met | Met 310 | Leu | Pro | Arg | Ala | Met 315 | Leu | Pro | Tyr | Leu | Tyr 320 | |
| Leu | Leu | Arg | Tyr . | Arg 325 | Gln | Leu : | Leu | Gly | Met 330 | Val . | Arg | Gly I | | Leu 335 | Pro | |
| Ser | Arg . | Arg | His (| Gln . | Ala | Ile 1 | | Thr : | He: | Ser | | | | | | |
| <210 <211 <212 | > 16 | 9 023 NA | | | | | | | | | | | | | | |
| <213 | > Ho | omo s | sapie | ens | | | | | | | | | | | | |
| <400: atgaa | | | cato | gcato | tte | gttgg | jaac | acct | ctac | са а | acti | ttaa | a ca | 22+0 | ctgg | 60 |
| aataa | | | | | | | | | | | | | | | | 60 |
| attg | | | | | | | | | | | | | | | | 120 |
| agato | | | | | | | | | | | | | | | | 180 |
| gtcca | | | | | | | | | | | | | | | | 240 |
| t ttg g | | | | | | | | | | | | | | | | 300 |
| | | | | | | | | | | | | | | | | 360 |
| agtgo | acgt | t. aa: | апаа | caac | gug | care: | gac | ayyı. | actt | Eg c | cctc | gtcca | a ac | catt | tcga | 420 |
| ctgac | tato | - gg. | rcati | taca | 9 ta | caay | acc | atcc | ggato | ca a | tttg | ggcct | tte | gggc | agct | 480 |
| tcctt | rant! | t ata | 3CaL | - - | Lgt | ctgg | gtc | tacto | cgaaq | gg to | catca | aaatt | taa | aaga | cggt | 540 |
| gttga | gage | . gr | | ecga | ttt. | gacat | cc (| cctga | cgat | tg ta | actci | ggta | tac | cacti | ttat | 600 |
| ttata | galdi | a caa | ctt | .cct | ttt | ccct | cta d | cctt | gatt | t to | ggtgt | gcta | tat | ttta | aatt | 660 |
| ttatg | Lata | ctt | ggga | gat | gtai | tcaac | ag a | aataa | ggat | g co | cagat | gctg | caa | tccc | agt | 720 |
| gtacca | aaaac | aga | ıgagt | gat | gaaq | gttga | ica a | aagat | ggtg | | gtgc je 38 | | ggt | agto | ttt | 780 |

| atcctgagtg ctgcccctta tcatgtgata caactggtga acttacagat ggaacagcco | = | | | | | | | | | | | |
|--|---|--|--|--|--|--|--|--|--|--|--|--|
| acactggcct tctatgtggg ttattacctc tccatctgtc tcagctatgc cagcagcagc | | | | | | | | | | | | |
| attaaccett ttetetacat eetgetgagt ggaaatttee agaaaegtet geeteaaate | | | | | | | | | | | | |
| caaagaagag cgactgagaa ggaaatcaac aatatgggaa acactctgaa atcacactt | | | | | | | | | | | | |
| | | | | | | | | | | | | |
| tag | | | | | | | | | | | | |
| <210> 40 <211> 340 <212> PRT <213> Homo sapiens | | | | | | | | | | | | |
| <400> 40 | | | | | | | | | | | | |
| Met Asn Pro Phe His Ala Ser Cys Trp Asn Thr Ser Ala Glu Leu Leu 1 5 10 15 | | | | | | | | | | | | |
| Asn Lys Ser Trp Asn Lys Glu Phe Ala Tyr Gln Thr Ala Ser Val Val 20 25 30 | | | | | | | | | | | | |
| Asp Thr Val Ile Leu Pro Ser Met Ile Gly Ile Ile Cys Ser Thr Gly 35 40 45 | | | | | | | | | | | | |
| Leu Val Gly Asn Ile Leu Ile Val Phe Thr Ile Ile Arg Ser Arg Lys 50 60 | | | | | | | | | | | | |
| Lys Thr Val Pro Asp Ile Tyr Ile Cys Asn Leu Ala Val Ala Asp Leu 65 70 75 80 | | | | | | | | | | | | |
| Val His Ile Val Gly Met Pro Phe Leu Ile His Gln Trp Ala Arg Gly 85 90 95 | | | | | | | | | | | | |
| Gly Glu Trp Val Phe Gly Gly Pro Leu Cys Thr Ile Ile Thr Ser Leu 100 105 110 | | | | | | | | | | | | |
| Asp Thr Cys Asn Gln Phe Ala Cys Ser Ala Ile Met Thr Val Met Ser 115 120 125 | | | | | | | | | | | | |
| Val Asp Arg Tyr Phe Ala Leu Val Gln Pro Phe Arg Leu Thr Arg Trp 130 135 140 | | | | | | | | | | | | |
| Arg Thr Arg Tyr Lys Thr Ile Arg Ile Asn Leu Gly Leu Trp Ala Ala 145 150 150 155 160 | | | | | | | | | | | | |
| Ser Phe Ile Leu Ala Leu Pro Val Trp Val Tyr Ser Lys Val Ile Lys 165 170 175 | | | | | | | | | | | | |
| Phe Lys Asp Gly Val Glu Ser Cys Ala Phe Asp Leu Thr Ser Pro Asp 180 185 190 | , | | | | | | | | | | | |
| Asp Val Leu Trp Tyr Thr Leu Tyr Leu Thr Ile Thr Thr Phe Phe Phe 195 200 205 | 2 | | | | | | | | | | | |
| Pro Leu Pro Leu Ile Leu Val Cys Tyr Ile Leu Ile Leu Cys Tyr Thr 210 215 220 | : | | | | | | | | | | | |
| Trp Glu Met Tyr Gln Gln Asn Lys Asp Ala Arg Cys Cys Asn Pro Ser 225 230 235 240 | 5 | | | | | | | | | | | |
| Val Pro Lys Gln Arg Val Met Lys Leu Thr Lys Met Val Leu Val Leu Page 39 | 1 | | | | | | | | | | | |

WO 01/36471 245 250 255 Val Val Val Phe Ile Leu Ser Ala Ala Pro Tyr His Val Ile Gln Leu 260 265 Val Asn Leu Gln Met Glu Gln Pro Thr Leu Ala Phe Tyr Val Gly Tyr Tyr Leu Ser Ile Cys Leu Ser Tyr Ala Ser Ser Ser Ile Asn Pro Phe Leu Tyr Ile Leu Leu Ser Gly Asn Phe Gln Lys Arg Leu Pro Gln Ile Gln Arg Arg Ala Thr Glu Lys Glu Ile Agn Asn Met Gly Asn Thr Leu 330 Lys Ser His Phe 340 <210> 41 <211> 24 <212> DNA <213> Artificial Sequence <220> <221> misc_feature
<223> Novel Sequence <400> 41 cttgcagaca tcaccatggc agcc 24 <210> 42 <211> 24 <212> DNA <213> Artificial Sequence <220> <221> misc_feature <223> Novel Sequence <400> 42 gtgatgctct gagtactgga ctgg 24

PCT/US00/31509

<210> 43
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<221> misc_feature
<223> Novel Sequence

<400> 43
gaagctgtga agagtgatgc 20

<210> 44 <211> 24 <212> DNA

Page 40

| <213> | Artificial Sequence | | |
|----------------------------------|--|---------|----|
| <220> <221> <223> | misc_feature Novel Sequence | | |
| <400> gtcagca | 44 aata ttgataagca gcag | | 24 |
| <210> <211> <212> <213` | | | |
| <220> <221> <223> | misc_feature Novel Sequence | | |
| <400> ccatgg | 45 ggaa cgattctgtc agctacg | | 27 |
| <211> <212> | | | |
| | misc_feature Novel Sequence | | |
| <400> gctato | 46 cctg aagccagtet tgtg | | 24 |
| <210> <211> <212> <213> | 47 26 DNA Artificial Sequence | | |
| <220> <221> <223> | - | • | |
| <400> ccagg | 47 atgtt gtgtcaccgt ggtggc | | 26 |
| <210> <211> <212> <213> | 26 DNA | | |
| <220> <221> <223> | misc_feature | | |
| <400> cacag | 48 cgctg cagccctgca gctggc | Page 41 | 26 |

| WO 01/36471 | |
|-------------|---------------|
| | PCT/US00/3150 |

| <210 | | |
|--------|--------------------------|----|
| <211 | | |
| <212 | | |
| <213 | > Artificial Sequence | |
| | | |
| <220 | | |
| <221 | > misc_feature | |
| <223 | > Novel Sequence | |
| | | |
| <400 | > 49 | |
| | ctctcg tagggatgaa ccagac | |
| | cagggacgaa ccagac | 26 |
| | | |
| <210 | > 50 | |
| <211> | > 26 | |
| <212> | DNA | |
| <213> | Artificial Sequence | |
| | | |
| <220> | | |
| <221> | misc_feature | |
| <223> | Novel Sequence | |
| | · | |
| <400> | 50 | |
| | | |
| cccgc | acagg tgggaagcac ctgtgg | 26 |
| | | |
| <210> | 51 | |
| <211> | 23 | |
| <212> | | |
| <213> | Artificial Sequence | |
| | | |
| <220> | | |
| <221> | misc_feature | |
| <223> | Novel Sequence | |
| | | |
| <400> | 51 | |
| <400> | | |
| gcctgt | gaca ggaggtaccc tgg | 23 |
| | | |
| <210> | 52 | |
| <211> | | |
| <212> | | |
| <213> | | |
| | | |
| <220> | | |
| <221> | misc_feature | |
| <223> | Novel Sequence | |
| | | |
| < 400s | | |
| <400> | | |
| catate | cctc cgagtgtcca gcggc | 25 |
| | | |
| <210> | 53 | |
| <211> | 31 | |
| <212> | DNA | |
| <213> | Artificial Sequence | |
| | | |
| <220> | | |
| <221> | misc_feature | |
| | | |

Page 42

| , | WO 01/36471 | PC 1/0500 |
|----------------------------------|-------------------------------------|-----------|
| <223> | Novel Sequence | |
| <400> gcatgg | 53 Jagag aaaatttatg toottgoaac o | 31 |
| <210><211><211><212><212><213> | | |
| <220> <221> | misc_feature Provel Sequence | |
| <400> caagaa | 54 acagg teteatetaa gagetee | 27 |
| <210><211><211><212><213> | 26 DNA | |
| | misc_feature Novel Sequence | |
| <400> gctgt | 55 tgcca tgacgtccac ctgcac | 26 |
| <210><211><211><212><213> | 26 DNA | |
| <220> <221> <223> | misc_feature | |
| <400> ggaca | · 56 agttca aggtttgcct tagaac | 26 |
| <2103 <2113 <2123 <2133 | > 23 > DNA | |
| <2203 <2213 <2233 | > misc_feature | |
| <400 | > 57 cgatac tgctcctatg ctc | 23 |
| <210 <211 | > 58 > 26 | |

| <212 | > DNA | |
|----------------|---------------------------------------|----|
| <213 | > Artificial Sequence | |
| | | |
| <220 | | |
| <221 | > misc_feature | |
| <223 | > Novel Sequence | |
| | | |
| <400 | > 58 | |
| | tccact gaaagtccag tgatcc | |
| g cug. | secure galagicolag igator | 26 |
| | | |
| <210 | > 59 | |
| <211> | · 26 | |
| | DNA | |
| <213> | Artificial Sequence | |
| | | |
| <220> | | |
| <221> | misc_feature | |
| \2237 | Novel Sequence | |
| | | |
| <400> | 59 | |
| tttct | gagca tggatccaac catctc | |
| | , , , , , , , , , , , , , , , , , , , | 26 |
| | | |
| <210> | | |
| <211> | | |
| <212> | | |
| <213> | Artificial Sequence | |
| <220> | | |
| | misc_feature | |
| <223> | Novel Sequence | |
| | novor bequence | |
| | | |
| <400> | | |
| ctgtct | gaca gggcagaggc tcttc | 25 |
| | | 25 |
| <210> | 61 | |
| <211> | | |
| <212> | | |
| <213> | Artificial Sequence | |
| | · · · · · · · · · · · · · · · · · · · | |
| <220> | | |
| <221> | | |
| <223> | Nove Sequence | |
| | | |
| <400> | 61 | |
| | | |
| ggaacc | cgta tagacccagc gtcgctcc | 28 |
| | | |
| <210> | 62 | |
| <211> | 28 | |
| <212> | DNA | |
| <213> | Artificial Sequence | |
| -222 | | |
| <220> <221> | | |
| <221> <223> | misc_feature | |
| ~~~>/ | Novel Sequence | |
| | | |
| <400> | 62 | |
| | | |

Page 44

| ggaggtt | gcg ccttagcgac agatgacc | 28 |
|----------------------------------|--|----|
| <210> <211> <212> <213> | 63 22 DNA Artificial Sequence | |
| <220> <221> <223> | misc_feature Novel Sequence | |
| <400> ctgcac | 63 ecgg aciettgete tg | 22 |
| <210> <211> <212> <213> | 64 25 DNA Artificial Sequence | |
| <220> <221> <223> | misc_feature Novel Sequence | |
| <400> gtctgc | 64 ttgt tcagtgccac tcaac | 25 |
| <210> <211> <212> <213> | 26 DNA | |
| | misc_feature Novel Sequence | |
| <400> tatct | 65 gcaat totattotag otootg | 26 |
| <210> <211> <212> <213> | 26 DNA | |
| <220> <221> <223> | misc_feature | |
| <400> tgtco | 66 ctaat aaagtcacat gaatgc | 26 |
| <210><211><211><212><213> | | |

<220>

| W O (///304// | | PCT/US00/31509 |
|--|--------|----------------|
| <221> misc_feature <223> Novel Sequence | | |
| <400> 67 ggagacaacc atgaatgagc | cac | 23 |
| <210> 68 <211> 24 | | |
| <212> DNA <213> Artificial Seque | ∍nce | |
| <220> <221> misc_feature <223> Novel Sequence | | |
| <400> 68 tatttcaagg gttgtttgag t | aac | 24 |
| <210> 69 <211> 27 <212> DNA <213> Artificial Seque | nce | |
| <220> <221> misc_feature <223> Novel Sequence | | |
| <400> 69 ggcaccagtg gaggttttct ga | agcatg | 27 |
| <210> 70 <211> 27 <212> DNA <213> Artificial Sequer | nce | |
| <220> <221> misc_feature <223> Novel Sequence | | |
| <400> 70 ctgatggaag tagaggctgt cc | | 27 |
| <210> 71 <211> 23 <212> DNA <213> Artificial Sequen | ce | |
| <220> <221> misc_feature <223> Novel Sequence | | |
| <400> 71 cctggcgagc cgctagcgcc ato | j | 23 |
| <210> 72 | | |

| <212> | 23 DNA Artificial Sequence | |
|----------------------------------|--|----|
| <220> <221> <223> | misc_feature Novel Sequence | |
| <400> atgage | 72 cctg ccaggccctc agt | 23 |
| <210> <211> <212> <213> | 73 27 DNA Artificial Sequence | |
| <220> <221> <223> | | |
| <400> ctgcga | 73 tgcc cacactcaat acttctg | 27 |
| <210> <211> <212> <213> | | |
| <220> <221> <223> | misc_feature Novel Sequence | |
| <400> aagga | 74 cccta cacttggtgg atctcag | 27 |
| <210> <211> <212> <213> | 22 DNA | |
| <400> gctgg | 75 agcat tcactaggcg ag | 22 |
| <210><211><211><212><213> | 24 DNA | |
| <220> <221> <223> | misc_feature | |
| <4000 agato | > 76 cctggt tcttggtgac aatg | 24 |
| <210 | > 77 | |

| <211 | > 24 | |
|----------------|-----------------------------|----|
| | > DNA | |
| <213 | > Artificial Sequence | |
| | | |
| <220 | | |
| <221 | > misc_feature | |
| <223 | > Novel Sequence | |
| | | |
| <400 | > 77 | |
| | | |
| agec | atccct gccaggaagc atgg | 24 |
| | | |
| <210 | > 78 | |
| | > 27 | |
| <212 | > DNA | |
| <213 | Artificial Sequence | |
| | · | |
| <220 | | |
| Z222 | nisc feature Novel Sequence | |
| \223/ | Novel Sequence | |
| | | |
| <400> | .78 | |
| ccaga | ctgtg gactcaagaa ctctagg | |
| | , | 27 |
| -0.0 | | |
| <210> | 79 | |
| <211> | 28 DNA | |
| | | |
| \213 / | Artificial Sequence | |
| <220> | | |
| <221> | misc feature | |
| <223> | Novel Sequence | |
| | • | |
| 400- | | |
| <400> | | |
| agtee | acgaa caatgaatcc atttcatg | 28 |
| | | |
| <210> | 80 | |
| <211> | | |
| <212> | | |
| <213> | Artificial Sequence | • |
| | · | |
| <220> | | |
| <223> | misc_feature | |
| 12237 | Novel Sequence | |
| | | |
| <400> | | |
| atcatg | tcta gactcatggt gatcc | |
| | | 25 |
| <210s | 01 | |
| <210> <211> | 81 30 | |
| <211> | | |
| <213> | DNA Artificial Company | |
| | Artificial Sequence | |
| <220> | | |
| <221> | misc feature | |
| <223> | Novel Sequence | |
| | | |

| WO 01/36471 | FC1/ |
|--|------|
| | |
| <400> 81 ggggagggaa agcaaaggtg gteeteetgg | 30 |
| <210> 82 <211> 30 <212> DNA <213> Artificial Sequence | |
| <220> <221> misc_feature <223> Novel Sequence | |
| <400> 82 ccaggagaac cacctttgct ttccctcccc | 30 |
| <210> 83 <211> 1356 <212> DNA <213> Homo sapiens | |
| <400> 83 atggagtcct cacccatccc ccagtcatca gggaactctt ccactttggg gagggtccct | 60 |
| caaaccccag gtccctctac tgccagtggg gtcccggagg tggggctacg ggatgttgct | 120 |
| toggaatotg tggccctctt cttcatgctc ctgctggact tgactgctgt ggctggcaat | 180 |
| gccgctgtga tggccgtgat cgccaagacg cctgccctcc gaaaatttgt cttcgtcttc | 240 |
| cacctetgee tggtggacet getggetgee etgaceetea tgeecetgge eatgetetee | 300 |
| agetetgee totttgacca egecetettt ggggaggtgg cetgeegeet etaettgttt | 360 |
| ctgagcgtgt gctttgtcag cctggccatc ctctcggtgt cagccatcaa tgtggagcgc | 420 |
| tactattacg tagtccaccc catgcgctac gaggtgcgca tgacgctggg gctggtggcc | 480 |
| totgtgctgg tgggtgtgtg ggtgaaggco ttggccatgg ottotgtgcc agtgttggga | 540 |
| agggtctcct gggaggaagg agetcccagt gtccccccag gctgttcact ccagtggagc | 600 |
| cacagtgcct actgccagct ttttgtggtg gtctttgctg tcctttactt tctgttgccc | 660 |
| ctgctcctca tacttgtggt ctactgcage atgttccgag tggcccgcgt ggctgccatg | 720 |
| cagcacgggc cgctgcccac gtggatggag acaccccggc aacgctccga atctctcagc | 780 |
| agccgctcca cgatggtcac cagctcgggg gcccccaga ccaccccaca ccggacgttt | 840 |
| gggggaggga aagcaaaggt ggtteteetg getgtggggg gacagtteet getetgttgg | 900 |
| ttgccctact tctctttcca cctctatgtt gccctgagtg ctcagcccat ttcaactggg | 960 |
| caggtggaga gtgtggtcac ctggattggc tacttttgct tcacttccaa ccctttcttc | 1020 |
| tatggatgtc tcaaccggca gatccggggg gagctcagca agcagtttgt ctgcttcttc | 1080 |
| aagccagctc cagaggagga gctgaggctg cctagccggg agggctccat tgaggagaac | 1140 |
| ttcctgcagt tccttcaggg gactggctgt ccttctgagt cctgggtttc ccgacccta | 1200 |
| cccagccca agcaggagcc acctgctgtt gactttcgaa tcccaggcca gatagctgag | 1260 |
| cccagcccca agcaggagcc acctgctgtt gacttegga tectaggar tectaggar page 49 | |

1356

gagacetetg agtteetgga geageaacte accagegaea teateatgte agacagetae ctccgtcctg ccgcctcacc ccggctggag tcatga <210> 84 <211> 451 <212> PRT <213> Homo sapiens <400> 84 Met Glu Ser Ser Pro Ile Pro Gln Ser Ser Gly Asn Ser Ser Thr Leu Gly Arg Val Pro Gln Thr Pro Gly Pro Ser Thr Ala Ser Gly Val Pro Glu Val Gly Leu Arg Asp Val Ala Ser Glu Ser Val Ala Leu Phe Phe Met Leu Leu Asp Leu Thr Ala Val Ala Gly Asn Ala Ala Val Met Ala Val Ile Ala Lys Thr Pro Ala Leu Arg Lys Phe Val Phe His Leu Cys Leu Val Asp Leu Leu Ala Ala Leu Thr Leu Met Pro Leu Ala Met Leu Ser Ser Ser Ala Leu Phe Asp His Ala Leu Phe Gly Glu Val Ala Cys Arg Leu Tyr Leu Phe Leu Ser Val Cys Phe Val Ser Leu 120 Ala Ile Leu Ser Val Ser Ala Ile Asn Val Glu Arg Tyr Tyr Val Val His Pro Met Arg Tyr Glu Val Arg Met Thr Leu Gly Leu Val Ala Ser Val Leu Val Gly Val Trp Val Lys Ala Leu Ala Met Ala Ser Val Pro Val Leu Gly Arg Val Ser Trp Glu Glu Gly Ala Pro Ser Val Pro 185 Pro Gly Cys Ser Leu Gln Trp Ser His Ser Ala Tyr Cys Gln Leu Phe Val Val Val Phe Ala Val Leu Tyr Phe Leu Leu Pro Leu Leu Leu Ile Leu Val Val Tyr Cys Ser Met Phe Arg Val Ala Arg Val Ala Ala Met Gln His Gly Pro Leu Pro Thr Trp Met Glu Thr Pro Arg Gln Arg Ser Glu Ser Leu Ser Ser Arg Ser Thr Met Val Thr Ser Ser Gly Ala Pro 265 Gln Thr Thr Pro His Arg Thr Phe Gly Gly Gly Lys Ala Lys Val Val Page 50

| | | 275 | | | | | 280 | | | | | 285 | | | | |
|--------------------------|--------------------------|---------------------------|------------|------------|------------|------------|------------|------------|------------|------------|--------------|------------|------------|------------|------------|-----|
| Leu | Leu 290 | Ala | Val | Gly | Gly | Gln 295 | Phe | Leu | Leu | Cys | Trp 300 | Leu | Pro | Tyr | Phe | |
| Ser 305 | Phe | His | Leu | Tyr | Val 310 | Ala | Leu | Ser | Ala | Gln 315 | Pro | Ile | Ser | Thr | Gly 320 | |
| Gln | Val | Glu | Ser | Val 325 | Val | Thr | Trp | Ile | Gly 330 | Tyr | Phe | Cys | Phe | Thr 335 | Ser | |
| Asn | Pro | Phe | Phe 340 | Tyr | Gly | Cys | Leu | Asn 345 | Arg | Gln | Ile | Arg | Gly 350 | Glu | Leu | |
| Ser | Lys | Gln 355 | Phe | Val | Cys | Phe | Phe 360 | Lys | Pro | Ala | Pro | Glu 365 | Glu | G! u | Leu | |
| Arg | Leu 370 | Pro | Ser | Arg | Glu | Gly 375 | Ser | Ile | Glu | Glu | Asn 380 | Phe | Leu | Gln | Phe | |
| Leu 385 | Gln | Gly | Thr | Gly | Cys 390 | Pro | Ser | Glu | Ser | Trp 395 | Val | Ser | Arg | Pro | Leu 400 | |
| Pro | Ser | Pro | Lys | Gln 405 | Glu | Pro | Pro | Ala | Val 410 | Asp | Phe | Arg | Ile | Pro 415 | Gly | |
| Gln | Ile | Ala | Glu 420 | | Thr | Ser | Glu | Phe 425 | Leu | Glu | Gln | Gln | Leu 430 | Thr | Ser | |
| Asp | Ile | 11e 435 | | Ser | Asp | Ser | Tyr 440 | Leu | Arg | Pro | Ala | Ala 445 | Ser | Pro | Arg | |
| Leu | Glu 450 | Ser | | | | | | | | | | | | | | |
| <21 <21 <21 <21 | 1> 2> | 85 28 DNA Homo | sap | iens | | | | | | | | | | | | |
| <40 cag | | 85 gca | aaga | ccac | ca t | cato | atc | | | | | | | | | 28 |
| <21 | 0> | 86 | | | | | | | | | | | • | | | |
| <21 <21 <21 | 2> | 28 DNA Homo | sap | oiens | 6 | | | | | | | | | | | |
| <40 gat | | 86 gatg | gtgg | jtctt | tg (| cctto | ctg | | | | | | | | | 28 |
| <21 <21 | 10> 11> 12> 13> | 87 1043 DNA Homo | l o sap | pien | S | | | | | | | | | | | |
| <40 ato |)0> ggaga | 87 agaa | aati | ttat | gtc (| cttg | caaco | ca t | ccat | ctcc | g ta | tcaga | aaat | gga | accaaat | 60 |
| | | | | | | | | | | | | | | | agaattt | 120 |
| tt | ccca | attg | tat | atct | gat | aata | tttt | tc t | gggg | agtc | t t g | ggaa | atgg | gtt | gtccata | 180 |

| tatgttttco | tgcagcctta | taagaagtcc | acatctgtga | acgttttcat | gctaaatctg | 240 |
|------------|--------------|------------|------------|------------|------------|-------|
| gccatttcag | g atctcctgtt | cataagcacg | cttcccttca | gggctgacta | ttatcttaga | 300 |
| ggctccaatt | ggatatttgg | agacctggcc | tgcaggatta | tgtcttattc | cttgtatgtc | 360 |
| aacatgtaca | gcagtattta | tttcctgacc | gtgctgagtg | ttgtgcgttt | cctggcaatg | 420 |
| gttcacccct | ttcggcttct | gcatgtcacc | agcatcagga | gtgcctggat | cctctgtggg | 480 |
| atcatatgga | tccttatcat | ggcttcctca | ataatgctcc | tggacagtgg | ctctgagcag | 540 |
| aacggcagtg | tcacatcatg | cttagagctg | aatctctata | aaattgctaa | gctgcagacc | 600 |
| atgaactata | ttgccttggt | ggtgggctgc | ctgctgccat | ttttcacact | cagcatctgt | 660 |
| tatctgctga | tcattcgggt | tctgttaaaa | gtggaggtcc | cagaatcggg | gctgcgggtt | 720 |
| tctcacagga | aggcaaagac | caccatcatc | atcaccttga | tcatcttctt | cttgtgtttc | 780 |
| ctgccctatc | acacactgag | gaccgtccac | ttgacgacat | ggaaagtggg | tttatgcaaa | 840 |
| gacagactgc | ataaagcttt | ggttatcaca | ctggccttgg | cagcagccaa | tgcctgcttc | 900 |
| aatcctctgc | tctattactt | tgctggggag | aattttaagg | acagactaaa | gtctgcactc | 960 . |
| agaaaaggcc | atccacagaa | ggcaaagaca | aagtgtgttt | tccctgttag | tgtgtggttg | 1020 |
| agaaaggaaa | caagagtata | a | | | | 1041 |
| | | | | | | |

<210> 88 <211> 346

<212> PRT

<213> Homo sapiens

<400> 88

Met Glu Arg Lys Phe Met Ser Leu Gln Pro Ser Ile Ser Val Ser Glu $1 \ \ \,$ 5

Met Glu Pro Asn Gly Thr Phe Ser Asn Asn Asn Ser Arg Asn Cys Thr 20 25 30

Ile Glu Asn Phe Lys Arg Glu Phe Phe Pro Ile Val Tyr Leu Ile Ile 35 40 45

Phe Phe Trp Gly Val Leu Gly Asn Gly Leu Ser Ile Tyr Val Phe Leu 50

Gln Pro Tyr Lys Lys Ser Thr Ser Val Asn Val Phe Met Leu Asn Leu 65 70 75 80

Ala Ile Ser Asp Leu Leu Phe Ile Ser Thr Leu Pro Phe Arg Ala Asp 85 90 95

Tyr Tyr Leu Arg Gly Ser Asn Trp Ile Phe Gly Asp Leu Ala Cys Arg 100 105 110

Ile Met Ser Tyr Ser Leu Tyr Val Asn Met Tyr Ser Ser Ile Tyr Phe
115 120 125

Leu Thr Val Leu Ser Val Val Arg Phe Leu Ala Met Val His Pro Phe 130 135 140

| Arg 145 | Leu | Leu | His | Val | Thr 150 | Ser | Ile | Arg | Ser | Ala 155 | Trp | Ile | Leu | Cys | Gly 160 | |
|--------------------------|--------------------------|-------------------------|---------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|----|
| Ile | Ile | Trp | Ile | Leu 165 | Ile | Met | Ala | Ser | Ser 170 | Ile | Met | Leu | Leu | Asp 175 | Ser | |
| Gly | Ser | Glu | Gln 180 | Asn | Gly | Ser | Val | Thr 185 | Ser | Cys | Leu | Glu | Leu 190 | Asn | Leu | |
| Tyr | Lys | Ile 195 | Ala | Lys | Leu | Gln | Thr 200 | Met | Asn | Tyr | Ile | Ala 205 | Leu | Val | Val | |
| Gly | Cys 210 | Leu | Leu | Pro | Phe | Phe 215 | Thr | Leu | Ser | Ile | Cys 220 | Tyr | Leu | Leu | Ile | |
| 11e 225 | Arg | Val | Leu | Leu | Lys 230 | Val | Glu | Val | Pro | Glu 235 | Ser | Gly | Leu | Arg | Val 240 | |
| Ser | His | Arg | Lys | Ala 245 | Lys | Thr | Thr | Ile | 11e 250 | Ile | Thr | Leu | Ile | Ile 255 | Phe | |
| Phe | Leu | Cys | Phe 260 | Leu | Pro | Tyr | His | Thr 265 | Leu | Arg | Thr | Val | His 270 | Leu | Thr | |
| Thr | Trp | Lys 275 | | Gly | Leu | Cys | Lys 280 | Asp | Arg | Leu | His | Lys 285 | Ala | Leu | Val | |
| Ile | Thr 290 | | Ala | Leu | Ala | Ala 295 | Ala | Asn | Ala | Суѕ | Phe 300 | Asn | Pro | Leu | Leu | |
| Tyr 305 | T yr | Phe | Ala | Gly | Glu 310 | Asn | Phe | Lys | Asp | Arg 315 | Leu | Lys | Ser | Ala | Leu 320 | |
| Arg | Lys | Gly | His | Pro 325 | | Lys | Ala | Lys | Thr 330 | Lys | Cys | Val | Phe | Pro 335 | Val | |
| Ser | Val | Trp | Leu 340 | | Lys | Glu | Thr | Arg 345 | Val | | | | | | | |
| <21 <21 <21 <21 | 1> 2> | 89 28 DNA Arti | fici | al S | Seque | nce | | | | | | | | | | |
| <22 <22 <22 | 1> | | el Se | | | | | | | - | | | J | | | |
| <40 cca | 0> .gtgc | 89 caaa | gcta | agaa | aag t | gato | ttc | | | | | | | | | 28 |
| <21 | .0> .1> .2> .3> | 90 28 DNA Art: | ific | ial : | Seque | ence | | | | | | | | | | |
| <22 | 20> 21> 23> | | c_fea el S | | | | | | | | | | | | | |
| | 00> | 90 cact | ttc | ttag | ctt · | tgca | ctgg | | | | | | | | | 28 |

| <211> 1527 <212> DNA | |
|--|------|
| <212> DNA <213> Homo sapiens | |
| | |
| <400> 91 atgacgteca cetgeaceaa cageaegege gagagtaaca geageeacae gtgeatgeee | 60 |
| ctctccaaaa tgcccatcag cctggcccac ggcatcatcc gctcaaccgt gctggttatc | 120 |
| tteetegeeg cetetttegt eggeaacata gtgetggege tagtgttgea gegeaageeg | 180 |
| cagetgetge aggtgaceaa cegttttate tttaacetee tegteacega cetgetgeag | 240 |
| atttegeteg tggeceeetg ggtggtggee acetetgtge etetettetg geceeteaac | 300 |
| agccacttet geaeggeest ggttageete acceaectgt tegeettege cagegteaae | 360 |
| accattgtcg tggtgtcagt ggatcgctac ttgtccatca tccaccctct ctcctacccg | 420 |
| tocaagatga cocagogoog oggttacotg otoototatg goacotggat tgtggooato | 480 |
| ctgcagagca ctcctccact ctacggctgg ggccaggctg cctttgatga gcgcaatgct | 540 |
| ctctgctcca tgatctgggg ggccagcccc agctacacta ttctcagcgt ggtgtccttc | 600 |
| atcgtcattc cactgattgt catgattgcc tgctactccg tggtgttctg tgcagcccgg | 660 |
| aggcagcatg ctctgctgta caatgtcaag agacacagct tggaagtgcg agtcaaggac | 720 |
| tgtgtggaga atgaggatga agagggagca gagaagaagg aggagttcca ggatgagagt | 780 |
| gagtttcgcc gccagcatga aggtgaggtc aaggccaagg agggcagaat ggaagccaag | 840 |
| gacggcagcc tgaaggccaa ggaaggaagc acggggacca gtgagagtag tgtagaggcc | 900 |
| aggggcagcg aggaggtcag agagagcagc acggtggcca gcgccggcag catggagggt | 960 |
| aaggaaggca gcaccaaagt tgaggagaac agcatgaagg cagacaaggg tcgcacagag | 1020 |
| gtcaaccagt gcagcattga cttgggtgaa gatgacatgg agtttggtga agacgacatc | 1080 |
| aatttcagtg aggatgacgt cgaggcagtg aacatcccgg agagcctccc acccagtcgt | 1140 |
| cgtaacagca acagcaacce tectetgeee aggtgetace agtgeaaage taagaaagtg | 1200 |
| atottoatoa toattttoto otatgtgota toootggggo ootactgott tttagcagto | 1260 |
| ctggccgtgt gggtggatgt cgaaacccag gtaccccagt gggtgatcac cataatcatc | 1320 |
| tggcttttct tcctgcagtg ctgcatccac ccctatgtct atggctacat gcacaagacc | 1380 |
| attaagaagg aaatccagga catgctgaag aagttcttct gcaaggaaaa gcccccgaaa | 1440 |
| gaagatagcc acccagacct gcccggaaca gagggtggga ctgaaggcaa gattgtccct | 1500 |
| tectacgatt etgetaettt teettga | 1527 |

<210> 92 <211> 508 <212> PRT <213> Homo sapiens

<210> 91

Page 54

<400> 92

Met Thr Ser Thr Cys Thr Asn Ser Thr Arg Glu Ser Asn Ser Ser His 10 Thr Cys Met Pro 20 Leu Ser Lys Met Pro 25 Leu Ala His Gly Ile Ser Leu Arg Ser Thr Val Leu Val Ile Phe Leu Ala Ala Ser Phe Val Gly And And Arg Ser Phe Val Gly Arg Ser Thr Val Leu Val Ile Phe Leu Ala Ala Ser Phe Val Gly Arg Ser P

Asn Ile Val Leu Ala Leu Val Leu Gln Arg Lys Pro Gln Leu Leu Gln

Val Thr Asn Arg Phe Ile Phe Asn Leu Leu Val Thr Asp Leu Leu Gln 65 70 75 80

Ile Ser Leu Val Ala Pro Trp Val Val Ala Thr Ser Val Pro Leu Phe 85 90 95

Trp Pro Leu Asn Ser His Phe Cys Thr Ala Leu Val Ser Leu Thr His 100 105 110

Leu Phe Ala Phe Ala Ser Val Asn Thr Ile Val Val Ser Val Asp 115 . 120 . 125

Arg Tyr Leu Ser Ile Ile His Pro Leu Ser Tyr Pro Ser Lys Met Thr 130 135 140

Gln Arg Arg Gly Tyr Leu Leu Leu Tyr Gly Thr Trp Ile Val Ala Ile 145 150 155 160

Leu Gln Ser Thr Pro Pro Leu Tyr Gly Trp Gly Gln Ala Ala Phe Asp 165 170 175

Glu Arg Asn Ala Leu Cys Ser Met Ile Trp Gly Ala Ser Pro Ser Tyr 180 185 190

Thr Ile Leu Ser Val Val Ser Phe Ile Val Ile Pro Leu Ile Val Met 195 200 205

Ile Ala Cys Tyr Ser Val Val Phe Cys Ala Ala Arg Arg Gln His Ala 210 215 220

Leu Leu Tyr Asn Val Lys Arg His Ser Leu Glu Val Arg Val Lys Asp 225 230 235 240

Cys Val Glu Asn Glu Asp Glu Glu Gly Ala Glu Lys Lys Glu Glu Phe 245 250 255

Gln Asp Glu Ser Glu Phe Arg Arg Gln His Glu Gly Glu Val Lys Ala 260 265 270

Lys Glu Gly Arg Met Glu Ala Lys Asp Gly Ser Leu Lys Ala Lys Glu 275 280 285

Gly Ser Thr Gly Thr Ser Glu Ser Ser Val Glu Ala Arg Gly Ser Glu 290 295 300

Glu Val Arg Glu Ser Ser Thr Val Ala Ser Asp Gly Ser Met Glu Gly 305 310 315 320

Lys Glu Gly Ser Thr Lys Val Glu Glu Asn Ser Met Lys Ala Asp Lys 325 330 335

Gly Arg Thr Glu Val Asn Gln Cys Ser Ile Asp Leu Gly Glu Asp Asp Met Glu Phe Gly Glu Asp Asp Ile Asn Phe Ser Glu Asp Asp Val Glu 365 Ala Val Asn Ile Pro Glu Ser Leu Pro Pro Ser Arg Arg Asn Ser Asn 375 Ser Asn Pro Pro Leu Pro Arg Cys Tyr Gln Cys Lys Ala Lys Lys Val Ile Phe Ile Ile Phe Ser Tyr Val Leu Ser Leu Gly Pro Tyr Cys Phe Leu Ala Val Leu Ala Val Trp Val Asp Val Glu Thr Gln Val Pro 425 Gln Trp Val Ile Thr Ile Ile Ile Trp Leu Phe Phe Leu Gln Cys Cys Ile His Pro Tyr Val Tyr Gly Tyr Met His Lys Thr Ile Lys Lys Glu Ile Gln Asp Met Leu Lys Lys Phe Phe Cys Lys Glu Lys Pro Pro Lys 470 475 Glu Asp Ser His Pro Asp Leu Pro Gly Thr Glu Gly Gly Thr Glu Gly 485 Lys Ile Val Pro Ser Tyr Asp Ser Ala Thr Phe Pro <210> 93 <211> 29 <212> DNA <213> Artificial Sequence <220> <221> misc_feature <223> Novel Sequence <400> 93 gccgccaccg cgccaagagg aagattggc 29 <210> 94 <211> 29 <212> DNA <213> Artificial Sequence <220> <221> misc_feature <223> Novel Sequence

29

<210> 95 <211> 1092 <212> DNA

<400> 94

gccaatcttc ctcttggcgc ggtggcggc

<213> Homo sapiens

| <400> 95 atgggccccg | gcgaggcgct | gctggcgggt | ctcctggtga | tggtactggc | cgtggcgctg | 60 |
|------------------------|--------------|------------|------------|------------|--------------|------|
| ctatccaacg | cactggtgct | gctttgttgc | gcctacagcg | ctgagctccg | cactcgagcc | 120 |
| tcaggcgtcc | tcctggtgaa | tctgtcgctg | ggccacctgc | tgctggcggc | gctggacatg | 180 |
| cccttcacgc | tgctcggtgt | gatgcgcggg | cggacaccgt | cggcgcccgg | cgcatgccaa | 240 |
| gtcattggct | tcctggacac | cttcctggcg | tccaacgcgg | cgctgagcgt | ggcggcgctg | 300 |
| agcgcagacc | agtggctggc | agtgggcttc | ccactgcgct | acgccggacg | cctgcgaccg | 360 |
| | gcctgctgct | | | | | 420 |
| | gctcgtggct | | | | | 480 |
| | ctgagcgtcc | | | | | 540 |
| | cgctggcggt | | | | | 600 |
| | agcgcatgga | | | | | 660 |
| | tgcggcagcg | • | | | | 720 |
| | gcattgctat | | | | | 780 |
| | agctcgtgcc | | | | | 840 |
| | acagcaaggo | | | | | 900 |
| | tcctggccgg | | | | | 960 |
| | , acagctctct | | | | | 1020 |
| | | | | | : ctgcctgcag | 1080 |
| cagacacact | | | | | | 1092 |
| - | | | | | | |

<210> 96

<211> 363 <212> PRT <213> Homo sapiens

<400> 96

Met Gly Pro Gly Glu Ala Leu Leu Ala Gly Leu Leu Val Met Val Leu

Ala Val Ala Leu Leu Ser Asn Ala Leu Val Leu Cys Cys Ala Tyr

Ser Ala Glu Leu Arg Thr Arg Ala Ser Gly Val Leu Leu Val Asn Leu

Ser Leu Gly His Leu Leu Leu Ala Ala Leu Asp Met Pro Phe Thr Leu

Leu Gly Val Met Arg Gly Arg Thr Pro Ser Ala Pro Gly Ala Cys Gln 65 70 75 80

Val Ile Gly Phe Leu Asp Thr Phe Leu Ala Ser Asn Ala Ala Leu Ser Page 57

90 95

Val Ala Ala Leu Ser Ala Asp Gln Trp Leu Ala Val Gly Phe Pro Leu 100 105 110

Arg Tyr Ala Gly Arg Leu Arg Pro Arg Tyr Ala Gly Leu Leu Gly 115 120 125

Cys Ala Trp Gly Gln Ser Leu Ala Phe Ser Gly Ala Ala Leu Gly Cys 130 135 140

Ser Trp Leu Gly Tyr Ser Ser Ala Phe Ala Ser Cys Ser Leu Arg Leu 145 150 155 160

Pro Pro Glu Pro Glu Arg Pro Arg Phe Ala Ala Phe Thr Ala Thr Leu 165 170 175

His Ala Val Gly Phe Val Leu Pro Leu Ala Val Leu Cys Leu Thr Ser 180 185 190

Leu Gln Val His Arg Val Ala Arg Ser His Cys Gln Arg Met Asp Thr 195 200 205

Val Thr Met Lys Ala Leu Ala Leu Leu Ala Asp Leu His Pro Ser Val 210 215 220

Arg Gln Arg Cys Leu Ile Gln Gln Lys Arg Arg Arg His Arg Ala Thr 225 235 240

Arg Lys Ile Gly Ile Ala Ile Ala Thr Phe Leu Ile Cys Phe Ala Pro $245 \hspace{1.5cm} 250 \hspace{1.5cm} 255 \hspace{1.5cm}$

Ala Gln Lys Gly Ile Leu Ser Lys Cys Leu Thr Tyr Ser Lys Ala Val 275 280 285

Ala Asp Pro Phe Thr Tyr Ser Leu Leu Arg Arg Pro Phe Arg Gln Val 290 295 300

Leu Ala Gly Met Val His Arg Leu Leu Lys Arg Thr Pro Arg Pro Ala 305 310 315 320

Ser Thr His Asp Ser Ser Leu Asp Val Ala Gly Met Val His Gln Leu 325 330 335

Leu Lys Arg Thr Pro Arg Pro Ala Ser Thr His Asn Gly Ser Val Asp 340 345 350

Thr Glu Asn Asp Ser Cys Leu Gln Gln Thr His 355

<210> 97

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<221> misc feature

<223> Novel Sequence

<400> 97 gatetetaga atggagteet cacceateee eeag

34

| <210> <211> <212> <213> | 98 36 DNA Artif | icial | Seque | ence | | | | |
|----------------------------------|---------------------------|--------------------|--------|------------|------------|------------|------------|------|
| <220> <221> <223> | | _featuı [Seque | | | | | | |
| <400> gatcgat | 98 tatc (| egtgact | tcca (| gccggggtga | ggcggc | | | 36 |
| <210> <211> <212> <213> | 99 2610 DNA Homo | sapie | ns an | d Rat | | | | |
| <400> atggag | 99 tcct (| caccca | tccc | ccagtcatca | gggaactctt | ccactttggg | gagggtccct | 60 |
| | | | | | | tggggctacg | | 120 |
| | | | | | | tgactgctgt | | 180 |
| | | | | | | gaaaatttgt | | 240 |
| | | | | | | tgcccctggc | | 300 |
| | | | | | | cctgccgcct | | 360 |
| | | | | | | cagccatcaa | | 420 |
| | | | | | | tgacgctggg | | 480 |
| tctgtg | ctgg | tgggtg | tgtg | ggtgaaggcc | ttggccatgg | cttctgtgcc | agtgttggga | 540 |
| agggto | tcct | gggagg | aagg | agctcccagt | gtccccccag | gctgttcact | ccagtggagc | 600 |
| cacagt | gcct | actgcc | agct | ttttgtggtg | gtctttgctg | tcctttactt | tctgttgccc | 660 |
| ctgctc | ctca | tacttg | tggt | ctactgcagc | atgttccgag | tggcccgcgt | ggctgccatg | 720 |
| cagcac | gggc | cgctgc | ccac | gtggatggag | acaccccggc | aacgctccga | atctctcagc | 780 |
| | | | | | | ccaccccaca | | 840 |
| | | | | | | | gctctgttgg | 900 |
| | | | | | | | ttcaactggg | 960 |
| | | | | | | | ccctttcttc | 1020 |
| | | | | | | | ctgcttcttc | 1080 |
| | | | | | | | tgaggagaac | 1140 |
| | | | | | | | ccgaccccta | 1200 |
| | | | | | | | gatagctgag | 1260 |
| | | | | | | | agacagctac | 1320 |
| 2-3-0 | | , | | | | Page 59 | | |

WO 01/36471 · ; PCT/US00/31509

| ctccgtcct | g ccgcctcac | c ccggctggaq | g tcagcgata | t ctgcagaat | t ccaccacact | 1380 |
|------------|--------------|--------------|--------------|-------------|--------------|-------|
| ggactagtg | g atccgagct | c ggtaccaago | ttgggctgca | ggtcgatgg | g ctgcctcggc | 1440 |
| aacagtaag | a ccgaggacc | a gcgcaacgag | gagaaggcgd | agcgcgagg | caacaaaaag | 1500 |
| atcgagaag | c agctgcaga | a ggacaagcag | gtctaccgg | ccacgcacco | g cctgctgctg | 1560 |
| ctgggtgct | g gagagtetge | g caaaagcacc | : attgtgaago | agatgaggat | cctacatgtt | 1620 |
| aatgggttta | a acggagagg | g cggcgaagag | gacccgcagg | ctgcaaggag | g caacagegat | 1680 |
| ggtgagaag | j ccaccaaagt | gcaggacatc | aaaaacaacc | tgaaggaggo | : cattgaaacc | 1740 |
| | | | | | tgagaaccag | 1800 |
| ttcagagtgg | actacattct | gagcgtgatg | aacgtgccaa | actttgactt | cccacctgaa | 1860 |
| | | | | | ctacgagcgc | 1,920 |
| | | cgactgtgcc | | | | 1980 |
| | | aagtgaccag | | | | 2040 |
| | | ccaggtggac | | | | 2100 |
| | | caagtggatc | | | | 2160 |
| | | ctacaacatg | | | | 2220 |
| | | cttcaagagc | | | | 2280 |
| | | caagcaagat | | | | 2340 |
| | | tccagagttc | | | | 2400 |
| | | acgcgtgacc | | | | 2460 |
| | | tggagatgga | | | | 2520 |
| | | ccgccgtgtc | | | | 2580 |
| | gccaatacga | | | | J J- | 2610 |
| | | | | | | |

<210> 100 <211> 869

<212> PRT

<213> Homo sapiens and Rat

<400> 100

Met Glu Ser Ser Pro Ile Pro Gln Ser Ser Gly Asn Ser Ser Thr Leu $1 \hspace{1cm} 1 \hspace{1cm}$

Gly Arg Val Pro Gln Thr Pro Gly Pro Ser Thr Ala Ser Gly Val Pro 20 25 30

Glu Val Gly Leu Arg Asp Val Ala Ser Glu Ser Val Ala Leu Phe Phe 35 40 45

Met Leu Leu Asp Leu Thr Ala Val Ala Gly Asn Ala Ala Val Met

Page 60

| 65 | | | | | Thr 70 | | | | | 15 | | | | | 80 |
|------------|------------|------------|------------|------------|--------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|--------------|
| His | Leu | Cys | Leu | Val 85 | Asp | Leu | Leu | Ala | Ala 90 | Leu | Thr | Leu | Met | Pro 95 | Leu |
| Ala | Met | Leu | Ser 100 | Ser | Ser | Ala | Leu | Phe 105 | Asp | His | Ala | Leu | Phe 110 | Gly | Glu |
| Val | Ala | Cys 115 | Arg | Leu | Tyr | Leu | Phe 120 | Leu | Ser | Val | Cys | Phe 125 | Val | Ser | Leu |
| Ala | Ile 130 | Leu | Ser | Val | Ser | Ala 135 | Ile | Asn | Val | Glu | Arg 140 | Tyr | Tyr | Tyr | Val |
| Val 145 | His | Pro | Met | Arg | Tyr 150 | Glu | Val | Arg | Met | Thr 155 | Leu | Gly | Leu | Val | Ala 160 |
| | | | | 165 | Val | | | | 1/0 | | | | | 1/3 | |
| Pro | Val | Leu | Gly 180 | Arg | Val | Ser | Trp | Glu 185 | Glu | Gly | Ala | Pro | Ser 190 | Val | Pro |
| Pro | Gly | Cys 195 | | Leu | Gln | Trp | Ser 200 | His | Ser | Ala | Tyr | Cys 205 | Gln | Leu | Phe |
| | 210 | | | | Val | 215 | | | | | 220 | | | | |
| 225 | | | | | Ser 230 | | | | | 235 | | | | | 240 |
| Gln | His | Gly | Pro | Leu 245 | Pro | Thr | Trp | Met | Glu 250 | Thr | Pro | Arg | Gln | Arg 255 | Ser |
| Glu | Ser | Leu | Ser 260 | | Arg | Ser | Thr | Met 265 | Val | Thr | Ser | Ser | Gly 270 | Ala | Pro |
| Gln | Thr | Thr 275 | | His | Arg | Thr | Phe 280 | Gly | Gly | Gly | Lys | Ala 285 | Ala | Val | Val |
| | 290 |) | | | Gly | 295 | | | | | 300 | | J | | |
| 305 | 5 | | | | 310 | | | | | 313 | | | | | 320 |
| | | | | 325 | • | | | | 330 |) | | | | 335 | |
| | | | 34 | 0 | | | | 345 |) | | | | 330 | , | ı Leu |
| | | 35 | 5 | | | | 360 | , | | | | 30. | , | | ı Leu |
| | 37 | 0 | | | | 375 |) | | | | 361 | , | | | n Phe |
| 38 | 5 | | | | 390 | נ | | | | 395 | , | | | | o Leu 400 |
| Pr | | | _ | | - C1. | , Dr | o Dra | . 11 | s Va | 3 Asr | n Ph | a Arc | r Il | e Pr | o Gly |

-

Gln Ile Ala Glu Glu Thr Ser Glu Phe Leu Glu Gln Gln Leu Thr Ser 420 425

Asp Ile Ile Met Ser Asp Ser Tyr Leu Arg Pro Ala Ala Ser Pro Arg 435 440 445

Leu Glu Ser Ala Ile Ser Ala Glu Phe His His Thr Gly Leu Val Asp
450 460

Pro Ser Ser Val Pro Ser Leu Gly Cys Arg Ser Met Gly Cys Leu Gly 465 470 475 480

Asn Ser Lys Thr Glu Asp Gln Arg Asn Glu Glu Lys Ala Gln Arg Glu 485 490 495

Ala Asn Lys Lys Ile Glu Lys Gln Leu Gln Lys Asp Lys Gln Val Tyr 500 505 510

Arg Ala Thr His Arg Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys 515 520 525

Ser Thr Ile Val Lys Gln Met Arg Ile Leu His Val Asn Gly Phe Asn 530 540

Gly Glu Gly Glu Glu Asp Pro Gln Ala Ala Arg Ser Asn Ser Asp 545 550 555 560

Gly Glu Lys Ala Thr Lys Val Gln Asp Ile Lys Asn Asn Leu Lys Glu 565 570 575

Ala Ile Glu Thr Ile Val Ala Ala Met Ser Asn Leu Val Pro Pro Val 580 585 585

Glu Leu Ala Asn Pro Glu Asn Gln Phe Arg Val Asp Tyr Ile Leu Ser 595 600 605

Val Met Asn Val Pro Asn Phe Asp Phe Pro Pro Glu Phe Tyr Glu His 610 615 620

Ala Lys Ala Leu Trp Glu Asp Glu Gly Val Arg Ala Cys Tyr Glu Arg 625 630 635 640

Ser Asn Glu Tyr Gln Leu Ile Asp Cys Ala Gln Tyr Phe Leu Asp Lys 645 650 655

Ile Asp Val Ile Lys Gln Ala Asp Tyr Val Pro Ser Asp Gln Asp Leu 660 665 670

Leu Arg Cys Arg Val Leu Thr Ser Gly Ile Phe Glu Thr Lys Phe Gln 675 680 685

Val Asp Lys Val Asn Phe His Met Phe Asp Val Gly Gln Arg Asp 690 695 700

Glu Arg Arg Lys Trp Ile Gln Cys Phe Asn Asp Val Thr Ala Ile Ile 705 710 715 720

Phe Val Val Ala Ser Ser Ser Tyr Asn Met Val Ile Arg Glu Asp Asn 725 730 735

Gln Thr Asn Arg Leu Gln Glu Ala Leu Asn Leu Phe Lys Ser Ile Trp
740 745 750

Asn Asn Arg Trp Leu Arg Thr Ile Ser Val Ile Leu Phe Leu Asn Lys Page 62 WO 01/36471 PCT/US00/31509

| | 755 | | | | | 760 | | | | | 765 | | | | |
|----------------------------------|--------------------------|----------------|---------------|------------|------------|-------|------|------------|------------|------------|------|------|------------|------------|-----|
| Gln Asp 770 | | Leu | Ala | Glu | Lys 775 | Val | Leu | Ala | Gly | Lys 780 | Ser | Lys | Ile | Glu | |
| Asp Tyr 785 | Phe | Pro | Glu | Phe 790 | Ala | Arg | Туr | Thr | Thr 795 | Pro | Glu | Asp | Ala | Thr 800 | |
| Pro Glu | Pro | Gly | Glu 805 | Asp | Pro | Arg | Val | Thr 810 | Arg | Ala | Lys | Tyr | Phe 815 | Ile | |
| Arg Asp | | 820 | | | | | 825 | | | | | 630 | | | |
| Tyr Cys | 835 | | | | | 840 | | | • | | 047 | | | | |
| Arg Val 850 | | Asn | Asp | Суѕ | Arg 855 | Asp | Ile | Ile | Gln | Arg 860 | Met | His | Leu | Arg | |
| Gln Tyr 865 | Glu | Leu | Leu | | | | | | | | | | | | |
| <210> <211> <212> <213> | 101 30 DNA Arti | fici | al S | eque | nce | | | | | | | | | | |
| <220> <221> <223> | | :_fea el Se | | | | | | | | | | | | | |
| <400> tctagaa | 101 atga | cgto | cacc | tg o | acca | acag | ıc | | | | | | | | 30 |
| <210> <211> <212> <213> | 102 34 DNA Art: | ifici | .al S | Seque | ence | | | | | | | | | | |
| <220> <221> <223> | mis Nov | c_fea el Se | ature eque | e nce | | | | | | | | v | | | |
| <400> gatato | 102 gcag | gaa | aagta | agc · | agaa | togta | ag g | aag | | | | | | | 34 |
| <210> <211> <212> <213> | 103 278 DNA Hom | 1 | pien | s an | d Ra | t | | | | | | | | | |
| <400> | 103 | cct | αcac | caa | cago | acqc | gc q | agag | taac | a gc | agco | acac | gtg | catgece | 60 |
| | | | | | | | | | | | | | | ggttatc | 120 |
| | | | | | | | | | | | | | | caagccg | 180 |
| | | | | | | | | | | | | | | gctgcag | 240 |
| | | | | | | | | | | | | | | | |

Page 63

| atttegeteg tggeceettg ggtggtggee acetetgtge etetettetg geceeteaae | 300 |
|--|------|
| agccacttct gcacggccct ggttagcctc acccacctgt tcgccttcgc cagcgtcaac | 360 |
| accattgtcg tggtgtcagt ggatcgctac ttgtccatca tccaccctct ctcctacccg. | 420 |
| tccaagatga cccagcgccg cggttacctg ctcctctatg gcacctggat tgtggccatc | 480 |
| ctgcagagca ctcctccact ctacggctgg ggccaggctg cctttgatga gcgcaatgct | 540 |
| ctctgctcca tgatctgggg ggccagcccc agctacacta ttctcagcgt ggtgtccttc | 600 |
| atcgtcattc cactgattgt catgattgcc tgctactccg tggtgttctg tgcagcccgg | 660 |
| aggcagcatg ctctgctgta caatg@caag agacacagct tggaagtgcg agtcaaggac | 720 |
| tgtgtggaga atgaggatga agagggagca gagaagaagg aggagttcca ggatgagagt | 780 |
| gagtttcgcc gccagcatga aggtgaggtc aaggccaagg agggcagaat ggaagccaag | 840 |
| gacggcagcc tgaaggccaa ggaaggaagc acggggacca gtgagagtag tgtagaggcc | 900 |
| aggggcagcg aggaggtcag agagagcagc acggtggcca gcgacggcag catggagggt | 960 |
| aaggaaggca gcaccaaagt tgaggagaac agcatgaagg cagacaaggg tcgcacagag | 1020 |
| gtcaaccagt gcagcattga cttgggtgaa gatgacatgg agtttggtga agacgacatc | 1080 |
| aatttcagtg aggatgacgt cgaggcagtg aacatcccgg agagcctccc acccagtcgt | 1140 |
| cgtaacagca acagcaaccc tectetgeec aggtgetace agtgeaaage tgetaaagtg | 1200 |
| atetteatea teatitiete etatgtgeta tecetgggge eetactgeti titageagte | 1260 |
| ctggccgtgt gggtggatgt cgaaacccag gtaccccagt gggtgatcac cataatcatc | 1320 |
| tggcttttct tcctgcagtg ctgcatccac ccctatgtct atggctacat gcacaagacc | 1380 |
| attaagaagg aaatccagga catgctgaag aagttcttct gcaaggaaaa gcccccgaaa | 1440 |
| gaagatagcc acccagacct gcccggaaca gagggtggga ctgaaggcaa gattgtccct | 1500 |
| tectaegatt etgetaettt teetgegata tetgeagaat teeaceaeae tggaetagtg | 1560 |
| gatccgaget eggtaccaag ettgggetge aggtegatgg getgeetegg caacagtaag | 1620 |
| accgaggacc agcgcaacga ggagaaggcg cagcgcgagg ccaacaaaaa gatcgagaag | 1680 |
| cagctgcaga aggacaagca ggtctaccgg gccacgcacc gcctgctgct gctgggtgct | 1740 |
| ggagagtctg gcaaaagcac cattgtgaag cagatgagga tcctacatgt taatgggttt | 1800 |
| aacggagagg gcggcgaaga ggacccgcag gctgcaagga gcaacagcga tggtgagaag | 1860 |
| gccaccaaag tgcaggacat caaaaacaac ctgaaggagg ccattgaaac cattgtggcc | 1920 |
| gccatgagca acctggtgcc ccccgtggag ctggccaacc ctgagaacca gttcagagtg | 1980 |
| gactacatte tgagegtgat gaaegtgeea aaetttgaet teecaeetga attetatgag | 2040 |
| catgccaagg ctctgtggga ggatgaggga gttcgtgcct gctacgagcg ctccaacgag | 2100 |
| taccagctga tegactgtge ccagtaette etggacaaga ttgatgtgat caagcaggee | 2160 |
| gactacgtgc caagtgacca ggacctgctt cgctgccgcg tcctgacctc tggaatcttt | 2220 |
| Page 64 | |

| gagaccaagt | tccaggtgga | caaagtcaac | ttccacatgt | tcgatgtggg | cggccagcgc | 2280 |
|--|------------|------------|------------|------------|------------|------|
| gatgaacgcc | gcaagtggat | ccagtgcttc | aatgatgtga | ctgccatcat | cttcgtggtg | 2340 |
| gccagcagca | gctacaacat | ggtcatccgg | gaggacaacc | agaccaaccg | tctgcaggag | 2400 |
| gctctgaacc | tcttcaagag | catctggaac | aacagatggc | tgcgtaccat | ctctgtgatc | 2460 |
| ctcttcctca | acaagcaaga | tctgcttgct | gagaaggtcc | tcgctgggaa | atcgaagatt | 2520 |
| gaggactact | ttccagagtt | cgctcgctac | accactcctg | aggatgcgac | tcccgagccc | 2580 |
| ggagaggacc | cacgcgtgac | ccgggccaag | tacttcatcc | gggatgagtt | tctgagaatc | 2640 |
| agcactgcta | gtggagatgg | acgtcactac | tgctaccctc | actttacctg | cgccgtggac | 2700 |
| | | cttcaacgac | | | | 2760 |
| | agctgctcta | | | | | 2783 |
| _ | | | | | | |
| <210> 104 <211> 926 <212> PRT <213> Hom | | nd Rat | | | | |
| <400> 104 | | | | | | |

Met Thr Ser Thr Cys Thr Asn Ser Thr Arg Glu Ser Asn Ser Ser His $1 \hspace{1cm} 5 \hspace{1cm} 10 \hspace{1cm} 15$

Thr Cys Met Pro Leu Ser Lys Met Pro Ile Ser Leu Ala His Gly Ile 20 25 30

Ile Arg Ser Thr Val Leu Val Ile Phe Leu Ala Ala Ser Phe Val Gly

Asn Ile Val Leu Ala Leu Val Leu Gln Arg Lys Pro Gln Leu Leu Gln

Val Thr Asn Arg Phe Ile Phe Asn Leu Leu Val Thr Asp Leu Leu Gln 65 70 80

Ile Ser Leu Val Ala Pro Trp Val Val Ala Thr Ser Val Pro Leu Phe 85 90 95

Trp Pro Leu Asn Ser His Phe Cys Thr Ala Leu Val Ser Leu Thr His

Leu Phe Ala Phe Ala Ser Val Asn Thr Ile Val Val Ser Val Asp

Arg Tyr Leu Ser Ile Ile His Pro Leu Ser Tyr Pro Ser Lys Met Thr

Gln Arg Arg Gly Tyr Leu Leu Leu Tyr Gly Thr Trp Ile Val Ala Ile 145 150 155 160

Leu Gln Ser Thr Pro Pro Leu Tyr Gly Trp Gly Gln Ala Ala Phe Asp

Glu Arg Asn Ala Leu Cys Ser Met Ile Trp Gly Ala Ser Pro Ser Tyr

Page 65

Thr Ile Leu Ser Val Val Ser Phe Ile Val Ile Pro Leu Ile Val Met Ile Ala Cys Tyr Ser Val Val Phe Cys Ala Ala Arg Arg Gln His Ala Leu Leu Tyr Asn Val Lys Arg His Ser Leu Glu Val Arg Val Lys Asp Cys Val Glu Asn Glu Asp Glu Glu Gly Ala Glu Lys Lys Glu Glu Phe Gln Asp Glu Ser Glu Phe Arg Arg Gln His Glu Gly Glu Val Lys Ala Lys Glu Gly Arg Met Glu Ala Lys Asp Gly Ser Leu Lys Ala Lys Glu Gly Ser Thr Gly Thr Ser Glu Ser Ser Val Glu Ala Arg Gly Ser Glu Glu Val Arg Glu Ser Ser Thr Val Ala Ser Asp Gly Ser Met Glu Gly Lys Glu Gly Ser Thr Lys Val Glu Glu Asn Ser Met Lys Ala Asp Lys Gly Arg Thr Glu Val Asn Gln Cys Ser Ile Asp Leu Gly Glu Asp Asp Met Glu Phe Gly Glu Asp Asp Ile Asn Phe Ser Glu Asp Asp Val Glu 360 Ala Val Asn Ile Pro Glu Ser Leu Pro Pro Ser Arg Arg Asn Ser Asn Ser Asn Pro Pro Leu Pro Arg Cys Tyr Gln Cys Lys Ala Ala Lys Val Ile Phe Ile Ile Phe Ser Tyr Val Leu Ser Leu Gly Pro Tyr Cys Phe Leu Ala Val Leu Ala Val Trp Val Asp Val Glu Thr Gln Val Pro 425 Gln Trp Val Ile Thr Ile Ile Ile Trp Leu Phe Phe Leu Gln Cys Cys Ile Gln Asp Met Leu Lys Lys Phe Phe Cys Lys Glu Lys Pro Pro Lys Glu Asp Ser His Pro Asp Leu Pro Gly Thr Glu Gly Gly Thr Glu Gly 490 Lys Ile Val Pro Ser Tyr Asp Ser Ala Thr Phe Pro Ala Ile Ser Ala Glu Phe His His Thr Gly Leu Val Asp Pro Ser Ser Val Pro Ser Leu 520 Gly Cys Arg Ser Met Gly Cys Leu Gly Asn Ser Lys Thr Glu Asp Gln

540 Page 66 WO 01/36471 PCT/US00/31509

Arg Asn Glu Glu Lys Ala Gln Arg Glu Ala Asn Lys Lys Ile Glu Lys Gln Leu Gln Lys Asp Lys Gln Val Tyr Arg Ala Thr His Arg Leu Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys Ser Thr Ile Val Lys Gln Met 580 585 590 Arg Ile Leu His Val Asn Gly Phe Asn Gly Glu Gly Glu Glu Asp Pro Gln Ala Ala Arg Ser Asn Ser Asp Gly Glu Lys Ala Thr Lys Val 615 Gln Asp Ile Lys Asn Asn Leu Lys Glu Ala Ile Glu Thr Ile Val Ala Ala Met Ser Asn Leu Val Pro Pro Val Glu Leu Ala Asn Pro Glu Asn Gln Phe Arg Val Asp Tyr Ile Leu Ser Val Met Asn Val Pro Asn Phe Asp Phe Pro Pro Glu Phe Tyr Glu His Ala Lys Ala Leu Trp Glu Asp Glu Gly Val Arg Ala Cys Tyr Glu Arg Ser Asn Glu Tyr Gln Leu Ile 690 700 695 Asp Cys Ala Gln Tyr Phe Leu Asp Lys Ile Asp Val Ile Lys Gln Ala 705 710 715 720 Asp Tyr Val Pro Ser Asp Gln Asp Leu Leu Arg Cys Arg Val Leu Thr Ser Gly Ile Phe Glu Thr Lys Phe Gln Val Asp Lys Val Asn Phe His Met Phe Asp Val Gly Gly Gln Arg Asp Glu Arg Arg Lys Trp Ile Gln Cys Phe Asn Asp Val Thr Ala Ile Ile Phe Val Val Ala Ser Ser Ser 770 780 . Tyr Asn Met Val Ile Arg Glu Asp Asn Gln Thr Asn Arg Leu Gln Glu 785 790 795 800 Ala Leu Asn Leu Phe Lys Ser Ile Trp Asn Asn Arg Trp Leu Arg Thr Ile Ser Val Ile Leu Phe Leu Asn Lys Gln Asp Leu Leu Ala Glu Lys Val Leu Ala Gly Lys Ser Lys Ile Glu Asp Tyr Phe Pro Glu Phe Ala 840 Arg Tyr Thr Thr Pro Glu Asp Ala Thr Pro Glu Pro Gly Glu Asp Pro Arg Val Thr Arg Ala Lys Tyr Phe Ile Arg Asp Glu Phe Leu Arg Ile 875 Ser Thr Ala Ser Gly Asp Gly Arg His Tyr Cys Tyr Pro His Phe Thr

| | | | 885 | | | | | 890 | | | | | 895 | | |
|----------------------------------|----------------------------|------------|-------|-------|-----|------------|------------|-----|-----|-----|------------|------------|-----|-----|----|
| Cys A | la Val | Asp 900 | Thr | Glu | Asn | Ile | Arg 905 | Arg | Val | Phe | Asn | Asp 910 | Cys | Arg | |
| Asp I | le Ile 915 | Gln | Arg | Met | His | Leu 920 | Arg | Gln | Tyr | Glu | Leu 925 | Leu | | | |
| <210><211><211><212><213> | 23 DNA | Fi o i - | .1 5 | | | | | | | | | | | | |
| <220> <221> <223> | misc_ | feat | ure | | ice | | | | | | | | | | |
| <400> catgta | 105 atgcc a | gcgt | cctg | ıc to | С | | | | | | | | | | 23 |
| <210> | 106 | | | | | | | | | | | | | | |
| <211> <212> <213> | 24 DNA Artif | ícia | l Se | quen | ce | | | | | | | | | | |
| <220> <221> <223> | misc_ Novel | | | e | | | | | | | | | | | |
| <400> gctatg | 106 scctg aa | agcca | agtc | t tg | tg | | | | | | | | | | 24 |
| <210> <211> <212> <213> | 107 25 DNA Artifi | icial | L Sed | quenc | ce | | | | | | | | | | |
| <220> <221> <223> | misc_f Novel | | | • | | | | | | | | • | | | |
| <400> gcacct | 107 gctc ct | gago | acct | tct: | cc | | | | | | | | | | 25 |
| <210><211><211><212><213> | 108 26 DNA Artifi | cial | Sea | reunc | e | | | | | | | | | | |
| <220> <221> <223> | misc_f Novel | eatu | re | | - | | | | | | | | | | |
| <400> cacageo | 108 gctg ca | gccc | tgca | gct | ggc | | | | | | | | | | 26 |
| | | | | | | | | | | | | | | | |

Page 68

<210> 109

| <211> | 24 | |
|----------------|-----------------------------|-----|
| <212> | | |
| <213> | Artificial Sequence | |
| .0005 | | |
| <220> | misc_feature | |
| <223> | Novel Sequence | |
| | • | |
| | | |
| <400> | 109 atga ctctgtccag cctg | 24 |
| ccageg | acya ococycoody oray | |
| | | |
| <210> | 110 | |
| <2!1> <212> | 24 DNA | |
| <213> | Artificial Sequence | |
| | | |
| <220> | mica forture | |
| <221> <223> | _ | |
| 12237 | 110102 0042-1101 | |
| | | |
| <400> | 110 | 24 |
| cagaca | cttg gcagggacga ggtg | |
| | | |
| <210> | 111 | |
| <211> | | |
| <212> <213> | DNA Artficial Sequence | |
| \213/ | With Total Codesing | |
| <220> | | |
| | misc_feature | |
| <223> | Novel Sequence | |
| | | |
| <400> | 111 | 26 |
| cttgtg | gtct actgcagcat gttccg | 20 |
| | | • |
| <210> | 112 | |
| <211> | | |
| <212> <213> | | |
| (213/ | Altilitial bequesso | |
| <220> | | |
| <221> | | |
| <223> | Novel Sequence | |
| | | |
| <400> | 112 | 25 |
| catato | ecete egagtgteea gegge | 2.3 |
| | | |
| <210> | 113 | |
| <211> | | |
| <212> <213> | | |
| \Z13> | Withing addresses | |
| <220> | | |
| <221> | | |
| <223> | Nove Sequence | |

| <400 | | |
|----------------|--|-----|
| atgg | atcctt atcatggctt cctc | 24 |
| <210 | > 114 | |
| <211 | | |
| | > DNA | |
| <213 | > Artificial Sequence | |
| <220 | | |
| <2213 | <pre> misc_feature > Novel Sequence</pre> | |
| ~~~ | Nover Sequence | |
| <400> | - 114 | |
| | acagg teteatetaa gagetee | 0.7 |
| | | 27 |
| <210> | 115 | |
| <211> | | |
| <212> <213> | DNA Artificial Sequence | |
| | | |
| <220> | misc_feature | |
| <223> | Novel Sequence | |
| | • | |
| <400> | 115 | |
| ctctg | atgcc atctgctgga ttcctg | 26 |
| | | 20 |
| <210> | | |
| <211> <212> | | |
| | Artificial Sequence | |
| <220> | | |
| | misc_feature | |
| <223> | Novel Sequence | |
| | | |
| <400> | 116 | |
| gtagto | cact gaaagtccag tgatcc | 26 |
| | • | |
| <210> <211> | 117 24 | |
| <212> | DNA | |
| <213> | Artificial Sequence | |
| <220> | | |
| <221> | <u>-</u> | |
| <223> | Novel Sequence | |
| . 4 6 5 | | |
| <400> | 117 cgat ggccaacagc gctc | |
| | -gac ggooddoago golo | 24 |
| <210> | 118 | |
| <211> | 24 | |
| <212> | DNA | |
| <213> | Artificial Sequence | |

| <220> | | |
|----------------|--------------------------|----|
| | misc feature | |
| | Novel Sequence | |
| | | |
| <400× | 118 | |
| <400> | cctt agcgacagat gacc | 24 |
| guigege | cet agegaeague gase | |
| | | |
| <210> | 119 | |
| <211> <212> | 23 DNA | |
| <213> | Artificial Sequence | |
| | | |
| <220> | | |
| <221> | misc_feature | |
| <223> | Novel Sequence | |
| | | |
| <400> | 119 | 22 |
| tcaacc | tgta tagcagcatc ctc | 23 |
| | | |
| <210> | 120 | |
| <211> | | |
| <212> | | |
| <213> | Artificial Sequence | |
| <220> | | |
| | misc feature | |
| | Novel Sequence | |
| | | |
| <400> | 120 | |
| | tagc agaatggtta gcc | 23 |
| 95-5 | | |
| .010 | 101 | |
| <210> <211> | 121 24 | |
| <211> | | |
| <213> | Artificial Sequence | |
| | | |
| <220> <221> | misc feature | |
| | Novel Sequence | |
| 1220 | | |
| | | |
| <400> | 121 | 24 |
| gacacc | tgtc agcggtcgtg tgtg | |
| | | |
| <210> | 122 | |
| <211> | | |
| <212> <213> | DNA Artificial Sequence | |
| -210/ | | |
| <220> | | |
| <221> | | |
| <223> | Novel Sequence | |
| | | |
| <400> | | |
| ctgate | ggaag tagaggctgt ccatctc | 27 |

| <210><211><211><212><213> | 24 | |
|----------------------------------|---|----|
| | misc_feature Novel Sequence | |
| <400> gcgctg | 123 ragcg cagaccagtg gctg | 24 |
| <210> <211> <212> <213> | DNA | |
| <220> <221> <223> | misc_feature Novel Sequence | |
| <400> cacggt | 124 gacg aagggcacga gctc | 24 |
| <210> <211> <212> <213> | | |
| | misc_feature Novel Sequence | |
| <400> agccat | 125 ccct gccaggaagc atgg | 24 |
| <210><211><211><212><213> | 126 25 DNA Artificial Sequence | |
| <220> <221> <223> | misc_feature Novel Sequence | |
| <400> ccaggta | 126 aggt gtgcagcaca atggc | 25 |
| <210> <211> <212> <213> | 127 25 DNA Artificial Sequence | |
| <220> <221> <223> | misc_feature Novel Sequence | |

| VO 01/36471 | PCT/US00/31509 |
|-------------|----------------|
| | |

| <400> ctgttc | aaca gggctggttg gcaac | 25 |
|----------------------------------|--|----|
| <210><211><211><212><213> | | |
| <220> <221> <223> | misc_feature Novel Sequence | |
| <400> atcatg | ·128 tcta gactcatggt gatcc | 25 |
| <210> <211> <212> <213> | 129 6 PRT Artificial Sequence | |
| <220> <221> <223> | | |
| <400> | 129 | |
| Thr Le | u Glu Ser Ile Met 5 | |
| <210><211><211><212><213> | 130 5 PRT Artificial Sequence | |
| <221> <223> | misc_feature Novel Sequence | |
| <400> | 130 | |
| Glu Ty | r Asn Leu Val 5 | |
| <210> <211> <212> <213> | 131 5 PRT Artificial Sequence | |
| <220> <221> <223> | misc_feature Novel Sequence | |
| <400> | 131 | |
| Asp Cys | s Gly Leu Phe 5 | |
| <210> | 132 | |

```
<211> 36
<212> PRT
<213> Artificial Sequence
<220>
<221> misc feature
<223> Novel Sequence
<400> 132
Gly Ala Thr Cys Ala Ala Gly Cys Thr Thr Cys Cys Ala Thr Gly Gly
Cys Gly Thr Gly Cys Thr Gly Cys Cys Thr Gly Ala Gly Cys Gly Ala
Gly Gly Ala Gly
<210> 133
<211> 53
<212> PRT
<213> Artificial Sequence
<220>
<221> misc feature
<223> Novel Sequence
<400> 133
Cys Ala Gly Gly Cys Cys Gly Cys Ala Gly Thr Cys Cys Thr Thr Cys 20 25 30
Ala Gly Gly Thr Thr Cys Ala Gly Cys Thr Gly Cys Ala Gly Gly Ala
Thr Gly Gly Thr Gly
```

| | | • . |
|--|---|-----|
| | | • |
| | • | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | · | |
| | • | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 25 May 2001 (25.05.2001)

(10) International Publication Number WO 01/36471 A3

| C12N 15/12 | | |
|------------|--|--|
| | | |
| | | |

60/242,343 60/243.019

20 October 2000 (20,10,2000) 24 October 2000 (24.10.2000)

US US

- (21) International Application Number: PCT/US00/31509
- (22) International Filing Date:

(51) International Patent Classification7:

16 November 2000 (16.11.2000)

(25) Filing Language:

English

C07K 14/705.

(26) Publication Language:

English

| Priority Data: | | |
|----------------|--------------------------------|----|
| 880,661/06 | 17 November 1999 (17.11.1999) | US |
| 60/166,099 | 17 November 1999 (17.11.1999) | US |
| 60/166,369 | 17 November 1999 (17.11.1999) | US |
| 60/171,900 | 23 December 1999 (23.12.1999) | US |
| 60/171,901 | 23 December 1999 (23.12.1999) | US |
| 60/171,902 | 23 December 1999 (23.12.1999) | US |
| 60/181,749 | 11 February 2000 (11.02.2000) | US |
| 60/189,258 | 14 March 2000 (14.03.2000) | US |
| 60/189.259 | 14 March 2000 (14.03.2000) | US |
| 60/195,898 | 10 April 2000 (10.04.2000) | US |
| 60/195,899 | 10 April 2000 (10.04.2000) | US |
| 60/196.078 | 10 April 2000 (10.04.2000) | US |
| 60/200,419 | 28 April 2000 (28.04.2000) | US |
| 60/203,630 | 12 May 2000 (12.05.2000) | US |
| 60/210,741 | 12 June 2000 (12.06.2000) | US |
| 60/210,982 | 12 June 2000 (12.06.2000) | US |
| 60/226,760 | 21 August 2000 (21.08.2000) | US |
| 60/235,418 | 26 September 2000 (26.09,2000) | US |
| 60/235,779 | 26 September 2000 (26.09.2000) | US |
| 60/242,332 | 20 October 2000 (20.10.2000) | US |

(71) Applicant (for all designated States except US): ARENA PHARMACEUTICALS, INC. [US/US]; 6166 Nancy Ridge Drive, San Diego, CA 92121 (US).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): CHEN, Ruoping [CN/US]; 5296 Timber Branch Way, San Diego, CA 92130 (US). DANG, Huong, T. [US/US]: 5352 Oak Park Drive, San Diego, CA 92105 (US). LOWITZ, Kevin, P. [US/US]: 8031 Caminito de Pizza #C, San Diego, CA 82108 (US).
- (74) Agents: MILLER, Suzanne, E. et al.; Woodcock Washburn Kurtz Mackiewicz & Norris LLP. One Liberty Place -46th Floor, Philadelphia, PA 19103 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO. NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: ENDOGENOUS AND NON-ENDOGENOUS VERSIONS OF HUMAN G PROTEIN-COUPLED RECEPTORS

4000 P3 Accumulation (cpm/mg protein 3000 2000 1000 0 CMV RUP12

IP3 Assay in 293 Cells

(57) Abstract: The invention disclosed in this patent document relates to transmembrane receptors, more particularly to a human G protein-coupled receptor for which the endogenous ligand is unknown ("orphan GPCR receptors"), and most particularly to mutated (non-endogenous) versions of the human GPCRs for evidence of constitutive activity.

WO 01/36471 A

WO 01/36471 A3



Published:

with international search report

(88) Date of publication of the international search report: 3 January 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Intel anal Application No PCT/US 00/31509

| A. CLASSI IPC 7 | FICATION OF SUBJECT MATTER C07K14/705 C12N15/12 | | | | |
|--|---|---|-----------------------|--|--|
| According to International Patent Classification (IPC) or to both national classification and IPC | | | | | |
| B. FIELDS | SEARCHED | | | | |
| Minimum do IPC 7 | ocumentation searched (classification system followed by classifical $C07K$ | ion symbols) | | | |
| | tion searched other than minimum documentation to the extent that | | | | |
| - | ata base consulted during the international search (name of data baternal, EMBL, STRAND, WPI Data, EMB | · | | | |
| C. DOCUMI | ENTS CONSIDERED TO BE RELEVANT | | | | |
| Category * | Citation of document, with indication, where appropriate, of the re | levant naceanas | Relevant to claim No. | | |
| Category | Citation of document, with indication, where appropriate, or the re- | вечани раззаусэ | neterall to damino. | | |
| X | WO 98 46620 A (MILLENNIUM PHARM 22 October 1998 (1998-10-22) claim 1; figures 1A,,2A; example | · | 1-4 | | |
| X | US 5 891 720 A (WOOLF ELIZABETH / 6 April 1999 (1999-04-06) Sequence No. 2 abstract | | 1-4 | | |
| | | - /- - | | | |
| X Funt | ner documents are listed in the continuation of box C. | χ Patent family members are listed | in annex. | | |
| 8 Canadal and | topolice of cited documents : | | | | |
| "A" docume consid | tegories of cited documents: ent defining the general state of the art which is not ered to be of particular relevance | *T* later document published after the inter or pnortly date and not in conflict with cited to understand the principle or the invention | the application but | | |
| "E" earlier o | locument but published on or after the international ate | "X" document of particular relevance; the cl cannot be considered novel or cannot | | | |
| | nt which may throw doubts on priority claim(s) or is cried to establish the publication date of another | involve an inventive step when the doc | cument is taken alone | | |
| citation | which is cited to establish the publication date of another citation or other special reason (as specified) "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the | | | | |
| other r | "O" document referring to an oral disclosure, use, exhibition or document is combined with one or more other such document, such combination being obvious to a person skilled in the art. | | | | |
| *P* document published prior to the international filing date but later than the priority date claimed *&* document member of the same patent family | | | | | |
| | Date of the actual completion of the international search Date of mailing of the international search report | | | | |
| 2 | 8 August 2001 | 1 9. 09. 01 | | | |
| Name and n | naiting address of the ISA | Authorized officer | | | |
| | European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tol (231-70) 340-2040 Tv 31 651 epo pl | | | | |
| | Tel. (+31-70) 340-2040. Tx. 31 651 epo nl. Fax: (+31-70) 340-3016 | Meyer, W | | | |

6

Inter anal Application No
PCT/US 00/31509

| PCT/US 00/31509 | | | | |
|--|--|--|--|--|
| (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT alegory Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. | | | | |
| Relevant to claim No. | | | | |
| 1-4 | | | | |
| 1-4 | | | | |
| 1-4 | | | | |
| 1-4 | | | | |
| 1-4 | | | | |
| | | | | |

Inter onal Application No
PCT/US 00/31509

| 0.40 | NOW DOCUMENTS CONSIDERED TO BE BE! E!/ANT | <u> </u> |
|------------|---|-----------------------|
| Category ° | ation) DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| Category | Citation of document, with indication, where appropriate, of the relevant passages | Helevan to claim to. |
| P,X | DATABASE EMBL 'Online! Accession Nr. Q9NTTO, 1 October 2000 (2000-10-01) COLLIER R.: "DJ680N.3 (G-Protein Coupled Receptors) (Fragment)" XP002168498 abstract | 1-4 |
| Р,Х | WO 00 22131 A (ARENA PHARMACEUTICALS INC ;GORE MARTIN (US); LIAW CHEN W (US); LIN) 20 April 2000 (2000-04-20) the whole document | 1-4 |
| X | DATABASE EMBL 'Online! AC: AC008728, 4 August 1999 (1999-08-04) DOE JOINT GENOME INSTITUTE: "Sequencing of Human Chromosome 5" XP002175776 abstract | 5-8 |
| А | WO 98 29439 A (SULLIVAN KATHLEEN ; MERCK & CO INC (US); TAN CARINA (US)) 9 July 1998 (1998-07-09) page 57; figure 13; example 14 | 5-8 |
| E | WO 01 14577 A (SMITHKLINE BEECHAM PLC;SMITHKLINE BEECHAM CORP (US)) 1 March 2001 (2001-03-01) page 30-31; claims 1,2 | 5–8 |
| Ε | EP 1 090 989 A (PFIZER LTD ;PFIZER (US)) 11 April 2001 (2001-04-11) Seq. Id. No. 1, 2 | 5–8 |
| X | DATABASE EMBL 'Online! AC: AC008754, 4 August 1999 (1999-08-04) DOE JOINT GENOME INSTITUTE: "Homo sapiens chomosome 19 clone CTD-3023J11, complete sequence" XP002175778 abstract | 9–12 |
| х | DATABASE EMBL 'Online! AC: AQ532303, 18 May 1999 (1999-05-18) ZHAO S ET AL.: "Use of BAC End Sequences from Library RPCI-11 for Sequence-Ready Map Building" XP002175779 abstract | 9-12 |
| | -/ | |
| | | |

6

Inter anal Application No
PCT/US 00/31509

| C/Combine | POOL MENTS CONCIDEDED TO BE DELEVANT | FC1/03 00/31309 |
|------------|---|-----------------------|
| Category * | ation) DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| Ρ,Χ | DATABASE EMBL 'Online! AC: AB038237, 4 May 2000 (2000-05-04) OHONO ET AL.: "Homo spiens mRNA for G proteine-coupled receptor C5L2, complete cds" XP002175947 abstract | 9-12 |
| P,X | WO 00 14229 A (ONO MITSUHARU ;KANNO KIMIYOSHI (JP); ASAHI CHEMICAL IND (JP); ISHI) 16 March 2000 (2000-03-16) page 98 -page 101; claim 4 page 103; claim 5 | 9-12 |
| E | WO 01 36471 A (ARENA PHARMACEUTICALS INC; CHEN RUOPING (US); DANG HUONG T (US); L) 25 May 2001 (2001-05-25) Seq. Id. No. 3 (claims 5-8) Seq. Id. No. 5 (claims 9-12) Seq. Id. No. 7 (claims 13-16) Seq. Id. No. 9 (claims 17-20) Seq. Id. No. 11 (claims 17-20) Seq. Id. No. 13 (claims 25-28) Seq. Id. No. 21 (claims 41-44) Seq. Id. No. 19, 23 (claims 45-48) Seq. ID. No. 25 (claims 49-52) | 5-28, 41-52 |
| E | EP 1 094 076 A (PFIZER LTD ;PFIZER (US)) 25 April 2001 (2001-04-25) Seq. Id. No. 1 | 9-12 |
| E | WO 01 31014 A (UPJOHN CO ; VOGELI GABRIEL (US); WOOD LINDA S (US); MERCHANT KALPAN) 3 May 2001 (2001-05-03) Sequence No. 5 | 13–16 |
| X | PATENT ABSTRACTS OF JAPAN vol. 1999, no. 09, 30 July 1999 (1999-07-30) & JP 11 098988 A (SMITHKLINE BEECHAM CORP), 13 April 1999 (1999-04-13) abstract & DATABASE EMBL 'Online! AC: E31720; E75225, 22 February 2001 (2001-02-22) JEFFREY L.M.D.D. AND BERGSUMA W.S.H.H.: "cDNA clone HeoAd54 encoding human seven-pass transmembrane receptor" abstract | 13-16 |
| X | US 5 955 308 A (BERGSMA DERK J ET AL) 21 September 1999 (1999-09-21) Sequence 1 | 13–16 |
| | / | |

Inte. .onal Application No
PCT/US 00/31509

| onen ° | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|---------|---|--------------------------|
| egory ° | Challen of document, with incleasing | |
| | EP 0 892 051 A (SMITHKLINE BEECHAM CORP) 20 January 1999 (1999-01-20) page 20 -page 21; claim 11 | 13–16 |
| P | DATABASE EMBL 'Online! AC: AP000808, 3 December 1999 (1999-12-03) HATTORI M. ET AL.: "Homo sapiens 171,539 genomic of 11q13" XP002175780 abstract | 17-20 |
| | WO 99 32519 A (FORTIN YVES ; LEMBO PAOLA (CA); AHMAD SULTAN (CA); BANVILLE DENIS () 1 July 1999 (1999-07-01) | 37–40 |
| | page 48 -page 54; claims 16,20,21 | 17-20, 57-60 37-40 |
| | page 55 -page 56; claim 25 page 52 -page 54; claim 21 | 37–40 |
| | DATABASE EMBL 'Online! AC011780 , 18 October 1999 (1999-10-18) BIRREN B., LINTON L., NUSBAUM C., LANDER E.: "Homo sapiens clone RP11-15H8, 31 unordered pieces." XP002175781 abstract | 21-24 |
| | JP 08 245697 A (TAKEDA CHEM IND LTD) 24 September 1996 (1996-09-24) claim 4; figures 1,2 | 21-24 |
| | WO 96 05302 A (FUJII RYO ;HOSOYA MASAKI (JP); OHGI KAZUHIRO (JP); FUKUSUMI SHOJI) 22 February 1996 (1996-02-22) page 263 -page 264; example 16 | 21-24 |
| P,X | DATABASE EMBL 'Online! AC: AL355310 , 5 May 2000 (2000-05-05) WALLIS, J: "Human DNA sequence from clone RP5-1160K1" XP002175782 abstract | 21-24 |
| x | DATABASE EMBL 'Online! AC: AQ001459, ADAMS M.D. ET AL.: "CIT-HSP-2286K19.TF CIT-HSP Homo sapiens genomic clone 2286K19, genomic survey sequence" XP002175783 abstract | 25-28 |
| A | EP 0 878 542 A (SMITHKLINE BEECHAM CORP) 18 November 1998 (1998-11-18) page 18 -page 19; claim 1 | 25-28 |
| | -/ | |

6

Inter onal Application No PCT/US 00/31509

| C.(Continua | ation) DOCUMENTS CONSIDERED TO BE RELEVANT | PCT/US 00/31509 |
|-------------|---|-----------------------|
| Category ° | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| Ρ,Χ | HEISE CHRISTOPHER E ET AL: "Characterization of the human cysteinyl leukotriene 2 receptor." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 275, no. 39, 29 September 2000 (2000-09-29), pages 30531-30536, XP002175775 ISSN: 0021-9258 the whole document & DATABASE EMBL 'Online! AC: AF254664, HEISE C.E. ET AL.: "Homo sapiens cysteinyl | Relevant to claim No. |
| | leukotriene receptor CYSLT2 gene, complete cds." abstract US 5 861 309 A (WEINSHANK RICHARD L ET AL) 19 January 1999 (1999-01-19) Sequence 1 | 29-32 |
| - | WO 01 09184 A (DELEERSNIJDER WILLY; NYS GUY (BE); ZHANG FAN (BE); SOLVAY PHARMACE) 8 February 2001 (2001-02-08) Sequence 1 page 6 -page 7; claims 1,15 | 29-32 |
| ,х | DATABASE EMBL 'Online! ACC16468, 1 December 1999 (1999-12-01) BIRREN B. ET AL.: "Homo sapiens clone RP11-14N15" XP002175784 abstract | 29-32 |
| | WO 99 48921 A (ORGANON NV ;SPEK PETRUS JOHANNES V D (NL); UNIV LELAND STANFORD JU) 30 September 1999 (1999-09-30) claims 2,4; figure 4 | 33–36 |
| , х | DATABASE EMBL 'Online! AL136106, 7 January 2000 (2000-01-07) BURTON J: "Human DNA sequence from clone RP11-15909" XP002175785 abstract | 33-36 |
| 30.7 | DATABASE EMBL 'Online! AC: AC008547, OE JOINT GENOME INSTITUTE STANFORD HUMAN GENOME CENTER.: "Homo sapiens chromosome 5 clone CTC-502M5, complete sequence." XP002175786 abstract | 41-44 |

Inte onal Application No
PCT/US 00/31509

| Contine | ation) DOCUMENTS CONSIDERED TO BE RELEVANT | |
|-----------|--|-----------------------|
| ategory ° | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| | WO 99 06552 A (GENSET SA ;LACROIX BRUNO (FR); DUCLERT AYMERIC (FR); DUMAS MILNE E) 11 February 1999 (1999-02-11) SEQ ID NO: 95 | 41-44 |
| | EP 0 612 845 A (AMERICAN CYANAMID CO) 31 August 1994 (1994-08-31) claim 2; figure 9 | 49-52 |
| | DATABASE EMBL 'Online! AC: AL065769, 29 May 1999 (1999-05-29) GEMPSCPÜE: "Drosophila melanogaster genome survey sequence TET3 end of BAC # BACR08K10 of RPCI-98 library from Drosophila melanogaster (fruit fly)" XP002175910 abstract | 49-52 |
| Ρ,Χ | DATABASE EMBL 'Online! AC: Al161458, 16 April 2000 (2000-04-16) BURTON J. ET AL.: "Human DNA sequence from clone RP11-163L4" XP002175911 abstract | 49–52 |
| A | BOYER JOSE L ET AL: "Molecular cloning and expression of an avian G protein-coupled P2Y receptor." MOLECULAR PHARMACOLOGY, vol. 52, no. 6, December 1997 (1997-12), pages 928-934, XP002175907 ISSN: 0026-895X the whole document | 53-56 |
| Ρ,Χ | DATABASE EMBL 'Online! AC: AC026756, 24 April 2000 (2000-04-24) ABOLA A.P. ET AL.: "omo sapiens chromosome 13 clone RP11-286P8, complete sequence" XP002175912 abstract | 53–56 |
| P,X | DATABASE EMBL 'Online! AC ACO27026, 27 April 2000 (2000-04-27) BIRREN B. ET AL.: "Homo sapiens chromosome 11, clone RP11-589F4" XP002175913 abstract | 57-60 |
| E | WO 01 16159 A (SMITHKLINE BEECHAM CORP) 8 March 2001 (2001-03-08) page 30; claim 1 | 57-60 |
|] | | |

6

Inte onal Application No
PCT/US 00/31509

| Category " | Citation of document, with indication, where appropriate, of the relevant passages | • |
|------------|--|-----------------------|
| | appropriate, or the relevant passages | Relevant to claim No. |
| х | DATABASE EMBL 'Online! AC: AC007104, 23 April 1999 (1999-04-23) STONE ET AL.: "Homo sapiens chomosome 4, 16 unordered pieces" XP002175914 abstract | 61-64 |
| E | WO 01 12673 A (MERCK PATENT GMBH; DUECKER KLAUS (DE)) 22 February 2001 (2001-02-22) Sequence 1, 2 page 39; claim 3 | 61-64 |
| X | JP 11 032770 A (ASAHI CHEM IND CO LTD) 9 February 1999 (1999-02-09) page 19; claim 7 | 65-68 |
| X | WO 98 56820 A (ELSHOURBAGY NABIL A ;SMITHKLINE BEECHAM CORP (US); LI XIAOTONG (US) 17 December 1998 (1998-12-17) page 30 -page 31; claims 1,2 | 69-72 |
| ١ | the whole document | 45-48 |
| (| DATABASE EMBL 'Online! AC: AC010984, 29 September 1999 (1999-09-29) | 73-76 |
| | WATERSON R.H.: "Homo sapiens chromosome 2 clone RP11-510C1" XP002175915 abstract | |
| | DATABASE EMBL 'Online! AC 008892, 15 July 1998 (1998-07-15) WEINSHANK R. H.: "5-Hydroxytryptamine 1B Receptor(-HT-1B) (Serotonin Receptor)" XP002175948 abstract | 73-76 |
| | MAHAIRAS GREGORY G ET AL: "Sequence-tagged connectors: A sequence approach to mapping and scanning the human genome." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES, vol. 96, no. 17, 17 August 1999 (1999-08-17), pages 9739-9744, XPO02175909 | 77-80 |
| | Aug. 17, 1999 ISSN: 0027-8424 the whole document | |
| | WO 01 07606 A (SMITHKLINE BEECHAM PLC) 1 February 2001 (2001-02-01) Sequence 2 page 31; claim 4 | 77-80 |
| | -/ | |
| 1 | , | |

Inte ional Application No
PCT/US 00/31509

| C (Centime | ation) DOCUMENTS CONSIDERED TO BE RELEVANT | FC1703 00/31309 | | |
|------------|--|-----------------|----------------------|--|
| | | Relevant to | elevant to claim No. | |
| | or a contract of the following states | , cavani i | | |
| P,X | Citation of document, with indication, where appropriate, of the relevant passages WO 00 49046 A (TERAO YASUKO ; WATANABE TAKUYA (JP); SHINTANI YASUSHI (JP); TAKEDA) 24 August 2000 (2000-08-24) claim 5; figure 1 | | -80 | |
| | | | | |
| | | | | |
| | | | | |

Inconational application No. PCT/US 00/31509

| Box I | Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet) | | | | | | |
|---|--|--|--|--|--|--|--|
| This Inte | ernational Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons: | | | | | | |
| 1. | Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely: | | | | | | |
| 2. | Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically: | | | | | | |
| 3. | Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a). | | | | | | |
| Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet) | | | | | | | |
| This Inte | ernational Searching Authority found multiple inventions in this international application, as follows: | | | | | | |
| | see additional sheet | | | | | | |
| | • - | | | | | | |
| 1. X | As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims. | | | | | | |
| 2. | As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee. | | | | | | |
| 3. | As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.: | | | | | | |
| 4. | No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: | | | | | | |
| Remark | on Protest The additional search fees were accompanied by the applicant's protest. X No protest accompanied the payment of additional search fees. | | | | | | |

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1998)

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-4

G protein-coupled receptor as characterized by SEQ.ID.2, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.1, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

2. Claims: 5-8

G protein-coupled receptor as characterized by SEQ.ID.4, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.3, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

3. Claims: 9-12

G protein-coupled receptor as characterized by SEQ.ID.6, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.5, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

4. Claims: 13-16

G protein-coupled receptor as characterized by SEQ.ID.8, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.7, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

5. Claims: 17-20

G protein-coupled receptor as characterized by SEQ.ID.10, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.9, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

6. Claims: 21-24

G protein-coupled receptor as characterized by SEQ.ID.12, its non-endogenous, constitutively activated version SEQ ID.84, a cDNA encoding said receptor as characterized by SEQ.ID.11, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

7. Claims: 25-28

G protein-coupled receptor as characterized by SEQ.ID.14, its non-endogenous, constitutively activated version SEQ.ID.88, a cDNA encoding said receptor as characterized by SEQ.ID.13, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

8. Claims: 29-32

G protein-coupled receptor as characterized by SEQ.ID.16, its non-endogenous, constitutively activated version SEQ.ID.92, a cDNA encoding said receptor as characterized by SEQ.ID.15, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

9. Claims: 33-36

G protein-coupled receptor as characterized by SEQ.ID.18, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.17, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

10. Claims: 37-40

G protein-coupled receptor as characterized by SEQ.ID.20, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.19, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

11. Claims: 41-44

G protein-coupled receptor as characterized by SEQ.ID.22, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.21, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

12. Claims: 45-48

G protein-coupled receptor as characterized by SEQ.ID.24, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.23, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

13. Claims: 49-52

G protein-coupled receptor as characterized by SEQ.ID.26, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.25, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

14. Claims: 53-56

G protein-coupled receptor as characterized by SEQ.ID.28, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.27, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

15. Claims: 57-60

G protein-coupled receptor as characterized by SEQ.ID.30, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.29, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

16. Claims: 61-64

G protein-coupled receptor as characterized by SEQ.ID.32, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.96, a plasmid comprising said SEQ.ID 95, and a host cell comprising said plasmid.

17. Claims: 65-68

G protein-coupled receptor as characterized by SEQ.ID.34, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.33, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

18. Claims: 69-72

G protein-coupled receptor as characterized by SEQ.ID.36, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.35, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

19. Claims: 73-76

G protein-coupled receptor as characterized by SEQ.ID.38, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.37, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

20. Claims: 77-80

G protein-coupled receptor as characterized by SEQ.ID.40, its non-endogenous, constitutively activated version, a cDNA encoding said receptor as characterized by SEQ.ID.39, a plasmid comprising said cDNA, and a host cell comprising said plasmid.

Information on patent family members

Inte onal Application No PCT/US 00/31509

| | | | PCI/ | US 00/31509 |
|---|---|------------------|------------------------------|--------------------------|
| Patent document cited in search repo | | Publication date | Patent family member(s) | Publication date |
| WO 9846620 | Α | 22-10-1998 | US 5891720 A AU 6973698 A | 06-04-1999 11-11-1998 |
| | | | EP 1007536 A | 14-06-2000 |
| US 5891720 | Α | 06-04-1999 | AU 6973698 A EP 1007536 A | 11-11-1998 14-06-2000 |
| | | | WO 9846620 A | 22-10-1998 |
| WO 9721731 | Α | 19-06-1997 | US 5750353 A | 12-05-1998 |
| | | | AU 715611 B AU 1334397 A | 03-02-2000 03-07-1997 |
| | | | CA 2239293 A | 19-06-1997 |
| | | | EP 0869975 A | 14-10-1998 |
| | | | JP 2000510324 T | 15-08-2000 |
| WO 0022131 | Α | 20-04-2000 | AU 6299199 A | 01-05-2000 |
| | | | AU 6430799 A EP 1121431 A | 01-05-2000 |
| | | | WO 0021987 A | 08-08-2001 20-04-2000 |
| | | | WO 0022129 A | 20-04-2000 |
| | | | AU 3790400 A | 13-06-2000 |
| | | | WO 0031258 A | 02-06-2000 |
| WO 9829439 | Α | 09-07-1998 | EP 0948529 A | 13-10-1999 |
| | | | EP 0948532 A EP 0960125 A | 13-10-1999 |
| | | | EP 0960125 A WO 9829440 A | 01-12-1999 09-07-1998 |
| | | | WO 9829441 A | 09-07-1998 |
| WO 0114577 | Α | 01-03-2001 | NONE | |
| EP 1090989 | A | 11-04-2001 | NONE | |
| WO 0014229 | Α | 16-03-2000 | AU 5449099 A | 27-03-2000 |
| | | | EP 1111049 A | 27-06-2001 |
| WO 0136471 | Α | 25-05-2001 | NONE | |
| EP 1094076 | Α | 25-04-2001 | NONE | |
| WO 0131014 | Α | 03-05-2001 | NONE | |
| JP 11098988 | Α | 13-04-1999 | US 5955308 A | 21-09-1999 |
| | | | EP 0892051 A | 20-01-1999 |
| JS 5955308 | Α | 21-09-1999 | EP 0892051 A | 20-01-1999 |
| | | | JP 11098988 A | 13-04-1999 |
| EP 0892051 | Α | 20-01-1999 | US 5955308 A | 21-09-1999 |
| | | | JP 11098988 A | 13-04-1999 |
| √O 9932519 | Α | 01-07-1999 | AU 1990499 A | 12-07-1999 |
| | | | BR 9814335 A | 10-10-2000 |
| | | | CN 1284966 T EP 1051434 A | 21-02-2001 |
| | | | NO 20003221 A | 15-11-2000 10-08-2000 |
| | | | PL 341524 A | 23-04-2001 |
| | | | TR 200001861 T | |

Information on patent family members

Int ional Application No PCT/US 00/31509

| | | T | | <u></u> | |
|---------------------------------------|-------|---------------------|-------------|--------------------------|--------------------------|
| Patent document cited in search repor | t | Publication date | | Patent family member(s) | Publication date |
| JP 08245697 | Α | 24-09-1996 | NONE | • | |
| WO 9605302 | Α | 22-02-1996 | AU | 4426296 A | 07-03-1996 |
| | , , | | CA | 2195768 A | 22-02-1996 |
| | | | EP | 0804575 A | 05-11-1997 |
| | | | JР | 9000268 A | 07-01-1997 |
| | | | US | 6114139 A | 05-09-2000 |
| EP 0878542 | Α | 18-11-1998 | CA | 2231740 A | 22-10-1998 |
| | | | CA | 2234399 A | 22-10-1998 |
| | | | EP | 0874047 A | 28-10-1998 |
| | | | JP | 11032784 A | 09-02-1999 |
| | · | | US | 6200775 B | 13-03-2001 |
| US 5861309 | Α | 19-01-1999 | AU | 718197 B | 06-04-2000 |
| | | • | AU | 3420797 A | 29-01-1998 |
| | | | AU | 677968 B | 15-05-1997 |
| | | | AU | 5165693 A | 26-04-1994 |
| | | | CA DE | 2145182 A | 14-04-1994 |
| | | | EP | 663014 T 1063291 A | 10-10-1996 27-12-2000 |
| | | | EP | 1063291 A 1063292 A | 27-12-2000 27-12-2000 |
| | | | EP | 0663014 A | 19-07-1995 |
| | | | ES | 2085247 T | 01-06-1996 |
| | | | GR | 95300067 T | 31-01-1996 |
| | | | JP | 8505044 T | 04-06-1996 |
| | | | WO | 9408040 A | 14-04-1994 |
| | | | ÜS | 6083705 A | 04-07-2000 |
| | | | ÜŠ | 5556753 A | 17-09-1996 |
| | | | US | 5714381 A | 03-02-1998 |
| | | | us | 6156518 A | 05-12-2000 |
| WO 0109184 | Α | 08-02-2001 | AU | 5985800 A | 19-02-2001 |
| WO 9948921 | Α | 30-09-1999 | EP | 1066324 A | 10-01-2001 |
| WO 9906552 | Α | 11-02-1999 | US | 6222029 B | 24-04-2001 |
| | | | AU | 8555598 A | 22-02-1999 |
| | | | EP | 1000150 A | 17-05-2000 |
| EP 0612845 | A | 31-08-1994 | AU | 5640494 A | 01-09-1994 |
| | | | JP | 7095889 A | 11-04-1995 |
| | | | US | 6258556 B | 10-07-2001 |
| | | | US | 6225080 B | 01-05-2001 |
| WO 0116159 | A | 08-03-2001 | NONE | | |
| WO 0112673 | Α | 22-02-2001 | NONE | | |
| JP 11032770 | A | 09-02-1999 | NONE | | |
| WO 9856820 | A | 17-12-1998 | AU EP | 7966098 A 1007563 A | 30-12-1998 14-06-2000 |
| WO 0107606 | Α | 01-02-2001 | NONE | | |
| WO 0049046 | Α | 24-08-2000 | AU JP 20 | 2573900 A 001017186 A | 04-09-2000 23-01-2001 |

Form PCT/ISA/210 (patent family annex) (July 1992)